

Documents

Jinap, S.^{a b}, Hasnol, N.D.S.^a, Sanny, M.^a, Jahurul, M.H.A.^c

Effect of organic acid ingredients in marinades containing different types of sugar on the formation of heterocyclic amines in grilled chicken

(2018) *Food Control*, 84, pp. 478-484. Cited 33 times.

DOI: 10.1016/j.foodcont.2017.08.025

^a Department of Food Science, Faculty of Food Science and Technology, Universiti Putra Malaysia, Serdang, Selangor 43400 UPM, Malaysia

^b Food Safety and Food Integrity (FOSFI), Institute of Tropical Agriculture and Food Security, Universiti Putra Malaysia, Serdang, Selangor 43400 UPM, Malaysia

^c Faculty of Food Science and Nutrition, Universiti Malaysia Sabah, Kota Kinabalu, Sabah 884000, Malaysia

Abstract

The aim of the study was to determine the use of alternative organic acids in formulating marinade ingredients to reduce heterocyclic amines (HCA) in grilled chicken (satay). Samples were marinated with table sugar, brown sugar, and honey with the addition of tamarind, lemon, lime, and calamansi for 24 h at 4 °C. The pH readings before and after marinating were measured. HCA concentrations before and after grilling were quantified. There was a significant difference ($p < 0.05$) in the combined HCAs among the control and marinated grilled chickens. Using lemon in marinades containing table sugar, concentrations of DiMeIQx were significantly reduced ($p < 0.006$) from 16.5 ng/g (low) to 8.30 ng/g (high) concentrations of organic acid ingredients. The mean pH of the treated samples was significantly lower ($p < 0.05$) than in the control samples. Calamansi was found to reduce HCAs in marinades containing table sugar and brown sugar, whereas tamarind in marinades containing honey. © 2017

Author Keywords

Calamansi; Heterocyclic amines; Lemon; Marinated grilled chicken; Tamarind

References

- Ables, P.A.F.R.H.
Enzymatic reaction mechanism
(2006), Oxford University Press
- Alaejos, M.S., Afonso, A.M.
Factors that affect the content of heterocyclic aromatic amines in foods
(2011) *Comprehensive Reviews in Food Science and Food Safety*, 10 (2), pp. 52-108.
- Anyasi, T.A., Jideani, A.I.O., Mchau, G.R.A.
Effect of organic acid pretreatment on some physical, functional and antioxidant properties of flour obtained from three unripe banana cultivars
(2015) *Food Chemistry*, 172, pp. 515-522.
- Barceló-Barrachina, E., Moyano, E., Galceran, M.T.
Determination of heterocyclic amines by liquid chromatography-quadrupole time-of-flight mass spectrometry
(2004) *Journal of Chromatography A*, 1054 (1-2), pp. 409-418.
- Berge, P., Ertbjerg, P., Larsen, L.M., Astruc, T., Vignon, X., Møller, A.J.
Tenderization of beef by lactic acid injected at different times post mortem
(2001) *Meat Science*, 57, pp. 347-357.
- Burke, R.M., Monahan, F.J.
Effect of breed-type on performance and carcass traits of intensively managed hair sheep
(2003) *Meat Science*, 63 (3), pp. 309-315.

- Cheng, K.W., Chen, F., Wang, M.
Inhibitory activities of dietary phenolic compounds on heterocyclic amine formation in both chemical model system and beef patties
(2007) *Molecular Nutrition and Food Research*, 51 (8), pp. 969-976.
- Chiu, C.P., Yang, D.Y., Chen, B.H.
Formation of heterocyclic amines in cooked chicken legs
(1998) *Journal of Food Protection*, 61 (6), pp. 712-719.
- Darias-Martín, J., Socas-Hernández, A., Díaz-Romero, C., Díaz-Díaz, E.
Comparative study of methods for determination of titrable acidity in wine
(2003) *Journal of Food Composition and Analysis*, 16, pp. 555-562.
- Emamgholizadeh, F.
Effects of marinades on the formation of heterocyclic amines in grilled beef steaks
(1993), Master of Science. M.D. Azad Tehran University
- Farhadian, A., Jinap, S., Hanifah, H.N., Zaidul, I.S.
Effects of meat preheating and wrapping on the levels of polycyclic aromatic hydrocarbons in charcoal-grilled meat
(2011) *Food Chemistry*, 124 (1), pp. 141-146.
- Felton, J.S., Fultz, E., Dolbeare, F.A., Knize, M.G.
Reduction of heterocyclic amine mutagens/carcinogens in fried beef patties by microwave treatment
(1994) *Food and Chemical Toxicology*, 32 (10), pp. 897-903.
- Giarratana, F., Muscolino, D., Beninati, C., Ziino, G., Giuffrida, A., Panebianco, A.
Activity of R(+) limonene on the maximum growth rate of fish spoilage organisms and related effects on shelf-life prolongation of fresh gilthead sea bream fillets
(2016) *International Journal of Food Microbiology*, 237, pp. 109-113.
- Giarratana, F., Muscolino, D., Panebianco, F., Patania, A., Benianti, C., Ziino, G.
Activity of R(+) limonene against anisakis larvae
(2015) *Italian Journal of Food Safety*, 4, pp. 209-211.
- Gibis, M.
Effect of oil marinades with garlic, onion, and lemon juice on the formation of heterocyclic aromatic amines in fried beef patties
(2007) *Journal of Agriculture and Food Chemistry*, 55.
- Gross, G.A., Grüter, A.
Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products
(1992) *Journal of Chromatography A*, 592 (1-2), pp. 271-278.
- Hasnol, N.D.S., Jinap, S., Sanny, M.
Effect of different types of sugars in a marinating formulation on the formation of heterocyclic amines in grilled chicken
(2014) *Food Chemistry*, 145, pp. 514-521.
- Hiu, Y.H., Barta, J., Cano, P.M.
Handbook of the fruits and fruit processing: Science and Technology
(2006), John Wiley & Sons Hoboken
- Hwang, D.K., Ngadi, M.
Kinetics of heterocyclic amines formation in meat emulsion at different fat contents
(2002) *LWT - Food Science and Technology*, 35 (7), pp. 600-606.
- IARC
Some naturally occurring Substances: Food items and constituents, heterocyclic

aromatic amines and mycotoxins

(1993) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, 56, pp. 165-195.

- Jang, A., Liu, X.D., Shin, M.H., Lee, B.D., Lee, S.K., Lee, J.H.
Antioxidative potential of raw breast meat from broiler chicks fed a dietary medicinal herb extract mix
(2008) *Poultry Science*, 87 (11), pp. 2382-2389.
- Jinap, S., Iqbal, S.Z., Selvam, R.M.P.
Effect of selected local spices marinades on the reduction of heterocyclic amines in grilled beef (satay)
(2015) *LWT - Food Science and Technology*, 63, pp. 919-926.
- Jinap, S., Iqbal, S.Z., Talib, N.H., Hasnol, N.D.S.
Heterocyclic aromatic amines in deep fried lamb meat: The influence of spices marination and sensory quality
(2016) *Journal of Food Science and Technology*, 53, pp. 1411-1417.
- Koohmaraie, M., Kent, M.P., Shackelford, S.D., Veiseth, E., Wheeler, T.L.
Meat tenderness and muscle growth: Is there any relationship?
(2003) *Meat Science*, 62 (3), pp. 345-352.
- Lu, F., Kuhnle, G.K., Cheng, Q.
Heterocyclic amines and polycyclic aromatic hydrocarbons in commercial ready-to-eat meat products on UK market
(2017) *Food Control*, 73, pp. 306-315.
- Messner, C., Murkovic, M.
Evaluation of a new model system for studying the formation of heterocyclic amines
(2004) *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*, 802 (1), pp. 19-26.
- Oz, F., Cakmak, I.H.
The effects of conjugated linoleic acid usage in meatball production on the formation of heterocyclic aromatic amines
(2016) *LWT - Food Science and Technology*, 65, pp. 1031-1037.
- Oz, F., Kizil, M., Zaman, A., Turhan, S.
The effects of direct addition of low and medium molecular weight chitosan on the formation of heterocyclic aromatic amines in beef chop
(2016) *LWT - Food Science and Technology*, 65, pp. 861-867.
- Salmon, Knize, M.G., Felton, J.S.
Effects of marinating on heterocyclic amine carcinogen formation in grilled chicken
(1997) *Food and Chemical Toxicology*, 35, pp. 433-441.
- Shin, H.S., Rodgers, W.J., Gomaa, E.A., Strasburg, G.M., Gray, J.I.
Inhibition of heterocyclic aromatic amine formation in fried ground beef patties by garlic and selected garlic-related sulfur compounds
(2002) *Journal of Food Protection*, 65.
- Skog, K.I., Johansson, M.A.E., Jägerstad, M.I.
Carcinogenic heterocyclic amines in model systems and cooked foods: A review on formation occurrence and intake
(1998) *Food and Chemical Toxicology*, 36, pp. 879-896.
- Tan, S.M., Lee, S.M., Dykes, G.A.
Buffering effect of chicken skin and meat protects *Salmonella enterica* strains against hydrochloric acid but not organic acid treatment
(2014) *Food Control*, 42, pp. 329-334.

- Turesky, R.J.
Formation and biochemistry of carcinogenic heterocyclic aromatic amines in cooked meats
(2007) *Toxicology Letters*, 168, pp. 219-227.
- Viegas, O., Novo, P., Pinto, E., Pinho, O., Ferreira, I.M.P.L.V.O.
Effect of charcoal types and grilling conditions on formation of heterocyclic aromatic amines (HAs) and polycyclic aromatic hydrocarbons (PAHs) in grilled muscle foods
(2012) *Food and Chemical Toxicology*, 50, pp. 2128-2134.

Correspondence Address

Jinap S.; Department of Food Science, Malaysia; email: jinap@upm.edu.my

Publisher: Elsevier Ltd

ISSN: 09567135

CODEN: FOOCE

Language of Original Document: English

Abbreviated Source Title: Food Control

2-s2.0-85037650789

Document Type: Article

Publication Stage: Final

Source: Scopus

ELSEVIER

Copyright © 2024 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

 RELX Group™