

Documents

Ullah, S.^a, Huyop, F.^{a h}, Wahab, R.A.^b, Huda, N.^c, Oyewusi, H.A.^{a g}, Sujana, I.G.A.^h, Saloko, S.^d, Andriani, A.A.S.P.R.^e, Mohamad, M.A.N.^f, Abdul Hamid, A.A.^f, Mohd Nasir, M.H.^f, Antara, N.S.^h, Gunam, I.B.W.^h

The first ITS1 profiling of honey samples from the Southeast Asian region Lombok, Bali and Banggi Island
(2024) *Scientific Reports*, 14 (1), art. no. 14122. .

DOI: 10.1038/s41598-024-64838-3

^a Department of Biosciences, Faculty of Science, Universiti Teknologi Malaysia, Johor Bahru, 81310, Malaysia

^b Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia, Johor Bahru, 81310, Malaysia

^c Faculty of Sustainable Agriculture, Universiti Malaysia Sabah, Sabah, Sandakan, 90509, Malaysia

^d Faculty of Food Technology and Agro Industry, University of Mataram, Nusa Tenggara Barat, Mataram, 83126, Indonesia

^e Faculty of Agriculture, Warmadewa University, Denpasar, Indonesia

^f Research Unit for Bioinformatics and Computational Biology (RUBIC), Kulliyah of Science, International Islamic University Malaysia, Bandar Indera Mahkota, Pahang, Kuantan, 25200, Malaysia

^g Biochemistry Unit, Department of Science Technology, The Federal Polytechnic, Ekiti State, Ado Ekiti, Nigeria

^h Bioindustry Laboratory, Department of Agro-Industrial Technology, Udayana University, Denpasar, Indonesia

Abstract

Southern Asian flowers offer honeybees a diversity of nectar. Based on its geographical origin, honey quality varies. Traditional methods are less authentic than DNA-based identification. The origin of honey is determined by pollen, polyphenolic, and macro-microorganisms. In this study, amplicon sequencing targets macro-microorganisms in eDNA using the ITS1 region to explore honey's geographical location and authentication. The variety of honey samples was investigated using ITS1 with Illumina sequencing. For all four honey samples, raw sequence reads showed 979,380 raw ITS1 amplicon reads and 375 ASVs up to the phylum level. The highest total number of 202 ASVs up to phylum level identified Bali honey with 211,189 reads, followed by Banggi honey with 309,207 a total number of 111 ASVs, and Lombok represents only 63 ASVs up to phylum level with several read 458,984. Based on Shannon and Chao1, honey samples from Bali (B2) and (B3) exhibited higher diversity than honey from Lombok (B1) and green honey from Sabah (B4), while the Simpson index showed that Banggi honey (B4) had higher diversity. Honey samples had significant variance in mycobiome taxonomic composition and abundance. *Zygosaccharomyces* and *Aspergillus* were the main genera found in Lombok honey, with percentages of 68.81% and 29.76% respectively. Bali honey samples (B2 and B3) were identified as having a significant amount of the genus *Aureobasidium*, accounting for 40.81% and 25% of the readings, respectively. The microbiome composition of Banggi honey (B4) showed a high presence of *Zygosaccharomyces* 45.17% and *Aureobasidium* 35.24%. The ITS1 analysis effectively distinguishes between honey samples of different origins and its potential as a discriminatory tool for honey origin and authentication purposes. © The Author(s) 2024.

Author Keywords

ASVs; Bali; Banggi; Illumina sequencing; ITS1; Lombok

Correspondence Address

Huyop F.; Department of Biosciences, Malaysia; email: fahrul@utm.my
Gunam I.B.W.; Bioindustry Laboratory, Indonesia; email: ibwgunam@unud.ac.id

Publisher: Nature Research

ISSN: 20452322

Language of Original Document: English

Abbreviated Source Title: Sci. Rep.

2-s2.0-85196283122

Document Type: Article

Publication Stage: Final

Source: Scopus

ELSEVIER

Copyright © 2024 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

 **RELX** Group™