





























- [22] Sekoai, P. T., Awosusi, A. A., Yoro, K. O., Singo, M., Oloye, O., Ayeni, A. O., ... Daramola, M. O. (2017). *Microbial cell immobilization in biohydrogen production: a short overview. Critical Reviews in Biotechnology*, 38(2), 157–171. doi:10.1080/07388551.2017.1312274
- [23] Seelert T, Ghosh D, Yargeau V. Improving biohydrogen production using *Clostridium beijerinckii* immobilized with magnetite nanoparticles. *Appl Microbiol Biotechnol*. 2015;99:4107–4116.
- [24] Jamali, N. S., Rashidi, N. F. D., Jahim, J. M., Thong, S. O., Jehlee, A., & Engliman, N. S. (2019). Thermophilic biohydrogen production from palm oil mill effluent: Effect of immobilized cells on granular activated carbon in fluidized bed reactor. *Food and Bioproducts Processing*. doi:10.1016/j.fbp.2019.07.012
- [25] Venkata Mohan, S., Vijaya Bhaskar, Y., Sarma PN., (2007). Biohydrogen production from chemical wastewater treatment in biofilm configured reactor operated in periodic discontinuous batch mode by selectively enriched anaerobic mixed consortia. *Water Res*;41: 2652–64.
- [26] Ginkel, v. S., Sung S, Lay JJ., 2001. Biohydrogen production as a function of pH and substrate concentration. *Environ Sci Technol*;35:4726–30.
- [27] Zagrodnik R, Laniecki M. The role of pH control on biohydrogen production by single stage hybrid dark- and photo-fermentation. *Bioresour Technol*. 2015 Oct;194:187-95. doi: 10.1016/j.biortech.2015.07.028. Epub 2015 Jul 14. PMID: 26196419.
- [28] Tiago Borges Ferreira, Gabriel Catucci Rego, Lucas Rodrigues Ramos, Camila Aparecida de Menezes, Laís Américo Soares, Isabel Kimiko Sakamoto, Maria Bernadete Amâncio Varesche, Edson Luiz Silva, HRT control as a strategy to enhance continuous hydrogen production from sugarcane juice under mesophilic and thermophilic conditions in AFBRs, *Int. J. Hydrogen Energy* 44 (36) (2019) 19719–19729, <https://doi.org/10.1016/j.ijhydene.2019.06.050>.
- [29] Su, C., Liu, Y., Yang, X., & Li, H. (2020). *Effect of hydraulic retention time on biohydrogen production from glucose in an internal circulation reactor. Energy & Fuels*. doi:10.1021/acs.energyfuels.9b033
- [30] Kumar G, Sen B, Sivagurunathan P, Lin C-Y. (2016). High rate hydrogen fermentation of cello-lignin fraction in de-oiled jatropha waste using hybrid immobilized cell system. *Fuel*;182:131e40
- [31] Albert, J., Wölfel, R., Bösmann, A., Wasserscheid, P., 2012. Selective oxidation of complex, water-insoluble biomass to formic acid using additives as reaction accelerators. *Energy & Environmental Science*. 5 (7): 7956.
- [32] Thauer, R.K., K. Jungermann & K. Decker 1977. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriological Reviews* 41(1): 100-180
- [33] Sivagurunathan, P., Sen, B., Chiu-Yue, L., 2014. Overcoming propionic acid inhibition of hydrogen fermentation by temperature shift strategy. *International Journal of Hydrogen Energy*. 39: 19232-19241.
- [34] Suksong, W., P. Kongjan, P. Prasertsan, T. Imai & O. Sompong 2016. Optimization and microbial community analysis for production of biogas from solid waste residues of palm oil mill industry by solid-state anaerobic digestion. *Bioresource technology* 214: 166-174.
- [35] Lutpi NA, Jahim JM, Mumtaz T, Abdul PM, Nor MTM. Physicochemical characteristics of attached biofilm on granular activated carbon for thermophilic biohydrogen production. *RSC Adv* 2015;5:19382e92.
- [36] Chu C-Y, Wu S-Y, Shen Y-C. Biohydrogen production performance in a draft tube bioreactor with immobilized cell. *Int J Hydrogen Energy* 2012;37:15658e65.
- [37] Zhang S, Lee Y, Kim T-H, Hwang S-J. Effects of OLRs and HRTs on hydrogen production from high salinity substrate by halophilic hydrogen producing bacterium (HHPB). *Bioresour Technol* 2013;141:227e32.