

Documents

Nasuruddin, M.H.^a, Che-Ahmad, A.^a, Ayeop, M.A.S.^a, Najmi, N.M.F.N.M.^b, Ahmad, M.W.^c

Pedobarography Study among Malay Population in Kuantan, Malaysia: A Pilot Study
(2022) *IUM Medical Journal Malaysia*, 21 (3), pp. 119-122.

DOI: 10.31436/imjm.v21i3.1755

^a Department of Orthopaedics, Traumatology, and Rehabilitation, International Islamic University, Malaysia

^b Department of Orthopaedic, Pahang Medical Centre, Malaysia

^c Department of Orthopaedic, University Malaysia Sabah, Malaysia

Abstract

INTRODUCTION: Pedobarography has been widely used in developed countries for few decades. However, in Malaysia it is still in its infancy. Studies showed that the normal values of pedobarography vary between races. To the best of our knowledge there is no standard value available for Malaysian or Southeast Asia. This study is designed to measure the pressure values in the normal foot of Malays in Pahang, Malaysia and its difference between different gender and body mass index (BMI). MATERIALS AND METHODS: A total of 400 feet of adult Malay subjects with no existing diabetes mellitus, lower limb and spine pain or problem are measured using Emed-q100 pedobarography device. RESULTS: 226 (56.5%) were females. 44.5% were with normal BMI followed by overweight (31.5%), and obese (24%). The mean-maximum-peak pressure (MPP) is 509kPa (SD 167) with no significant difference among gender and BMI. Most (38.5%, n=154) of the peak pressure area (PPA) are observed in 1st metatarsal head and big toe region (1MH&T), followed by 2nd metatarsal head (2MH) (34.3%, n=137). In the normal BMI group, 48.3% were in 1MH&T region while in the overweight and obese groups, 42.1% and 43.8%, respectively were in 2MH. This difference is significant ($\chi^2(df=8)=36.963$, $p<0.001$). There was no significant difference between PPA and gender. CONCLUSION: The MPP among Malays in this study was 509kPa(SD 167) and it is not affected by different genders or BMI. The PPA are most commonly fall on 1MH&T. There is a significant shift in the overweight and obese groups to the 2MHT. This finding can be used as initial reference for further studies, in Malaysia particularly. © 2022. IUM Medical Journal Malaysia

Author Keywords

Foot pressure; Pedobarography; Plantar pressure

References

- Hughes, J, Clark, P, Linge, K, Klenerman, L
A comparison of two studies of the pressure distribution under the feet of normal subjects using different equipment
(1993) *Foot Ankle*, 14 (9), pp. 514-519.
1
- Bacarin, T. A., Sacco, I. C. N., Hennig, E. M.
Plantar pressure distribution patterns during gait in diabetic neuropathy patients with a history of foot ulcers
(2009) *Clinics (Sao Paulo)*, 64 (2), pp. 113-120.
2
- Demirbüken, İ., Özgül, B., Timurtaş, E., Yurdalan, S. U., Çekin, M. D., Polat, M. G.
Gender and age impact on plantar pressure distribution in early adolescence
(2019) *Acta orthopaedica et traumatologica turcica*, 53 (5), pp. 215-220.
3
- Frykberg, R. G., Lavery, L. A., Pham, H., Harvey, C., Harkless, L., Veves, A.
Role of Neuropathy and High Foot Pressures in Diabetic Foot Ulceration
(1998) *Diabetes Care*, 21 (10), pp. 1714-1719.
4
- Stolwijk, N. M., Duysens, J., Louwerens, J. W. K., van de Ven, Y. H. M., Keijsers, N. L. W.
Flat Feet, Happy Feet Comparison of the Dynamic Plantar Pressure Distribution and Static Medial Foot Geometry between Malawian and Dutch Adults

(2013) *PLoS ONE*, 8 (2).

5

- Tuna, H., Birtane, M., Güldiken, S., Soysal, N. A., Taşpınar, Ö., Süt, N., Taştekin, N.
The effect of disease duration on foot plantar pressure values in patients with Type 2 Diabetes Mellitus

(2014) *Turkish Journal of Physical Medicine & Rehabilitation/ Turkiye Fiziksel Tip ve Rehabilitasyon Dergisi*, 60 (3).

6

- Fernandez-Segum, L. M., Diaz Mancha, J. A., Sanchez Rodriguez, R., Escamilla Martinez, E., Gomez Martin, B., Ramos Ortega, J.

Comparison of plantar pressures and contact area between normal and cavus foot
(2014) *Gait & Posture*, 39 (2), pp. 789-792.

7

- Maetzler, M., Bochdansky, T., Abboud, R. J.
Normal pressure values and repeatability of the Emed ST2 system

(2010) *Gait and Posture*, 32 (3), pp. 391-394.

8

- Putti, A. B., Arnold, G. P., Abboud, R. J.
Foot pressure differences in men and women
(2010) *Foot and Ankle Surgery*, 16 (1), pp. 21-24.

9

- Tomankova, K., Pridalova, M., Svoboda, Z., Cuberek, R.
Evaluation of plantar pressure distribution in relationship to body mass index in Czech women during walking

(2017) *Journal of the American Podiatric Medical Association*, 107 (3), pp. 208-214.

10

Correspondence Address

Nasuruddin M.H.; Department of Orthopaedics, Malaysia; email: drhaidar@iium.edu.my

Publisher: International Islamic University Malaysia

ISSN: 27352285

Language of Original Document: English

Abbreviated Source Title: IIUM Med. J. Malaysia.

2-s2.0-85133312603

Document Type: Article

Publication Stage: Final

Source: Scopus

ELSEVIER

Copyright © 2024 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

 **RELX Group™**