Individual, Host-Vector Interactions, and Environmental Risk Factors for *Plasmodium knowlesi* Malaria Among At-Risk Communities in Peninsular Malaysia: A Case-Control Study

By Pasi, H (Pasi, Hafizah) ; Mohamad, E (Mohamad, Emma) ; Azlan, AA (Azlan, Arina Anis) ; Hamzah, MR (Hamzah, Mohammad Rezal) ; Sulong, MR (Sulong, Mohd Rahim) ; Isa, A (Isa, Affendi) ; Genapathy, S (Genapathy, Sivasangari) ; Damanhuri, H (Damanhuri, Hazwani)

View Web of Science ResearcherID and ORCID (provided by Clarivate)

Source VECTOR-BORNE AND ZOONOTIC DISEASES
Volume: 25 Issue: 3 Page: 167-179
DOI: 10.1089/vbz.2024.0023

Published MAR 1 2025

Early Access NOV 2024

Indexed 2025-01-15

Document Type Article

Abstract Background: Highlighting the individual, host-vector interactions, and environmental risk factors for knowlesi malaria were consequential toward more focused and effective prevention and control strategies. This study aims to identify the

individual, host-vector interactions, and environmental risk factors for Plasmodium knowlesi malaria among at-risk communities in Peninsular Malaysia. Materials and Methods: A case-control study was conducted involving laboratory-confirmed cases of P. knowlesi malaria, while a locality-matched individual with no history of fever and tested negative for malaria was taken as control. Univariate and multiple logistic regression were applied to evaluate the potential risk factors among respondents using IBM SPSS Statistics for Windows, Version 26.0. Results: Results showed higher cases among males as compared to females (76.1% vs. 23.9%). Multiple logistic regression analysis showed being male is 3.51 higher risk ($p < 0.001$) to become a case. Respondents whose place of work or study is near the forest edge have 44.0% lower risk ($p = 0.030$), while those living in the Orang Asli village were 56.0% lower risk as compared to the organized village to become a case ($p = 0.035$). Conclusion: These findings demonstrated that gender emerges as an independent individual risk factor while residing near a forest edge, in an Orang Asli village, or occupying workers' longhouses situated in hilly areas lowered the environmental risk among respondents. These findings attested that alternative directions must be considered in addressing the known risk factors associated with this type of malaria and the design of prevention and control programs should be tailored to the unique characteristics of each population.

Keywords

Author Keywords: individual; host-vector interaction; environmental; risk factor; knowlesi malaria; at-risk communities

Keywords Plus: PARASITE; TRANSMISSION; SABAH; FALCIPARUM

Addresses

[1] Int Islamic Univ Malaysia Kuantan Campus, Dept Community Med, Kulliyyah Med, Kuantan 25200, Pahang, Malaysia

[2] Univ Kebangsaan Malaysia, Fac Social Sci & Humanities, Ctr Res Media & Commun, Bangi, Selangor, Malaysia

[3] Univ Malaysia Perlis, Fac Business & Commun, Kangar, Perlis, Malaysia

[4] Minist Hlth, Putrajaya, Malaysia

Categories/ Classification	Research Areas: Public, Environmental & Occupational Health; Infectious Diseases Citation Topics: 1 Clinical & Life Sciences › 1.217 Parasitology - Malaria, Toxoplasmosis & Coccidiosis › 1.217.59 Malaria Sustainable Development Goals: 03 Good Health and Well-being
Web of Science Categories	Public, Environmental & Occupational Health; Infectious Diseases
Language	English
Accession Number	WOS:001363194200001
PubMed ID	39585202
ISSN	1530-3667
eISSN	1557-7759
IDS Number	Z4T0M

– See fewer data fields

Citation Network

In Web of Science Core Collection

0 Citations

47
Cited References

Use in Web of Science

0
Last 180 Days

0
Since 2013

This record is from:

Web of Science Core Collection

- Science Citation Index Expanded (SCI-EXPANDED)