



MENU

Results for OPTIMIZING AN... &gt;

Optimizing Antibiotic Dosing: A Prospective Observational Study of Piperaci...



# Optimizing Antibiotic Dosing: A Prospective Observational Study of Piperacillin/Tazobactam Plasma Levels in Critically Ill Patients with Augmented Renal Clearance

**By** Negin, F (Negin, F.) ; Rahim, SAA (Rahim, S. A. Abdul) ; Nor, MBM (Nor, M. B. Mat) ; Mohamad, F (Mohamad, F.) ; Nordin, NS (Nordin, N. S.) ; Abdul-Aziz, MH (Abdul-Aziz, M. H.)

[View Web of Science ResearcherID and ORCID](#) (provided by Clarivate)

**Source** IIUM MEDICAL JOURNAL MALAYSIA  
Volume: 24 Issue: 3 Page: 48-53  
DOI: 10.31436/imjm.v24i03.2607

**Published** JUL 2025

**Indexed** 2025-08-12

**Document Type** Article

**Abstract** INTRODUCTION: Managing antibiotic dosing in critically ill patients presents challenges especially in achieving optimal therapeutic levels. Thus, we conducted a study to evaluate how augmented renal clearance (ARC) affects the attainment of pharmacokinetic/pharmacodynamic (PK/PD) targets in patients

receiving piperacillin/ tazobactam (PTZ) via continuous infusion.

**MATERIALS AND METHODS:** A singlecentred, prospective, observational study was conducted in intensive care unit at Sultan Ahmad Shah Medical Center @IIUM, Kuantan Pahang. A total of 43 adult patients with normal renal function treated as sepsis with standard PTZ doses via continuous infusion were included for the study and their blood were sampled for assessment of drug concentrations and PK/PD target attainment. **RESULTS:** There was substantial PK variability with 60% diagnosed with ARC and 37.2% of experienced piperacillin underexposure in which patients with ARC had significantly higher rates of underexposure at both distribution and steady-state phases. **CONCLUSION:** The high prevalence of ARC in these patients impacted the therapeutic PTZ levels and as many patients did not reach desired drug concentrations, there is increased risk of treatment failure without dose adjustment. These findings underscore the importance of individualized dosing strategies, particularly in critically ill patients with ARC, to optimize antibiotic therapy efficacy and mitigate the risk of inadequate treatment.

**Keywords**

**Author Keywords:** augmented renal clearance; critically ill; piperacillin-tazobactam; subtherapeutic

**Keywords Plus:** PHARMACOKINETICS

**Addresses**

<sup>1</sup> Kulliyyah Med, Dept Anaesthesia & Intens Care, Kuantan, Malaysia

<sup>2</sup> Kulliyyah Pharm, Dept Pharmaceut Chem, Kuantan, Malaysia

<sup>3</sup> Univ Queensland Ctr Clin Res UQCCR, Herston, Australia

**Categories/  
Classification**

Research Areas: General & Internal Medicine

Citation [1 Clinical & Life Sciences](#) > [1.23 Antibiotics & Antimicrobials](#) > [1.23.422 Antibiotic Pharmacokinetics](#)

Sustainable Development Goals: [03 Good Health and Well-being](#)

**Web of Science  
Categories**

Medicine, General & Internal

[+ See more data fields](#)

### Citation Network

In Web of Science Core Collection

0 Citations

14  
Cited References

### Use in Web of Science

00

Last 180 DaysSince 2013

### This record is from:

Web of Science Core Collection

- Emerging Sources Citation Index (ESCI)

### Suggest a correction

If you would like to improve the quality of the data in this record, please [Suggest a correction](#)