

[Alert Results](#) > [Low Bone Mineral Density in Men Living with HIV on Tenofovir Antiretroviral...](#)

Low Bone Mineral Density in Men Living with HIV on Tenofovir Antiretroviral Therapy: A Cross-Sectional Exploratory Study from a Malaysian Tertiary Hospital

By [Daud, MR](#) (Daud, Muhamad Riduan) ^[1]; [Periyasamy, P](#) (Periyasamy, Petrick) ^[1]; [Chin, KY](#) (Chin, Kok-Yong) ^[2]; [Kori, N](#) (Kori, Najma) ^[1]; [Zainulabid, UA](#) (Zainulabid, Ummu Afeera) ^[3]; [Yew, SQ](#) (Qian Yew, Sheng) ^[4]; [Kazmin, NEM](#) (Mohammad Kazmin, Nur Ezzaty) ^[5]; [Cheong, XK](#) (Khee Cheong, Xiong) ^[1]

[View Web of Science ResearcherID and ORCID](#) (provided by Clarivate)

Source [TROPICAL MEDICINE AND INFECTIOUS DISEASE](#)

[← View Journal Impact](#)

Volume: 11 Issue: 2

DOI: 10.3390/tropicalmed11020038

Article Number 38

Published JAN 29 2026

Indexed 2026-03-07

Document Type Article

Abstract Background and objectives: Low bone mineral density (BMD) is a recognized complication in people living with HIV (PLHIV) that remains under-addressed, particularly in Malaysia. Known contributing factors for low BMD include advanced age, HIV infection itself, and prolonged use of anti-retroviral therapy (ART), particularly tenofovir-based regimens. There are limited data on the burden of low BMD in the HIV population in Malaysia. This study aimed to determine the prevalence of low BMD among men living with HIV on tenofovir disoproxil fumarate (TDF) and to identify the possible associated factors compared to a group of healthy individuals matched for age and ethnicity. Methods: This is single-center cross-sectional study involved 112 men (56 HIV-positive individuals and 56 uninfected individuals matched for age and ethnicity) who underwent dual-energy X-ray absorptiometry (DXA) scans of the femoral neck and lumbar spine. Sociodemographic, clinical lifestyle, and laboratory data, including FRAX scores for those aged more than 40 years old, were collected. Results: The prevalence of low BMD at the femoral neck in HIV-infected men on tenofovir disoproxil fumarate was significantly higher compared to healthy individuals (32.1% vs. 16.1%; $p < 0.05$). Low BMD prevalence at the lumbar spine was higher in the HIV group (8.9% vs. 3.6%; $p = 0.463$) but was not statistically significant. Older age and low body mass index (BMI) were found to be significantly associated with reduced BMD in HIV patients. Chronic kidney disease stage 2 and 3a was linked with low femoral neck BMD. HIV-related factors (duration of illness, duration of ART exposure, and CD4+ counts) showed no significant associations to low BMD. The 10-year risk of major osteoporotic and hip fractures estimated by the FRAX tool was low in both groups, and no participant exceeded the recommended intervention threshold. Conclusions: Men with HIV on tenofovir disoproxil fumarate have a higher prevalence of low BMD, particularly at the femoral neck. Traditional risk factors were more closely associated with low BMD compared to HIV-related factors and specific markers, supporting the need for

routine bone health screening and preventive strategies in this population.

Keywords

Author Keywords: [HIV](#); [anti-retroviral therapy](#); [tenofovir](#); [bone mineral density](#); [DXA scan](#)

Keywords Plus: [INFECTION](#); [RECOMMENDATIONS](#); [MANAGEMENT](#); [DISEASE](#)

Author Information

Corresponding Address: Khee Cheong, Xiong (corresponding author)

▼ Univ Kebangsaan Malaysia, Fac Med, Dept Med, Infect Dis Unit, Cheras 56000, Kuala Lumpur, Malaysia

E-mail Addresses : cheongxk@gmail.com

Addresses :

▼ ¹ Univ Kebangsaan Malaysia, Fac Med, Dept Med, Infect Dis Unit, Cheras 56000, Kuala Lumpur, Malaysia

▼ ² Univ Kebangsaan Malaysia, Fac Med, Dept Pharmacol, Cheras 56000, Kuala Lumpur, Malaysia

▼ ³ Int Islamic Univ Malaysia, Dept Internal Med, Kulliyah Med, Kuantan 25200, Malaysia

▼ ⁴ Univ Kebangsaan Malaysia, Fac Med, Dept Publ Hlth, Cheras 56000, Kuala Lumpur, Malaysia

▼ ⁵ ...

 [Free Full Text from Publisher](#)

[View Full Text on ProQuest](#)

[Full Text Links](#) ▼

[Export](#) ▼

[Add To Marked List](#)

< 1 of 1 >

Classification [Medicine](#)

Web of Science Categories [Infectious Diseases](#); [Parasitology](#); [Tropical Medicine](#)

Funding



MENU



▼ View funding text

Funding agency	Grant number
Universiti Kebangsaan Malaysia	FF-2024-500

Language	English
Accession Number	WOS:001700952900001
PubMed ID	41746008
eISSN	2414-6366
IDS Number	EH1HU

— See fewer data fields

Journal information
