

Scopus

Document details

[Back to results](#) | 1 of 1

[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More...](#)
[Full Text](#)
[View at Publisher](#)

Renewable Energy

Volume 63, March 2014, Pages 617-623

Reduction of low grade iron ore pellet using palm kernel shell (Article)

Abd Rashid, R.Z.^a, Mohd. Salleh, H.^a, Ani, M.H.^a, Yunus, N.A.^a, Akiyama, T.^b, Purwanto, H.^a^aFaculty of Engineering, International Islamic University Malaysia, Jalan Gombak, P.O. Box 10, 50728 Kuala Lumpur, Malaysia^bCenter for Advanced Research of Energy and Materials, Hokkaido University, Sapporo 060-8628, Japan

Abstract

[View references \(28\)](#)

Effective use of local iron ore and biomass waste as energy and material resources in iron making is an interesting economic prospect since Malaysia imports iron ore to supply its domestic steel consumption while there is an abundance of biomass waste from the palm oil industry. In this work, a composite pellet made of Malaysian iron ore with palm kernel shell (PKS) waste was subjected to reduction tests using an electric tube furnace to investigate the effect of temperature and PKS content on reduction rate. Several iron ore samples taken from different mining locations were subjected to thermal and X-ray diffraction (XRD) analysis. The rate of iron ore reduction increased with increasing temperature up to 900°C. XRD analysis revealed that the original iron ore mainly contains iron oxide hydrate and was converted into simple hematite after heating and then become magnetite after reduction. The Fe content in the original ore increased almost 12% when 40wt% of PKS was used. The iron oxide was successfully reduced to magnetite and small amount of wustite when up to 20wt% of PKS was present in the mixture. Besides, 20wt% of PKS in reduction process can reduce CO₂ emissions by almost 18.69wt% as well as decrease carbon consumption by 19.78wt%. Thus, the utilization of biomass as a reducing agent for low grade iron ore reduction is an attractive method for upgrading iron ore as well as reducing CO₂ emissions. © 2013 Elsevier Ltd.

Author keywords

[Biomass](#) [CO₂ emission](#) [Iron making](#) [Iron ore](#) [Palm kernel shell waste](#) [Reduction](#)

Indexed keywords

[Metrics](#) [View all metrics](#)

15 Citations in Scopus

86th Percentile

1.16 Field-Weighted Citation Impact



PlumX Metrics

Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 15 documents

Comparison of numerical solutions for a mathematical model of theoretical combustion and conductivity variation of shell palm in an fixed bed reactor | Comparación entre soluciones numéricas de un modelo matemático de combustión teórica y variación de la conductividad del cuello de palma en un reactor de lecho fijo
Ahumada, L.M. , Audivet, C.P. , Bula, A.J.
(2017) *Informacion Tecnologica*

Fe-Assisted Hydrothermal Liquefaction of Lignocellulosic Biomass for Producing High-Grade Bio-Oil
Miyata, Y. , Sagata, K. , Hirose, M.

(2017) *ACS Sustainable Chemistry and Engineering*
Current status and potential of biomass utilization in ferrous metallurgical industry

Wei, R. , Zhang, L. , Cang, D.
(2017) *Renewable and Sustainable Energy Reviews*

[View all 15 citing documents](#)
Inform me when this document is cited in Scopus:

[Set citation alert](#)
[Set citation feed](#)

Carbon consumption
 Economic prospects
 Effect of temperature
 Increasing temperatures
 Iron making
 Material resources
 Palm kernel shells
 Steel consumption
 Carbon consumption
 Economic prospects
 Effect of temperature
 Increasing temperatures
 Iron making
 Material resources
 Palm kernel shells
 Steel consumption

Engineering controlled terms: Biomass Carbon dioxide Hydrates Iron ores Iron oxides Magnetite Reduction
 Biomass Carbon dioxide Hydrates Iron Iron and steel industry Iron ores
 Iron oxides Magnetite Ore analysis Palm oil Pelletizing Reduction Temperature

Engineering main heading: Iron ore reduction Iron ore reduction

GEOBASE Subject Index: biofuel carbon dioxide carbon emission energy resource iron ore iron oxide
 oil industry ore deposit renewable resource shell biomass carbon dioxide
 carbon emission energy resource fuel consumption iron ore oil industry reduction
 steel waste treatment

Regional Index: Malaysia Malaysia

Related documents

Carbon doped iron ore using palm kernel shell

Rusila, Z.A.R. , Purwanto, H. , Salleh, H.M.
(2013) Advanced Materials Research

Enhancement of magnetic properties of Malaysian iron ore by reduction roasting using oil palm empty fruit bunch

Abd Rashid, R.Z. , Yunus, N.A. , Mohd Salleh, H.
(2014) ISIJ International

Effect of reduction roasting by using bio-char derived from empty fruit bunch on the magnetic properties of Malaysian iron ore

Yunus, N.A. , Ani, M.H. , Salleh, H.M.
(2014) International Journal of Minerals, Metallurgy and Materials

[View all related documents based on references](#)

[Find more related documents in Scopus based on:](#)

[Authors >](#) [Keywords >](#)

ISSN: 09601481
Source Type: Journal
Original language: English

DOI: 10.1016/j.renene.2013.09.046
Document Type: Article

References (28)

[View in search results format >](#)

All [Export](#)  [Print](#)  [E-mail](#) [Save to PDF](#) [Create bibliography](#)

1 Murakami, T., Kasai, E.

Utilization of ores with high combined water content for composite iron coke ore-carbon

(2011) *ISIJ International*, 51 (8), pp. 1220-1226. Cited 8 times.
 doi: 10.2355/isijinternational.51.1220

[View at Publisher](#)