

INTERNATIONAL CONFERENCE ON ANTIOXIDANTS & DEGENERATIVE DISEASES (ICADD 2015)



“From Molecules to Human”

June 3 - 4, 2015

ISTANA HOTEL | KUALA LUMPUR, MALAYSIA

Organised by:
Antioxidants, Ageing and Degenerative Diseases Research Group
Universiti Kebangsaan Malaysia

www.icadd.org.my



DD-06

Assessment of Reactive Oxygen Species Status and Neuroprotection by Vitamin E in Chronic Cerebral Hypoperfusion Induced Neurodegeneration in Rats

Saveed S, Saxena AK, Azzubaidi MS, Talib NA

Basic Medical Sciences, Kulliyyah of Medicine; International Islamic University Malaysia, Kuantan Campus, Jalan Istana, Bandar Indera Mahkota 25200 Kuantan, Pahang, Malaysia

Background: Chronic cerebral hypoperfusion has been linked with cognitive decline in ageing and Alzheimer's disease (AD). A reduced cerebral blood flow (CBF) is considered to be a predictive marker for the progression into Alzheimer's disease. In order to tackle the neuropathological consequences of such a reduction in CBF a similar condition was created in experimental animals. Chronic cerebral hypoperfusion induced neurodegeneration was produced by common carotid artery occlusion (2 vessel occlusion, 2VO) in rats. Since oxidative stress, leading to neuronal apoptosis and death, is one of the mechanisms which is thought to play a significant role in chronic degenerative neurological disorders, the present study was designed to assess the neuroprotective role of Vitamin E in chronic cerebral hypoperfusion-induced neurodegeneration.

Methods: After acclimatization, thirty Sprague Dawley rats weighing 200-250 g were equally divided into three groups. Group A – sham control, Group B – 2VO, and Group C – 2VO-E (treated daily with Vitamin E, 100 mg/kg, orally following 2VO). On the 8th week, all the rats were euthanized and the hippocampi were isolated. Viable neuronal cell count in the hippocampal CA-1 region was estimated.

Results: There was significant difference in neuronal cell death in 2VO group as compared to sham control group. In Vitamin E-treated 2VO (2VO-E) rats, the viable neuronal cell count of the hippocampal CA-1 region was significantly higher as compared to the untreated 2VO group.

Conclusion: Vitamin E is an effective neuroprotective agent in chronic cerebral hypoperfusion-induced neurodegeneration in rats and can be effectively utilized in the management of AD. The antioxidant activity of Vitamin E is in progress and will be evaluated by the measurement of Isoprostane F2 through Enzyme-Linked Immunosorbent Assay.

Keywords: Vitamin E, chronic cerebral hypoperfusion, neurodegeneration, Alzheimer's disease, oxidative stress