

## Medical Research Symposium & Pacific Partnership in conjunction with

## Kuantan Research Day 2016

**International Research Network : The Way Forward**

# ABSTRACTS

**Organized by :**



**Kulliyah of Medicine, IIUM  
Research Management Centre, Kuantan**

In collaboration:



## Basic Health Sciences

## Poster

Abstract ID: 74

## DNA methylation of *DISC1* gene in schizophrenia using methylight Taqman assay

Nour El Huda Abd Rahim<sup>a</sup> | Mohd Nabil Fikri Rahim<sup>a</sup> | Norsidah Ku Zaifah<sup>a</sup> | Hanisah Mohd Noor<sup>b</sup> | Kartini Abdullah<sup>b</sup> | Norlelawati A. Talib<sup>c</sup>

<sup>a</sup>Department of Basic Medical Sciences, Kulliyyah of Medicine, International Islamic University Malaysia

<sup>b</sup>Department of Psychiatry, Kulliyyah of Medicine, International Islamic University Malaysia

<sup>c</sup>Department of Pathology and Laboratory Medicine, Kulliyyah of Medicine, International Islamic University Malaysia

**Introduction:** Significant evidences from functional studies have shown that *DISC1* gene has a role in the pathogenesis of schizophrenia, although the basis of the genetic defect has yet to be established. There has been a shift of emphasis from *DISC1* gene variations to other types of genetic defects namely copy number and epigenetic, both of which have not been well investigated. Thus, the aim of this study is to examine the DNA methylation status of *DISC1* gene in patients with schizophrenia. **Methods:** In this study, 239 subjects were included, 117 schizophrenia patients and 122 healthy controls. Genomic DNA was derived from peripheral blood and bisulfite converted. The DNA methylation level was quantitatively measured by Methylight Taqman analysis. Sociodemographic and the clinical parameters were noted. The severity of the clinical symptoms was assessed using Positive and Negative Syndrome Scale (PANSS). **Results:** The mean age and gender distribution between the study groups were similar. There was no significant difference in the methylation level of *DISC1* between the patients and control group. When patients were compared by age, duration of illness, age at diagnosis, body mass index, smoking status, PANSS score and types of antipsychotic treatment, the DNA methylation level of *DISC1*, did not show any significant difference ( $p>0.05$ ). **Conclusions:** This study found no significant difference in methylation level of *DISC1* gene between schizophrenia patients and healthy control. Therefore, it is suggested that aberrant DNA methylation of *DISC1* most probably does not contribute to the pathogenesis of schizophrenia.

**KEYWORDS:** Schizophrenia, DNA methylation, *DISC1*