



Document details

[Back to results](#) | 1 of 1

[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)

[Full Text](#) [View at Publisher](#)

Marine and Freshwater Behaviour and Physiology
Volume 49, Issue 3, 3 May 2016, Pages 159-171

Morphogenesis of free neuromasts in the larvae of brown-marbled grouper *Epinephelus fuscoguttatus* (Article)

Mukai, Y.^a Lim, L.S.^b

^aDepartment of Marine Science, Kulliyah of Science, International Islamic University Malaysia, Kuanatan, Malaysia

^bBorneo Marine Research Institute, University Malaysia Sabah, Kota Kinabalu, Malaysia

Abstract

[View references \(43\)](#)

Newly hatched larvae had one pair of free neuromasts behind the eyes. As the larvae grew, free neuromasts increased in number. The apical surface of sensory epithelium widened and subsequently elongated. The number of sensory hair cells increased and the directions of maximum sensitivity became both anteroposterior and dorsoventral on the trunk. Before notochord flexion, only the anteroposterior type was observed. After notochord flexion, two types of neuromasts were observed on the trunk. On the head, the orientation of free neuromasts formed a tangential line to concentric circles around the eyes and nostrils. Free neuromasts on the head could therefore receive stimuli from various angles from predators or zooplanktons. This suggests that these free neuromasts play a role in compensating for a dead angle of vision, and an important role in detecting zooplankton under scotopic vision. Canal organs were observed on the head and operculum in 40-d-old animals. © 2016 Informa UK Limited, trading as Taylor & Francis Group.

SciVal Topic Prominence

Topic: Lateral Line System | Zebrafish | Artificial lateral

Prominence percentile: 84.606

Author keywords

[cupulae](#) [fish larvae](#) [Free neuromasts](#) [lateral line](#) [maximum sensitivity](#) [neuromast morphology](#)

Indexed keywords

GEOBASE Subject Index: [fish](#) [larva](#) [morphogenesis](#) [morphology](#) [physiological response](#) [zooplankton](#)

Species Index: [Animalia](#) [Epinephelus fuscoguttatus](#)

Metrics [View all metrics >](#)

3 Citations in Scopus

25th percentile

0.25 Field-Weighted Citation Impact



PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 3 documents

Identification of Candidate Growth-Related SNPs and Genes Using GWAS in Brown-Marbled Grouper (*Epinephelus fuscoguttatus*)

Yang, Y. , Wu, L. , Wu, X. (2020) *Marine Biotechnology*

Immunity, feed, and husbandry in fish health management of cultured *Epinephelus fuscoguttatus* with reference to *Epinephelus coioides*

Chieng, C.C.Y. , Daud, H.M. , Yusoff, F.M. (2018) *Aquaculture and Fisheries*

Feeding and development of warm water marine fish larvae in early life

Hu, J. , Liu, Y. , Ma, Z. (2018) *Emerging Issues in Fish Larvae Research*

[View all 3 citing documents](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

[Set citation feed >](#)

References (43)

View in search results format >

All Export Print E-mail Save to PDF Create bibliography

- 1 Blaxter, J.H.S.

Development of sense organs and behaviour of teleost larvae with special reference to feeding and predator avoidance

(1986) *Transactions of the American Fisheries Society*, 115 (1), pp. 98-114. Cited 497 times.
doi: 10.1577/1548-8659(1986)115[98:NLFCD0]2.0.CO;2

[View at Publisher](#)

- 2 Blaxter, J.H.S.

Structure and development of the lateral line
(1987) *Biol Rev*, 62, pp. 471-514. Cited 92 times.

Development of sensory organs in larvae of African catfish *Clarias gariepinus*

Mukai, Y. , Tuzan, A.D. , Lim, L.S.
(2008) *Journal of Fish Biology*

Neuromast formation in the prehatching embryos of the cod fish, *Gadus macrocephalus* *Tilesius*

Otsuka, M. , Nagai, S.-I.
(1997) *Zoological Science*

[View all related documents based on references](#)

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

- 3 Blaxter, J.H.S., Fuiman, L.A.

Function of the free neuromasts of marine teleost larvae
(1989) *The mechanosensory lateral line*, pp. 481-499. Cited 92 times.
Coombs S., Görner P., Münz H., (eds), New York(NY): Springer-Verlag

- 4 Blaxter, J.H.S., Gray, J.A.B., Best, A.C.G.

Structure and development of the free neuromasts and lateral line system of the herring

(1983) *Journal of the Marine Biological Association of the United Kingdom*, 63 (2), pp. 247-260. Cited 61 times.
doi: 10.1017/S0025315400070648

[View at Publisher](#)

- 5 Diaz, J.P., Prié-Granié, M., Kentouri, M., Varsamos, S., Connes, R.

Development of the lateral line system in the sea bass

(2003) *Journal of Fish Biology*, 62 (1), pp. 24-40. Cited 23 times.
doi: 10.1046/j.1095-8649.2003.00004.x

[View at Publisher](#)

- 6 Disler, N.N.

(1971) *Lateral line sense organs and their importance in fish behavior*. Cited 27 times.
Mills H., Yariv M., (eds), Jerusalem: Israel Program for Science Translation

Related documents

Structure and development of free neuromasts in barramundi, *Lates calcarifer* (block)

Mukai, Y. , Chai, L.L. , Shaleh, S.R.M.
(2007) *Zoological Science*

- 7 Flock, Å.

Electron microscope and electrophysiological studies on the lateral line canal organ
(1965) *Acta Otolaryngol Suppl*, 199, pp. 1-90. Cited 211 times.