

## On classification of $m$ -dimensional algebras

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 819 012012

(<http://iopscience.iop.org/1742-6596/819/1/012012>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.48.222.12

This content was downloaded on 06/04/2017 at 10:14

Please note that [terms and conditions apply](#).

You may also be interested in:

### [ALGEBRAS THAT ARE DEFORMABLE INTO FREE ALGEBRAS](#)

A A Boyarkin

### [A Complex Higher-Dimensional Lie Algebra with Real and Imaginary Structure Constants as Well as Its Decomposition](#)

Zhang Yu-Feng

### [Classification of weakly commutative complex homogeneous spaces](#)

Ivan V Losev

### [Some examples of finitely generated algebras](#)

V T Markov

### [Defining relations for nilpotent subalgebras of modular classical Lie algebras](#)

A S Dzhumadil'daev and Sh Sh Ibraev

### [Bilinear forms on fermionic Novikov algebras](#)

Zhiqi Chen and Fuhai Zhu

### [Bianchi spaces and their three-dimensional isometries as S-expansions of two-dimensional isometries](#)

Ricardo Caroca, Igor Kondrashuk, Nelson Merino et al.

### [Diassociative algebras and their derivations](#)

I M Rikhsiboev, I S Rakhimov and W Basri

### [On Valdivia-type universal topological algebras](#)

L B Shapiro

# On classification of $m$ -dimensional algebras

**U. Bekbaev**

Department of Science in Engineering, Faculty of Engineering, International Islamic University Malaysia, P.O. Box 10, 50728 Kuala Lumpur, Malaysia

E-mail: [bekbaev@iium.edu.my](mailto:bekbaev@iium.edu.my)

**Abstract.** A constructive approach to the classification and invariance problems, with respect to basis changes, of the finite dimensional algebras is offered. A construction of an invariant open, dense (in the Zariski topology) subset of the space of structure constants of algebras is given. A classification of all algebras with structure constants from this dense set is given by providing canonical representatives of their orbits. A finite system of generators for the corresponding field of invariant rational functions of structure constants is shown.

## 1. Introduction

The classification of finite dimensional algebras is an important problem in Algebra. For example, the classification of finite dimensional simple and semi-simple associative algebras by Wedderburn, the classification of finite dimensional simple and semi-simple Lie algebras by Cartan are well known. Their classifications are examples of structural (basis free, invariant) approaches to the classification problem of algebras. A such approach used to classify one class of algebras can not be used with the same purpose to another class. Usually the obtained classification covers only a marginal part and the main bulk of the considered algebras is left unclassified. The structural approach becomes more difficult and unclear when one considers more general types of algebras. Another disadvantage of such approach is that the classification is assumed to be only with respect to the general linear group. In reality one may be interested also in classification of algebras with respect to specific changes of basis.

Another approach to the classification and invariants problems of finite dimensional algebras is coordinate (basis based, structure constants) approach. In contrast to the first approach the coordinate approach provides classification of all algebras from the bulky part of all given dimensional algebras. In this sense these two approaches are complementary to each other.

For the small dimensional cases for such approach one can see [1-3]. In any finite dimensional algebra case a basis based approach is considered in [4]. In this paper we also consider the classification and invariants problems of finite dimensional algebras. Though there are some intersecting results in this paper with those of [4] our approach to these problems is quite different, constructive and used tools are more elementary than of [4]. We note that our approach is applicable in the case of some classical subgroups of general linear group as well [5].

In general for the given dimension we provide a method how to construct an invariant, open, dense subset of the space of structure constants of algebras and classify all algebras who's systems of structure constants are from this set. The classification is given by providing the structure constants of the canonical algebra for each such algebra. We provide a basis for the field of invariant rational functions of structure constants as well.



The paper is organized as follows. The key results which are used to obtain the classification and invariants of algebras are presented in Section 2. Section 3 can be considered as a realization of Section 2 results in the case of representation of general linear group in the space of structure constants of algebras.

## 2. Preliminaries

In this section we consider a linear representation of a subgroup of the general linear group and under an assumption prove some general results on classification and invariance problems with respect to this subgroup.

Let  $n, m$  be any natural numbers,  $\tau : (G, V) \rightarrow V$  be a fixed linear algebraic representation of an algebraic subgroup  $G$  of  $GL(m, F)$  on  $V$  over  $F$ , where  $F$  is any field and  $V$  is  $n$ -dimensional vector space over  $F$ . Further we consider this representation under the following assumption:

**Assumption.** There exists a nonempty  $G$ -invariant subset  $V_0$  of  $V$  and an algebraic map  $P : V_0 \rightarrow G$  such that

$$P(\tau(g, \mathbf{v})) = P(\mathbf{v})g^{-1} \quad (2.1)$$

whenever  $\mathbf{v} \in V_0$  and  $g \in G$ .

Note that due to this Assumption  $G$ -stabilizer of any  $\mathbf{v} \in V_0$  is trivial. The following result is a classification theorem for elements of  $V_0$  with respect to  $G$ .

**Theorem 2.1.** *Elements  $\mathbf{u}, \mathbf{v} \in V_0$  are  $G$ -equivalent, that is  $\mathbf{u} = \tau(g, \mathbf{v})$  for some  $g \in G$ , if and only if  $\tau(P(\mathbf{u}), \mathbf{u}) = \tau(P(\mathbf{v}), \mathbf{v})$ .*

*Proof.* If  $\mathbf{u} = \tau(g, \mathbf{v})$  then  $\tau(P(\mathbf{u}), \mathbf{u}) = \tau(P(\tau(g, \mathbf{v})), \tau(g, \mathbf{v})) =$

$$\tau(P(\mathbf{v})g^{-1}, \tau(g, \mathbf{v})) = \tau(P(\mathbf{v}), \tau(g^{-1}, \tau(g, \mathbf{v}))) = \tau(P(\mathbf{v}), \mathbf{v}).$$

Visa versa, if  $\tau(P(\mathbf{u}), \mathbf{u}) = \tau(P(\mathbf{v}), \mathbf{v})$  then

$$\tau(P(\mathbf{u})^{-1}P(\mathbf{v}), \mathbf{v}) = \tau((P(\mathbf{u}))^{-1}, \tau(P(\mathbf{v}), \mathbf{v})) = \tau((P(\mathbf{u}))^{-1}, \tau(P(\mathbf{u}), \mathbf{u})) = \mathbf{u}$$

that is  $\mathbf{u} = \tau(g, \mathbf{v})$ , where  $g = P(\mathbf{u})^{-1}P(\mathbf{v})$ . □

So this theorem provides for any  $\mathbf{v} \in V_0$  the canonical representative, namely  $\tau(P(\mathbf{v}), \mathbf{v})$ , of its  $G$ -orbit and this canonical representative is same for all elements of the orbit. It shows also that the system of components of  $\tau(P(\mathbf{x}), \mathbf{x})$  is a separating system of invariants for the  $G$ -orbits in  $V_0$ , where  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  is an algebraic independent system of variables over  $F$ .

Further in this paper it is assumed that  $F$  is an algebraically closed field of characteristic zero,  $V_0$  in the above Assumption is dense (in Zariski topology) in  $V$ . In such case for any  $\mathbf{u}, \mathbf{v} \in V_0$  one has  $P(\tau(P(\mathbf{u}), \mathbf{v})) = P(\mathbf{v})P(\mathbf{u})^{-1}$  and due to density of  $V_0$  in  $V$  one has

$$P(\tau(P(\mathbf{y}), \mathbf{x})) = P(\mathbf{x})P(\mathbf{y})^{-1}, \quad (2.2)$$

where  $\mathbf{y} = (y_1, y_2, \dots, y_n)$  is also an algebraic independent system of variables over  $F$ .

**Theorem 2.2.** *The field of  $G$ -invariant rational functions  $F(\mathbf{x})^G$  is generated over  $F$  by the system of components of  $\tau(P(\mathbf{x}), \mathbf{x})$  and moreover the equality  $P(\tau(P(\mathbf{x}), \mathbf{x})) = I_m$  holds true, where  $I_m$  stands for  $m$ -order identity matrix.*

*Proof.* It is evident that all components of  $\tau(P(\mathbf{x}), \mathbf{x})$  are in  $F(\mathbf{x})^G$ . If  $f(\mathbf{x}) = f(\tau(g, \mathbf{x}))$  for all  $g \in G$  then, in particular,  $f(\mathbf{x}) = f(\tau(P(\mathbf{u}), \mathbf{x}))$  whenever  $\mathbf{u} \in V_0$ . It implies, as far as  $V_0$  is dense in  $V$ , that for the variable vector  $\mathbf{y}$  the equality

$$f(\mathbf{x}) = f(\tau(P(\mathbf{y}), \mathbf{x}))$$

holds true. In  $\mathbf{y} = \mathbf{x}$  case one gets that  $f(\mathbf{x}) = f(\tau(P(\mathbf{x}), \mathbf{x}))$  and  $P(\tau(P(\mathbf{x}), \mathbf{x})) = I_m$  due to (2.2). □

**Corollary 2.3.** *The field  $F(\mathbf{x})$  is generated over  $F(\mathbf{x})^G$  by the system of components of  $P(\mathbf{x})$ .*

*Proof.* Indeed  $F(\mathbf{x})^G(P(\mathbf{x})) = F(\tau(P(\mathbf{x}), \mathbf{x}))(P(\mathbf{x})) = F(\tau(P(\mathbf{x}), \mathbf{x}), P(\mathbf{x}))$  and  $\tau(P(\mathbf{x})^{-1}, \tau(P(\mathbf{x}), \mathbf{x})) = \mathbf{x}$  and therefore  $F(\mathbf{x})^G(P(\mathbf{x})) = F(\mathbf{x})$ .  $\square$

**Proposition 2.4.** *The equality  $\text{trdeg} F(P(\mathbf{x}))/F = \dim G$  holds true.*

*Proof.* To prove the equality it is enough to show equality of the vanishing ideals of  $P(\mathbf{x})$  and  $G$ . If polynomial  $p$  vanished on  $P(\mathbf{x})$ , that is  $p[P(\mathbf{x})] = 0$ , then  $p[P(\tau(g, \mathbf{x}))] = p[P(\mathbf{x})g^{-1}] = 0$ . In particular,  $p[g] = 0$  for any  $g \in G$  that is  $p$  vanishes on  $G$  as well.

If  $p[g] = 0$  for any  $g \in G$  then, in particular,  $p[P(\mathbf{u})] = 0$  for any  $\mathbf{u} \in V_0$ . Due to density of  $V_0$  in  $V$  one has  $p[P(\mathbf{x})] = 0$ .  $\square$

**Theorem 2.5.** *The equality  $\text{trdeg} F(\mathbf{x})^G/F = n - \dim G$  holds true.*

*Proof.* Let  $\widetilde{P(\mathbf{x})}$  stand for any system of entries of  $P(\mathbf{x})$  which is a transcendence basis for the field  $F(P(\mathbf{x}))$  over  $F$ . We show that the system  $\widetilde{P(\mathbf{x})}$  is algebraic independent over  $F(\mathbf{x})^G$  as well. Indeed let  $p[(y_{ij})_{i,j=1,2,\dots,m}]$  be any polynomial over  $F(\mathbf{x})^G$  for which  $p[\widetilde{P(\mathbf{x})}] = 0$  that is  $p_{\mathbf{v}}[\widetilde{P(\mathbf{v})}] = 0$  for all  $\mathbf{v} \in V_1$ , where  $V_1$  is a  $G$ -invariant nonempty open subset of  $V_0$ , where  $p_{\mathbf{v}}[(y_{ij})_{i,j=1,2,\dots,m}]$  stands for the polynomial obtained from  $p[(y_{ij})_{i,j=1,2,\dots,m}]$  by substitution  $\mathbf{v}$  for  $\mathbf{x}$ . The equality  $0 = p_{\mathbf{v}}[\widetilde{P(\mathbf{v})}] = p_{\tau(g, \mathbf{v})}[P(\tau(g, \mathbf{v}))] = p_{\mathbf{v}}[P(\mathbf{v})g^{-1}]$  implies that  $p_{\mathbf{v}}[g] = 0$  for any  $g \in G$ . Therefore  $p_{\mathbf{v}}[\widetilde{P(\mathbf{x})}] = 0$ , that is  $p_{\mathbf{v}}[(y_{ij})_{i,j=1,2,\dots,m}]$  is zero polynomial for any  $\mathbf{v} \in V_1$ . It means that  $p[(y_{ij})_{i,j=1,2,\dots,m}]$  is zero polynomial itself. Now due to  $F \subset F(\mathbf{x})^G \subset F(\mathbf{x})$ ,  $\text{tr.deg.} F(\mathbf{x})/F = n$  and **Corollary 2.3** one has the required result.  $\square$

**Remark 2.6.** *According to Theorem 2.2 for entries of  $\tau(P(\mathbf{x}), \mathbf{x})$  one has following  $m^2$  equalities  $P(\tau(P(\mathbf{x}), \mathbf{x})) = I_m$ . If it enables one to express some  $\dim G$  entries of  $\tau(P(\mathbf{x}), \mathbf{x})$  by the rest of its  $n - \dim G$  entries, which may happen in some cases, one gets rationality of the extension  $F \subset F(\mathbf{x})^G$  due to Theorem 2.5.*

In particular, for  $G = GL(m, F)$  case due to **Theorem 2.2** one has the following result.

**Corollary 2.7.** *The transcendence degree of  $F(\mathbf{x})^{GL(m, F)}$  over  $F$  equals to  $n - m^2$  and the field extension  $F(\mathbf{x})^{GL(m, F)} \subset F(\mathbf{x})$  is a pure transcendental extension. For a transcendental basis one can take the system of components of  $P(\mathbf{x})$ .*

**Remark 2.8.** *The Assumption may be productive in a slightly different form as well: There exists a nonempty  $G$ -invariant subset  $V_0$  of  $V$  and an algebraic map  $P : V_0 \rightarrow GL(m, F)$  such that  $P(\tau(g, \mathbf{v})) = P(\mathbf{v})g^{-1}$  whenever  $\mathbf{v} \in V_0$  and  $g \in G$ . In this case Theorem 2.1 can be formulated in the following form: Elements  $\mathbf{u}, \mathbf{v} \in V_0$  are  $G$ -equivalent if and only if  $\tau(P(\mathbf{u}), \mathbf{u}) = \tau(P(\mathbf{v}), \mathbf{v})$  and  $P(\mathbf{u})^{-1}P(\mathbf{v}) \in G$ . For the classical subgroups of  $GL(m, F)$  the relation  $P(\mathbf{u})^{-1}P(\mathbf{v}) \in G$  can be written in terms of equality of some  $G$ -invariants on  $P(\mathbf{u})$  and  $P(\mathbf{v})$ , for example, in  $G = O(m, F)$ -the orthogonal group case it means  $P(\mathbf{u})P(\mathbf{u})^t = P(\mathbf{v})P(\mathbf{v})^t$ . So in such cases for the classical subgroups the separating system of invariants can be listed easily. But whether the separating system generates the corresponding field of invariant functions is unclear even one admits density of  $V_0$  in  $V$ . Therefore one comes to the following problem.*

**Problem 2.9.** *If some system of invariants  $\{f_1(x), f_2(x), \dots, f_k(x)\}$  from  $F(\mathbf{x})^G$  for which  $V_0 \subset \text{Dom} f_i$ ,  $i = 1, 2, \dots, k$  and  $V_0$  is dense in  $V$ , separates all orbits in  $V_0$  does it imply that the field extension  $F(f_1(x), f_2(x), \dots, f_k(x)) \subset F(\mathbf{x})^G$  is an algebraic extension? In other words in this case is it true that*

$$\text{trdeg} F(f_1(x), f_2(x), \dots, f_k(x))/F = \text{trdeg} F(\mathbf{x})^G/F?$$

With relation to Corollary 2.7 the following natural problem can be considered as well.

**Problem 2.10.** *Under the Assumption for  $G = GL(m, F)$  is it true that  $F \subset F(\mathbf{x})^{GL(m, F)}$  is also a pure transcendental extension?*

### 3. Classification of algebras

#### 3.1. General case

In this paper we use the standard notation (the Einstein notation) for tensors as well as the matrix representation for tensors which is more convenient in dealing with equivalence and invariance problems of tensors with respect to basis changes. The use of matrix representation for tensors makes the descriptions more transparent as well. Further the classification problem is considered only with respect to the general linear group.

Let us consider any  $m$  dimensional algebra  $W$  with multiplication  $\cdot$  given by a bilinear map  $(\mathbf{u}, \mathbf{v}) \mapsto \mathbf{u} \cdot \mathbf{v}$ . If  $e = (e^1, e^2, \dots, e^m)$  is a basis for  $W$  then one can represent the bilinear map by a matrix  $A \in Mat(m \times m^2; F)$  such that

$$\mathbf{u} \cdot \mathbf{v} = eA(u \otimes v)$$

for any  $\mathbf{u} = eu, \mathbf{v} = ev$ , where  $u = (u_1, u_2, \dots, u_m), v = (v_1, v_2, \dots, v_m)$  are column vectors. So the algebra  $W$  (binary operation, bilinear map, tensor) is presented by the matrix  $A \in Mat(m \times m^2; F)$ -the matrix of structure constants of  $W$  with respect to the basis  $e$ . Further we deal only with such matrices of rank  $m$ .

If  $e' = (e'^1, e'^2, \dots, e'^m)$  is also a basis for  $W$ ,  $g \in G = GL(m, F)$ ,  $e'g = e$  and  $\mathbf{u} \cdot \mathbf{v} = e'B(u' \otimes v')$ , where  $\mathbf{u} = e'u', \mathbf{v} = e'v'$ , then

$$\mathbf{u} \cdot \mathbf{v} = eA(u \otimes v) = e'B(u' \otimes v') = eg^{-1}B(gu \otimes gv) = eg^{-1}B(g \otimes g)(u \otimes v)$$

as far as  $\mathbf{u} = eu = e'u' = eg^{-1}u', \mathbf{v} = ev = e'v' = eg^{-1}v'$ . Therefore the equality

$$B = gA(g^{-1})^{\otimes 2} \quad (3.1)$$

is valid.

Now let  $\tau$  stand for the representation of  $G = GL(m, F)$  on the  $n = m^3$  dimensional vector space  $V = Mat(m \times m^2; F)$  defined by

$$\tau : (g, A) \mapsto B = gA(g^{-1} \otimes g^{-1}).$$

To have **Theorems 2.1, 2.3, 2.5** for this case we will construct a map  $P : V_0 \rightarrow GL(m, F)$  with property (2.1) in the following way. For any natural number  $k$  due to (3.1) one has

$$B^{\otimes k} = g^{\otimes k} A^{\otimes k} (g^{-1})^{\otimes 2k} \quad (3.2)$$

Let us consider all its possible contractions with respect to  $k$  upper and  $k$  lower indexes. It is clear that the result of each of such contraction will be  $f(B) = f(A)(g^{-1})^{\otimes k}$  type equality, where  $f(A)$  is a row vector with  $m^k$  entries.

In  $k = 1$  case one gets the following  $2^1 1! = 2$  different row equalities:  $\mathbf{Tr}_1(B) = \mathbf{Tr}_1(A)g^{-1}$ ,  $\mathbf{Tr}_2(B) = \mathbf{Tr}_2(A)g^{-1}$ , where  $\mathbf{Tr}_1(A)$  stands for the row vector with entries  $A_{j,i}^j = \sum_{j=1}^n A_{j,i}^j$ - the contraction on the first upper and lower indexes and  $\mathbf{Tr}_2(A)$  stands for the row vector with entries  $A_{i,j}^j = \sum_{j=1}^n A_{i,j}^j$ - the contraction on the first upper and second lower indexes

In  $k = 2$  case one gets the following  $2^2 2! + 2^1 1! = 10$  different row equalities:

$$\mathbf{Tr}_i(B) \otimes \mathbf{Tr}_j(B) = (\mathbf{Tr}_i(A) \otimes \mathbf{Tr}_j(A))(g^{-1})^{\otimes 2}, \quad \mathbf{Tr}_i(B)B = \mathbf{Tr}_i(A)A(g^{-1})^{\otimes 2},$$

where  $i, j = 1, 2$ , and

$$(B_{j,p}^i B_{i,q}^j) = (A_{j,p}^i A_{i,q}^j)(g^{-1})^{\otimes 2}, \quad (B_{j,p}^i B_{q,i}^j) = (A_{j,p}^i A_{q,i}^j)(g^{-1})^{\otimes 2},$$

$$(B_{p,j}^i B_{i,q}^j) = (A_{p,j}^i A_{i,q}^j)(g^{-1})^{\otimes 2}, \quad (B_{p,j}^i B_{q,i}^j) = (A_{p,j}^i A_{q,i}^j)(g^{-1})^{\otimes 2}.$$

In any  $k$  case only the number of contractions of  $A^{\otimes k}$  when all  $k$  different upper indexes are contracted with lower indexes of different  $A$  is

$$(2k) \times (2(k-1)) \times (2(k-2)) \times \dots \times 2 = 2^k k!.$$

In general it is nearly clear that the corresponding resulting system of  $2^k k!$  rows depending on the variable matrix  $A := \mathbf{x} = (x_{j,k}^i)_{i,j,k=1,2,\dots,m}$  is linear independent over  $F$ . But for big enough  $k$  the inequality  $2^k k! \geq m^k$  holds true as well. Therefore in general for big enough  $k$  it is possible to choose  $m^k$  contractions (rows) among the all contractions of  $\mathbf{x}^{\otimes k}$  for which the matrix  $Q(\mathbf{x})$  consisting of these  $m^k$  rows is a nonsingular matrix. For the matrix  $Q(\mathbf{x})$  one has equality  $Q(\mathbf{y}) = Q(\mathbf{x})(g^{-1})^{\otimes k}$  whenever  $g \in G$ ,  $\mathbf{y} = g\mathbf{x}(g^{-1})^{\otimes 2}$ .

Now note that for any  $A \in \{\mathbf{x} : \det(Q(\mathbf{x})) \neq 0\}$  and  $g \in G$  one has, for example,

$$(B \otimes (\mathbf{Tr}_1(B))^{\otimes k-2})Q(B)^{-1} =$$

$$g(A \otimes (\mathbf{Tr}_1(A))^{\otimes k-2})(g^{-1})^{\otimes k}(Q(A)(g^{-1})^{\otimes k})^{-1} = g(A \otimes (\mathbf{Tr}_1(A))^{\otimes k-2})Q(A)^{-1}.$$

Therefore if  $P(A)^{-1}$  stands for arbitrary nonsingular  $m \times m$  size sub-matrix of

$$(A \otimes (\mathbf{Tr}_1(A))^{\otimes k-2})Q(A)^{-1}$$

then one has the equality

$$P(B)^{-1} = gP(A)^{-1},$$

where  $g \in G$ ,  $B = gA(g^{-1})^{\otimes 2}$ . It implies that whenever  $A \in V_0 = \{A : \det(P(A)) \det(Q(A)) \neq 0\}$  the equality  $P(B) = P(A)g^{-1}$  holds true for any  $g \in G$  and  $B = gA(g^{-1})^{\otimes 2}$ . Note that

$$V_0 = \{A : \det(P(A)) \det(Q(A)) \neq 0\}$$

is a  $G$ -invariant, open and dense subset of  $V$ .

Therefore we have the following classification theorem for algebras whose matrices of structure constants are in  $V_0$ .

**Theorem 3.1.** *Two algebras with matrices of structure constants  $A, B \in V_0$  are same (isomorphic) algebras if and only if*

$$P(A)A(P(A)^{-1} \otimes P(A)^{-1}) = P(B)B(P(B)^{-1} \otimes P(B)^{-1}).$$

This result means that for any algebra  $W$ , given by a matrix of structure constants  $A \in V_0$ , for the canonical representative of algebras isomorphic to  $W$  one can take the algebra whose matrix of structure constants is given by  $P(A)A(P(A)^{-1} \otimes P(A)^{-1})$ .

**Theorem 3.2.** *The field of  $G$ -invariant rational functions  $F(\mathbf{x})^G$  of structure constants defined by variable matrix  $\mathbf{x} = (x_{j,k}^i)_{i,j,k=1,2,\dots,m}$  is generated by the system of entries of  $P(\mathbf{x})\mathbf{x}(P(\mathbf{x})^{-1} \otimes P(\mathbf{x})^{-1})$  over  $F$ , that is the equality*

$$F(\mathbf{x})^G = F(P(\mathbf{x})\mathbf{x}(P(\mathbf{x})^{-1} \otimes P(\mathbf{x})^{-1}))$$

holds true.

**Theorem 3.3.** *The transcendence degree of  $F(\mathbf{x})^G$  over  $F$  equals to  $m^3 - m^2$  and the field extension  $F(\mathbf{x})^G \subset F(\mathbf{x})$  is a pure transcendental extension.*

**Remark 3.4.** *One of the main results (Theorem 1) of [4] states that the field extension  $F \subset F(\mathbf{x})^{GL(m,F)}$  is a pure transcendental extension, which so far we could not get by our approach. Theorem 3.3 can be considered as a complementary result to that Theorem 1.*

Now let us consider two and three dimensional algebra cases.

**Example 3.5.** *Two dimensional ( $m = 2$ ) case. Let*

$$A = \begin{pmatrix} A_{1,1}^1 & A_{1,2}^1 & A_{2,1}^1 & A_{2,2}^1 \\ A_{1,1}^2 & A_{1,2}^2 & A_{2,1}^2 & A_{2,2}^2 \end{pmatrix}$$

*be a matrix of structure constants with respect to a basis. In this case at  $k = 1$  already  $2^1 1! = m^1$  and therefore for the rows of  $P(A)$  on can take*

$$\mathbf{Tr}_1(A) = (A_{1,1}^1 + A_{2,1}^2, A_{1,2}^1 + A_{2,2}^2) \text{ and } \mathbf{Tr}_2(A) = (A_{1,1}^1 + A_{1,2}^2, A_{2,1}^1 + A_{2,2}^2)$$

*and  $V_0$  consists of all  $A$  for which*

$$\det P(A) = (A_{1,1}^1 + A_{2,1}^2)(A_{2,1}^1 + A_{2,2}^2) - (A_{1,2}^1 + A_{2,2}^2)(A_{1,1}^1 + A_{1,2}^2) \neq 0.$$

*To see matrix of structure constants of the corresponding canonical algebra one should evaluate  $P(A)A(P(A)^{-1})^{\otimes 2}$ . System of entries of  $P(\mathbf{x})\mathbf{x}(P(\mathbf{x})^{-1})^{\otimes 2}$  presents a system of generators for the corresponding field of invariants rational functions, where  $\mathbf{x} = \begin{pmatrix} x_{1,1}^1 & x_{1,2}^1 & x_{2,1}^1 & x_{2,2}^1 \\ x_{1,1}^2 & x_{1,2}^2 & x_{2,1}^2 & x_{2,2}^2 \end{pmatrix}$ .*

On full classification of two dimensional algebras one can see [1].

**Example 3.6.** *Three dimensional ( $m = 3$ ) case. Let*

$$A = \begin{pmatrix} A_{1,1}^1 & A_{1,2}^1 & A_{1,3}^1 & A_{2,1}^1 & A_{2,2}^1 & A_{2,3}^1 & A_{3,1}^1 & A_{3,2}^1 & A_{3,3}^1 \\ A_{1,1}^2 & A_{1,2}^2 & A_{1,3}^2 & A_{2,1}^2 & A_{2,2}^2 & A_{2,3}^2 & A_{3,1}^2 & A_{3,2}^2 & A_{3,3}^2 \\ A_{1,1}^3 & A_{1,2}^3 & A_{1,3}^3 & A_{2,1}^3 & A_{2,2}^3 & A_{2,3}^3 & A_{3,1}^3 & A_{3,2}^3 & A_{3,3}^3 \end{pmatrix}$$

*be a matrix of the structure constants with respect to a basis.*

*In this case at  $k = 1$  one has  $2^1 1! < 3^1$ . At  $k = 2$  already  $2^2 2! + 2^1 1! = 10 > 3^2$  and the following 10 equalities*

$$\mathbf{Tr}_i(B) \otimes \mathbf{Tr}_j(B) = \mathbf{Tr}_i(A) \otimes \mathbf{Tr}_j(A)(g^{-1})^{\otimes 2}, \quad \mathbf{Tr}_i(B)B = \mathbf{Tr}_i(A)A(g^{-1})^{\otimes 2},$$

*where  $i, j = 1, 2$ ,*

$$(B_{j,p}^i B_{i,q}^j) = (A_{j,p}^i A_{i,q}^j)(g^{-1})^{\otimes 2}, \quad (B_{j,p}^i B_{q,i}^j) = (A_{j,p}^i A_{q,i}^j)(g^{-1})^{\otimes 2},$$

$$(B_{p,j}^i B_{i,q}^j) = (A_{p,j}^i A_{i,q}^j)(g^{-1})^{\otimes 2}, \quad (B_{p,j}^i B_{q,i}^j) = (A_{p,j}^i A_{q,i}^j)(g^{-1})^{\otimes 2}$$

*hold true.*

*Therefore for  $Q(A)$ , for example, one can take the following matrix*

$$Q(A) = \begin{pmatrix} A_{i,1}^i A_{j,1}^j & A_{i,1}^i A_{j,2}^j & A_{i,1}^i A_{j,3}^j & A_{i,2}^i A_{j,1}^j & A_{i,2}^i A_{j,2}^j \\ A_{i,1}^i A_{1,j}^j & A_{i,1}^i A_{2,j}^j & A_{i,1}^i A_{3,j}^j & A_{i,2}^i A_{1,j}^j & A_{i,2}^i A_{2,j}^j \\ A_{1,i}^i A_{j,1}^j & A_{1,i}^i A_{j,2}^j & A_{1,i}^i A_{j,3}^j & A_{2,i}^i A_{j,1}^j & A_{2,i}^i A_{j,2}^j \\ A_{1,i}^i A_{1,j}^j & A_{1,i}^i A_{2,j}^j & A_{1,i}^i A_{3,j}^j & A_{2,i}^i A_{1,j}^j & A_{2,i}^i A_{2,j}^j \\ A_{i,j}^i A_{1,1}^j & A_{i,j}^i A_{1,2}^j & A_{i,j}^i A_{1,3}^j & A_{i,j}^i A_{2,1}^j & A_{i,j}^i A_{2,2}^j \\ A_{j,i}^i A_{1,1}^j & A_{j,i}^i A_{1,2}^j & A_{j,i}^i A_{1,3}^j & A_{j,i}^i A_{2,1}^j & A_{j,i}^i A_{2,2}^j \\ A_{j,1}^i A_{i,1}^j & A_{j,1}^i A_{i,2}^j & A_{j,1}^i A_{i,3}^j & A_{j,2}^i A_{i,1}^j & A_{j,2}^i A_{i,2}^j \\ A_{j,1}^i A_{1,i}^j & A_{j,1}^i A_{2,i}^j & A_{j,1}^i A_{3,i}^j & A_{j,2}^i A_{1,i}^j & A_{j,2}^i A_{2,i}^j \\ A_{1,j}^i A_{i,1}^j & A_{1,j}^i A_{i,2}^j & A_{1,j}^i A_{i,3}^j & A_{2,j}^i A_{i,1}^j & A_{2,j}^i A_{i,2}^j \\ A_{i,2}^i A_{j,3}^j & A_{i,3}^i A_{j,1}^j & A_{i,3}^i A_{j,2}^j & A_{i,3}^i A_{j,3}^j \\ A_{i,2}^i A_{3,j}^j & A_{i,3}^i A_{1,j}^j & A_{i,3}^i A_{2,j}^j & A_{i,3}^i A_{3,j}^j \\ A_{2,i}^i A_{j,3}^j & A_{3,i}^i A_{j,1}^j & A_{3,i}^i A_{j,2}^j & A_{3,i}^i A_{j,3}^j \\ A_{2,i}^i A_{3,j}^j & A_{3,i}^i A_{1,j}^j & A_{3,i}^i A_{2,j}^j & A_{3,i}^i A_{3,j}^j \\ A_{i,j}^i A_{2,3}^j & A_{i,j}^i A_{3,1}^j & A_{i,j}^i A_{3,2}^j & A_{i,j}^i A_{3,3}^j \\ A_{j,i}^i A_{2,3}^j & A_{j,i}^i A_{3,1}^j & A_{j,i}^i A_{3,2}^j & A_{j,i}^i A_{3,3}^j \\ A_{j,2}^i A_{i,3}^j & A_{j,3}^i A_{i,1}^j & A_{j,3}^i A_{i,2}^j & A_{j,3}^i A_{i,3}^j \\ A_{j,2}^i A_{3,i}^j & A_{j,3}^i A_{1,i}^j & A_{j,3}^i A_{2,i}^j & A_{j,3}^i A_{3,i}^j \\ A_{2,j}^i A_{i,3}^j & A_{3,j}^i A_{i,1}^j & A_{3,j}^i A_{i,2}^j & A_{3,j}^i A_{i,3}^j \end{pmatrix}.$$

For  $P(A)^{-1}$  one can take any  $3 \times 3$  size nonsingular sub-matrix of  $AQ(A)^{-1}$ .

### 3.2. Commutative and anti-commutative algebra cases

For the classification purpose instead of all  $m$  dimensional algebras one can consider only such commutative or anti-commutative algebras. The commutativity (anti-commutativity) of the binary operation in terms of the corresponding matrix  $A$  means  $A_{j,k}^i = A_{k,j}^i$  (respectively,  $A_{j,k}^i = -A_{k,j}^i$ ) for all  $i, j, k = 1, 2, \dots, m$ . So in commutative (anti-commutative) algebra case for the  $V$  we consider  $V =$

$$\{A \in Mat(m \times m^2; F) : A_{j,k}^i = A_{k,j}^i (\text{resp. } A_{j,k}^i = -A_{k,j}^i) \text{ for all } i, j, k = 1, 2, \dots, m.\}$$

Note that in commutative (anti-commutative) case the dimension of  $V$  is  $\frac{m^2(m+1)}{2}$  (respectively,  $\frac{m^2(m-1)}{2}$ ).

To have **Theorems 2.1, 2.2, 2.5** for these cases one can construct a map  $P : V_0 \rightarrow GL(m, F)$  with property (2.1) in a similar way as in the general algebra case. Consider once again equality (3.2) and all its possible contractions with respect to  $k$  upper and  $k$  lower indexes.

In commutative (anti-commutative) case at  $k = 1$  one gets the following  $1! = 1$  row equality:  $\text{Tr}_1(B) = \text{Tr}_1(A)g^{-1} = \text{Tr}_2(A)g^{-1}$  as far as  $A_{j,k}^i = A_{k,j}^i$  (respectively,  $\text{Tr}_1(B) = \text{Tr}_1(A)g^{-1} = -\text{Tr}_2(A)g^{-1}$  as far as  $A_{j,k}^i = -A_{k,j}^i$ ) for all  $i, j, k = 1, 2, \dots, m$ .

In  $k = 2$  case one gets the following  $2! + 1! = 3$  different row equalities:

$$\text{Tr}_1(B) \otimes \text{Tr}_1(B) = (\text{Tr}_1(A) \otimes \text{Tr}_1(A))(g^{-1})^{\otimes 2},$$

$$\text{Tr}_1(B)B = \text{Tr}_1(A)A(g^{-1})^{\otimes 2}, \quad (B_{j,p}^i B_{i,q}^j) = (A_{j,p}^i A_{i,q}^j)(g^{-1})^{\otimes 2}.$$

In any  $k$  case only the number of contractions of  $A^{\otimes k}$  when all  $k$  different upper indexes are contracted with lower indexes of different  $A$  is  $k!$ . Once again in general it is nearly clear



that the corresponding resulting system of  $k!$  rows depending on variable matrix  $A := \mathbf{x} = (x_{j,k}^i)_{i,j,k=1,2,\dots,m}$ , where  $x_{j,k}^i = x_{k,j}^i$  (respectively,  $x_{j,k}^i = -x_{k,j}^i$ ) for all  $i, j, k = 1, 2, \dots, m$ , is linear independent over  $F$ . But for big enough  $k$  the inequality  $k! \geq m^k$  holds true as well. Therefore in general for big enough  $k$  it is possible to choose  $m^k$  contractions (rows) among the all contractions of  $\mathbf{x}^{\otimes k}$  for which the matrix  $Q(\mathbf{x})$  consisting of these  $m^k$  rows is nonsingular. For the matrix  $Q(\mathbf{x})$  one has equality  $Q(\mathbf{y}) = Q(\mathbf{x})(g^{-1})^{\otimes k}$  whenever  $g \in G$ ,  $\mathbf{y} = g\mathbf{x}(g^{-1})^{\otimes 2}$ .

Therefore if  $P(A)^{-1}$  stands for arbitrary  $m \times m$ -size nonsingular sub-matrix of  $(A \otimes (\text{Tr}_1(A))^{\otimes k-2})Q(A)^{-1}$  then one has the equality  $P(B)^{-1} = gP(A)^{-1}$ , where  $g \in G$ ,  $B = gA(g^{-1})^{\otimes 2}$ . It implies that whenever  $A \in V_0 = \{A \in V : \det(P(A)) \det(Q(A)) \neq 0\}$  the equality  $P(B) = P(A)g^{-1}$  holds true for any  $g \in G$ , where  $B = gA(g^{-1})^{\otimes 2}$ . Note that

$$V_0 = \{A \in V : \det(P(A)) \det(Q(A)) \neq 0\}$$

is a  $G$ -invariant, open and dense subset of  $V$ .

Therefore for such algebras one can state the following results.

**Theorem 3.7.** *Two commutative (anti-commutative) algebras with the matrices of structure constants  $A, B \in V_0$  are the same algebras if and only if*

$$P(A)A(P(A)^{-1} \otimes P(A)^{-1}) = P(B)B(P(B)^{-1} \otimes P(B)^{-1}).$$

**Theorem 3.8.** *The field of  $G$ -invariant rational functions  $F(\mathbf{x})^G$  of the structure constants presented by the matrix  $\mathbf{x} = ((x_{j,k}^i)_{i,j,k=1,2,\dots,m})$  of the variable commutative (respectively, anti-commutative) algebras, where  $x_{j,k}^i = x_{k,j}^i$  (respectively,  $x_{j,k}^i = -x_{k,j}^i$ ) for all  $i, j, k = 1, 2, \dots, m$ , is generated by the system of entries of  $P(\mathbf{x})\mathbf{x}(P(\mathbf{x})^{-1} \otimes P(\mathbf{x})^{-1})$  over  $F$ , that is the equality*

$$F(\mathbf{x})^G = F(P(\mathbf{x})\mathbf{x}(P(\mathbf{x})^{-1} \otimes P(\mathbf{x})^{-1}))$$

holds true.

**Theorem 3.9.** *In commutative (anti-commutative) algebra case the transcendence degree of  $F(\mathbf{x})^G$  over  $F$  equals to  $\frac{m^2(m-1)}{2}$  (respectively,  $\frac{m^2(m-3)}{2}$ ,  $m \geq 3$ ) and the field extension  $F(\mathbf{x})^G \subset F(\mathbf{x})$  is a pure transcendental extension.*

## Acknowledgments

This research is supported by the MOE of Malaysia under grant FRGS14-153-0394.

## References

- [1] Michel G and Elisabeth R 2011 *African J. Math. Phys.* **10** 81–91
- [2] Durán D R et al 2003 *Lin. Alg. Appl.* **364** 1–12
- [3] Hernández Encinas L et al 2004 *Lin. Alg. Appl.* **387** 69–82
- [4] Popov V 2011 *arXiv: 1411.6570*
- [5] Bekbaev U 2016 *Jour. of Phys.: Conf. Ser.* **697** 012005