

Web of Science

Search

Search Results

My Tools

Search History

Marked List

 Look Up Full Text


Save to EndNote online

Add to Marked List

1 of 1

Thyroxine (T-4) Transfer from Blood to Cerebrospinal Fluid in Sheep Isolated Perfused Choroid Plexus: Role of Multidrug Resistance-Associated Proteins and Organic Anion Transporting Polypeptides

By: Zibara, K (Zibara, Kazem)^[1,2]; El Zein, N (El Zein, Nabil)^[1,2]; Sabra, M (Sabra, Mirna)^[1,3]; Hneino, M (Hneino, Mohammad)^[1,4]; Harati, H (Harati, Hayat)^[3]; Mohamed, W (Mohamed, Wael)^[5,6]; Kobeissy, FH (Kobeissy, Firas H.)^[7]; Kassem, N (Kassem, Nouhad)^[1,3]

[View ResearcherID and ORCID](#)

FRONTIERS IN NEUROLOGY

Volume: 8

Article Number: 214

DOI: 10.3389/fneur.2017.00214

Published: MAY 23 2017

[View Journal Impact](#)

Abstract

Thyroxine (T-4) enters the brain either directly across the blood-brain barrier (BBB) or indirectly via the choroid plexus (CP), which forms the blood-cerebrospinal fluid barrier (B-CSF-B). In this study, using isolated perfused CP of the sheep by single-circulation paired tracer and steady-state techniques, T4 transport mechanisms from blood into lateral ventricle CP has been characterized as the first step in the transfer across the B-CSF-B. After removal of sheep brain, the CPs were perfused with I-125-T-4 and C-14-mannitol. Unlabeled T-4 was applied during single tracer technique to assess the mode of maximum uptake (U-max) and the net uptake (U-net) on the blood side of the CP. On the other hand, in order to characterize T-4 protein transporters, steady-state extraction of I-125-T-4 was measured in presence of different inhibitors such as probenecid, verapamil, BCH, or indomethacin. Increasing the concentration of unlabeled-T-4 resulted in a significant reduction in U-max%, which was reflected by a complete inhibition of T-4 uptake into CP. In fact, the obtained U-net% decreased as the concentration of unlabeled-T-4 increased. The addition of probenecid caused a significant inhibition of T-4 transport, in comparison to control, reflecting the presence of a carrier mediated process at the basolateral side of the CP and the involvement of multidrug resistance-associated proteins (MRPs: MRP1 and MRP4) and organic anion transporting polypeptides (Oatp1, Oatp2, and Oatp14). Moreover, verapamil, the P-glycoprotein (P-gp) substrate, resulted in similar to 34% decrease in the net extraction of T-4, indicating that MDR1 contributes to T-4 entry into CSF. Finally, inhibition in the net extraction of T-4 caused by BCH or indomethacin suggests, respectively, a role for amino acid "L" system and MRP1/Oatp1 in mediating T-4 transfer. The presence of a carrier-mediated transport mechanism for cellular uptake on the basolateral membrane of the CP, mainly P-gp and Oatp2, would account for the efficient T-4 transport from blood to CSF. The current study highlights a carrier-mediated transport mechanism for T4 movement from blood to brain at the basolateral side of B-CSF-B/CP, as an alternative route to BBB.

Keywords

Author Keywords: transport; thyroid hormone; blood-cerebrospinal fluid barrier; blood-brain barrier; efflux; uptake

KeyWords Plus: THYROID-HORMONE TRANSPORTERS; CARRIER-MEDIATED TRANSPORT; BRAIN-BARRIER; P-GLYCOPROTEIN; RAT-BRAIN; PRIMARY CULTURE; LEAD-EXPOSURE; CELLS; TRANSTHYRETIN; SYSTEM

Author Information

Citation Network

0 Times Cited

[60 Cited References](#)
[View Related Records](#)
 [Create Citation Alert](#)
(data from Web of Science Core Collection)

All Times Cited Counts

0 in All Databases

0 in Web of Science Core Collection

0 in BIOSIS Citation Index

0 in Chinese Science Citation Database

0 in Data Citation Index

0 in Russian Science Citation Index

0 in SciELO Citation Index

Usage Count

Last 180 Days: 1

Since 2013: 1

[Learn more](#)

This record is from:

Web of Science Core Collection
- Science Citation Index Expanded

Suggest a correction

If you would like to improve the quality of the data in this record, please [suggest a correction](#).

Reprint Address: Zibara, K; Kassem, N (reprint author)

+ Lebanese Univ, ER045, PRASE, Beirut, Lebanon.

Reprint Address: Zibara, K (reprint author)

+ Lebanese Univ, Biol Dept, Fac Sci, Beirut, Lebanon.

Reprint Address: Kassem, N (reprint author)

+ Lebanese Univ, Neurosci Res Ctr, Fac Med Sci, Beirut, Lebanon.

Reprint Address: Kobeissy, FH (reprint author)

+ Amer Univ Beirut, Dept Biochem & Mol Genet, Beirut, Lebanon.

Addresses:

+ [1] Lebanese Univ, ER045, PRASE, Beirut, Lebanon

+ [2] Lebanese Univ, Biol Dept, Fac Sci, Beirut, Lebanon

+ [3] Lebanese Univ, Neurosci Res Ctr, Fac Med Sci, Beirut, Lebanon

+ [4] Lebanese Univ, Fac Publ Hlth, Med Lab Dept, Beirut, Lebanon

+ [5] Int Islamic Univ Malaysia, Kulliyah Med, Basic Med Sci Dept, Kuantan, Pahang, Malaysia

[6] Menoufia Med Sch, Neurosci Unit, Cairo, Egypt

+ [7] Amer Univ Beirut, Dept Biochem & Mol Genet, Beirut, Lebanon

E-mail Addresses: kzibara@ul.edu.lb; firasko@gmail.com; nouhad.kassem@hotmail.com

Publisher

FRONTIERS MEDIA SA, PO BOX 110, EPFL INNOVATION PARK, BUILDING I, LAUSANNE, 1015, SWITZERLAND

Categories / Classification

Research Areas: Neurosciences & Neurology

Web of Science Categories: Clinical Neurology; Neurosciences

Document Information

Document Type: Article

Language: English

Accession Number: WOS:000401944200001

PubMed ID: 28588548

ISSN: 1664-2295

Journal Information

Impact Factor: [Journal Citation Reports](#)

Other Information

IDS Number: EV7GP

Cited References in Web of Science Core Collection: 60

Times Cited in Web of Science Core Collection: 0