

Document details

[< Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)[Full Text](#) [View at Publisher](#)IIUM Engineering Journal
Volume 19, Issue 1, 2018, Pages 15-23

Investigation of bioflocculant as dewatering aid in sludge treatment

(Article)

Jami, M.S. [✉](#), Mel, M., Ariff, A.R.M., Abdulazeez, Q.M. [👤](#)

Department of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, Jalan Gombak, Kuala Lumpur, Malaysia

Abstract

[View references \(15\)](#)

Sludge treatment is one of the most important and expensive steps in water and wastewater treatment plants. Chemical conditioners such as polyaluminum chloride, aluminum sulfate, Fenton's reagent, gypsum, and polyacrylamide can produce byproducts that cause health and environmental problems. *Moringa oleifera* (MO) seeds can be used as a natural alternative to chemical conditioners. The bioactive materials have to be extracted from MO seeds for better performance. In this study, the treatment methods of MO seeds were the bioactive extraction by NaCl (1 M) and oil extraction by hexane solvent, as well as the untreated (crude) seeds powder. Synthetic sludge samples were prepared using kaolin suspension (5% w/v). The most effective coagulant-form was determined based on the values of settling velocity (V_s) and sludge volume index (SVI). Results showed that extraction by NaCl gave the best results of 0.41 cm/min of settling velocity and 63.39 ml/g of SVI. A SVI value greater than 150 ml/g indicates poor settling qualities whereas the control sludge of the current study was 100 ml/g. The most effective coagulant-form was optimized with respect to three process conditions: MO seeds dosage, mixing speed, and contact time. The experiments were designed using 2 Level Factorial-Design by Design-Expert software. The optimum process conditions were seeds dosage of 3246 mg/l, mixing speed of 102 rpm, and mixing time of 29 min. MO seeds can be considered as a natural coagulant that can be used as main sludge conditioner. © 2018 International Islamic University Malaysia-IIUM.

Author keywords

[Dewatering](#) [Environmental friendly](#) [Moringa oleifera](#) [Sludge treatment](#)

ISSN: 1511788X

Source Type: Journal

Original language: English

DOI: 10.31436/iiumej.v19i1.735

Document Type: Article

Publisher: International Islamic University Malaysia-IIUM

References (15)

[View in search results format >](#) All [Export](#) [Print](#) [E-mail](#) [Save to PDF](#) [Create bibliography](#)

- 1 Henderson, J.P., Heinecke, J.W.
Myeloperoxidase and eosinophil peroxidase: Phagocyte enzymes for halogenation in humans

(2003) *Handbook of Environmental Chemistry*, Part P 3, pp. 201-214. Cited 7 times.<http://www.springer.com/series/698>

doi: 10.1007/b10448

[View at Publisher](#)Metrics [?](#)

0 Citations in Scopus

0 Field-Weighted

Citation Impact

PlumX Metrics [v](#)

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

Related documents

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

- 2 Verlicchi, P., Zambello, E.
Pharmaceuticals and personal care products in untreated and treated sewage sludge: Occurrence and environmental risk in the case of application on soil - A critical review
(2015) *Science of the Total Environment*, 538, pp. 750-767. Cited 89 times.
www.elsevier.com/locate/scitotenv
doi: 10.1016/j.scitotenv.2015.08.108

View at Publisher
-
- 3 Abdul Hamid, S.H., Lananan, F., Khatoon, H., Jusoh, A., Endut, A.
A study of coagulating protein of *Moringa oleifera* in microalgae bio-flocculation
(2016) *International Biodeterioration and Biodegradation*, 113, pp. 310-317. Cited 7 times.
www.elsevier.com/inca/publications/store/4/0/5/8/9/9
doi: 10.1016/j.ibiod.2016.03.027

View at Publisher
-
- 4 LeBlanc, R.J., Matthews, P., Richard, R.P.
(2008) *Global atlas of excreta*
wastewater sludge, and biosolids management: moving forward the sustainable and welcome uses of a global resource
-
- 5 Bichi, M.H.
A review of the applications of *Moringa oleifera* seeds extract in water treatment
(2013) *Civ. Environ. Res*, 3 (8), pp. 1-10. Cited 7 times.
-
- 6 Abdulazeez, Q.M., Jami, M.S., Alam, M.Z.
Effective sludge dewatering using *Moringa oleifera* seeds extract combined with aluminum sulphate
(2016) *ARNP Journal of Engineering and Applied Sciences*, 11 (1), pp. 372-381.
http://www.arnpjournals.org/jeas/research_papers/rp_2016/jeas_0116_3348.pdf
-
- 7 Gold, M., Dayer, P., Faye, M.C.A.S., Clair, G., Seck, A., Niang, S., Morgenroth, E., (...), Strande, L.
Locally produced natural conditioners for dewatering of faecal sludge
(2016) *Environmental Technology (United Kingdom)*, 37 (21), pp. 2802-2814. Cited 7 times.
<http://www.tandf.co.uk/journals/titles/09593330.asp>
doi: 10.1080/09593330.2016.1165293

View at Publisher
-
- 8 Kansal, S.K., Kumari, A.
Potential of *M. oleifera* for the treatment of water and wastewater
(2014) *Chemical Reviews*, 114 (9), pp. 4993-5010. Cited 39 times.
<http://pubs.acs.org/journal/chcreay>
doi: 10.1021/cr400093w

View at Publisher
-