

[< Back to results](#) | 1 of 1[↗ Export](#) [↓ Download](#) [🖨 Print](#) [✉ E-mail](#) [Save to PDF](#) [☆ Add to List](#) [More... >](#)Indian Journal of Natural Products and Resources
Volume 8, Issue 1, 2017, Pages 54-62

Open Access

Mimosa pudica L.: A comparative study via in vitro analysis and GC Q-TOF MS profiling on conventional and supercritical fluid extraction using food grade ethanol (Article)

Tasnuva, S.T.^a, Sahena, F.^b, Qamar, U.A.^a, Kashif, G.^c, Juliana, J.^a, Zaidul, I.S.M.^a [✉](#) [👤](#)^aFaculty of Pharmacy, International Islamic University Malaysia, Kuantan, Pahang DM, Malaysia^bFaculty of Science, International Islamic University Malaysia, Kuantan, Pahang DM, Malaysia^cDepartment of Food Science and Nutrition, King Saud University, Riyadh, Saudi Arabia

Abstract

[View references \(40\)](#)

The present study compared conventional (maceration) extraction (EtOHconv) and supercritical fluid extraction (SFE) methods as a mean of comparing conventional and green process for a weed namely *Mimosa pudica* L. to obtain a safe antidiabetic natural agent. In vitro analysis comprised of two different assays, antioxidant assay (determination of total phenolic content, total flavonoid content, and 2,2-diphenyl-1-picrylhydrazyl assay) and antidiabetic assay (inhibition of α -amylase and α -glucosidase). GC Q-TOF MS profiling for both extracts was done after derivetisation to confirm the presence of bioactive compounds. SFE was performed at 40 MPa pressure, 60 °C temperature and 5 mL/min CO₂ flow rate using 30 % ethanol (co-solvent) for 2 h. EtOHconv prepared using 95 % ethanol through conventional method (maceration) showed a good in vitro antioxidant potential and digestive enzymes inhibitory effect compared to supercritical fluid extract. α -amylase and α -glucosidase inhibitory activities for EtOHconv at 1 mg/mL were 30.08 % (± 5.23) and 38.29 % (± 2.52), whereas for standard acarbose it was 28.24 % (± 13.66) and 36.93 % (± 2.70), respectively. Supercritical fluid extract showed less potent in vitro antioxidant and digestive enzymes inhibitory effects (15.67 \pm 4.03- α -amylase, 28.36 \pm 2.01- α -glucosidase). GC Q-TOF MS analysis was done to confirm the presence of bioactive compounds in both the extracts. Although EtOHconv showed better results, SFE was found to contain more bioactive compounds associated with various pharmacological effects especially antioxidative as per GC Q-TOF MS results. SFE being a clean and green technology could be employed in future with more focus on method development and optimization to reproduce better and safe bioactive products from the neglected weed *M. pudica*. © 2017, National Institute of Science Communication and Information Resources (NISCAIR). All Rights Reserved.

Reaxys Database Information

[View Compounds](#)

Author keywords

Antioxidant Conventional extraction Derivetisation GC Q-TOF MS *Mimosa pudica* L SFE

ISSN: 09760504

Source Type: Journal

Original language: English

Document Type: Article

Publisher: National Institute of Science Communication and Information Resources (NISCAIR)

References (40)

[View in search results format >](#) All [Export](#) [🖨 Print](#) [✉ E-mail](#) [Save to PDF](#) [Create bibliography](#)

Metrics

0 Citations in Scopus

0 Field-Weighted
Citation Impact

PlumX Metrics

Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 0 documents

Inform me when this document
is cited in Scopus:

Related documents

Enrichment, in vitro, and
quantification study of
antidiabetic compounds from
neglected weed *Mimosa pudica*
using supercritical CO₂ and CO₂-
SohxhletTunna, T.S. , Sarker, M.Z.I. ,
Ghafoor, K.
(2018) *Separation Science and
Technology (Philadelphia)*Analyses and profiling of extract
and fractions of neglected weed
Mimosa pudica Linn. traditionally
used in Southeast Asia to treat
diabetesTunna, T.S. , Zaidul, I.S.M. ,
Ahmed, Q.U.
(2015) *South African Journal of
Botany*Erratum: L-glycine, a novel anti-
inflammatory,
immunomodulatory and
cytoprotective agent (Current
Opinion in Clinical Nutrition and
Metabolic Care (March 2003) 6
(229-240))

Zhong, Z. , Wheeler , Li, X.

- 1 Santana-Méridas, O., González-Coloma, A., Sánchez-Vioque, R.
Agricultural residues as a source of bioactive natural products

(2012) *Phytochemistry Reviews*, 11 (4), pp. 447-466. Cited 32 times.
doi: 10.1007/s11101-012-9266-0

[View at Publisher](#)

- 2 Conde, E., Reinoso, B.D., Gonzalez-Munoz, M.J., Andres Moure, A.
Recovery and concentration of antioxidants from industrial effluents and processing streams of underutilized vegetal biomass
(2013) *Food Pub Health*, 3, pp. 69-91. Cited 10 times.

- 3 Rout, S.P., Choudary, K.A., Kar, D.M., Das, L., Jain, A.
Plants in traditional medicinal system - Future source of new drugs

(2009) *International Journal of Pharmacy and Pharmaceutical Sciences*, (1), pp. 1-23. Cited 58 times.
<http://ijppsjournal.com/Vol%201%20Issue%201/101.pdf>

- 4 Ibañez, E., Herrero, M., Mendiola, J.A., Castro-Puyana, M.
Extraction and characterization of bioactive compounds with health benefits from marine resources: Macro and micro algae, cyanobacteria, and invertebrates

(2012) *Marine Bioactive Compounds: Sources, Characterization and Applications*, 9781461412472, pp. 55-98. Cited 81 times.

<http://dx.doi.org/10.1007/978-1-4614-1247-2>
ISBN: 978-146141247-2; 1461412463; 978-146141246-5
doi: 10.1007/978-1-4614-1247-2_2

[View at Publisher](#)

- 5 Moure, A., Cruz, J.M., Franco, D., Manuel Domínguez, J., Sineiro, J., Domínguez, H., Núñez, M.J., (...), Carlos Parajó, J.
Natural antioxidants from residual sources

(2001) *Food Chemistry*, 72 (2), pp. 145-171. Cited 965 times.
doi: 10.1016/S0308-8146(00)00223-5

[View at Publisher](#)

- 6 Chemat, F., Vian, M.A., Cravotto, G.
Green extraction of natural products: Concept and principles

(2012) *International Journal of Molecular Sciences*, 13 (7), pp. 8615-8627. Cited 236 times.

<http://www.mdpi.com/1422-0067/13/7/8615/pdf>
doi: 10.3390/ijms13078615

[View at Publisher](#)

- 7 Reverchon, E., De Marco, I.
Supercritical fluid extraction and fractionation of natural matter

(2006) *Journal of Supercritical Fluids*, 38 (2), pp. 146-166. Cited 560 times.
doi: 10.1016/j.supflu.2006.03.020

[View at Publisher](#)

- 8 (2014) *Weed Risk of Mimosa Pudica (Fabaceae)-Sensitive Plant*
Animal and Plant health Inspection Service, Raleigh, Version 1

(2003) *Current Opinion in Clinical Nutrition and Metabolic Care*

[View all related documents based on references](#)

[Find more related documents in Scopus based on:](#)

[Authors >](#) [Keywords >](#)