

[About IMJM](#) ▾[Submission Guidelines](#) ▾[Archives](#)[Contact us](#)[Home](#) › [Volume 17](#) › [IMJM Volume 17 Special Issue 1](#)

# IMJM Volume 17 Special Issue 1

📅 Posted on [July 18, 2018](#) | by  [Web Editor](#) | Posted in [Volume 17](#)

International Medical  
Journal Malaysia

Q3

Medicine  
(miscellaneous)

best quartile

SJR 2017

0.2

powered by scimagojr.com

## Previous Volumes

- [IMJM Volume 17 Special Issue 1](#)
- [Volume 17 No 1 \(April 2018\)](#)
- [Message from Chief Editor](#)
- [Volume 16 No 2 \(Dec 2017\)](#)
- [Volume 16 Supplementary Issue No 2](#)
- [Volume 16 Supplementary Issue No 1](#)
- [Volume 16 No 1 \(June 2017\)](#)
- [Volume 15 No 2 \(Dec 2016\)](#)
- [Volume 15 Supplementary Issue No 1](#)



Volume 1

# 2<sup>nd</sup> WORLD CONGRESS ON INTEGRATION AND ISLAMICISATION

FOCUS ON MEDICAL AND  
HEALTH CARE SCIENCES

21 – 23 OCTOBER 2016

[www.iium.edu.my/wcii](http://www.iium.edu.my/wcii)



In collaboration with  
Majlis Ugama Islam dan Adat Resam Melayu  
Pahang (MUIP)

- [Volume 15 No 1  
\(June 2016\)](#)

## Search IMJM article



## Related Links

- [The IIUM Official Website](#)
- [The Kulliyah of Medicine](#)

## EDITORIAL BOARD

### CHIEF EDITOR

Prof Dato Dr Tariq Abdul Razak  
MBBS, MCRP, M.Sc, MD, FRCP  
Department of Chairman Ibnu Sina of Chair Medicine  
Kulliyah of Medicine,  
International Islamic University Malaysia

### EDITORS

1. Assoc. Prof. Dr. Abdurezak Abdullah Hashi  
BA, MA, PhD  
Department of Biotechnonology

- Kulliyyah of Science  
International Islamic University Malaysia
2. Asst. Prof. Dr. Che Anuar Che Muhammad  
MBBS, MSc, PhD  
Department of Basic of Medical Sciences (Anatomy)  
Kulliyyah of Pharmacy  
International Islamic University Malaysia
3. Assoc. Prof Dr. Mohammad Arif Shahar  
MBBS (IIUM); MMed. (Int. Med.), UKM; Consultant Physician  
(Endocrinology)  
Department of Internal Medicine  
Kulliyyah of Medicine  
International Islamic University Malaysia
4. Asst. Prof. Dr. Intan Azura Shahdan  
BSc, MPhil, PhD  
Department of Biomedical Sciences  
Kulliyyah of Allied Health Science  
International Islamic University Malaysia
5. Assoc. Prof. Dr. Kamarul Ariffin Khalid  
DSc, MSur, MMedSc, LRCPL, & LCRI, MB, BCh, BAO  
Department of orthopedics  
Kulliyyah of Medicine  
International Islamic University Malaysia
6. Prof. Dr. Naseer Mohammad Amjad  
MBBS, MS, FCRS, DIS  
Department of Surgery  
Kulliyyah of Medicine  
International Islamic University Malaysia
7. Assoc. Prof. Dr. Aminuddin Che Ahmad  
MD, M Orth, HUG, TUM  
Department of Orthopedics, Traumatology & Rehabilitation  
Kulliyyah of Medicine  
International Islamic University Malaysia
8. Prof. Dato' Dr. Ariff Osman  
MBBS (Malaya). Master of Medicine (Anesthesiology), UKM  
Department of Anesthesiology & Intensive Care  
Kulliyyah of Medicine  
International Islamic University Malaysia

Show  entries Search:

Name	Size	Hits

Name	Size	Hits
<a href="#">A Case Report on Blood in Breastmilk: Safety Issues and Islamic Ruling</a>	438.2 KiB	45
<a href="#">A Pilot Survey on Knowledge, Attitude and Practice Towards the Islamisation of Knowledge As Perceived by the Staff of Kulliyah of Allied Health Sciences, International Islamic University Malaysia</a>	602.6 KiB	34
<a href="#">Addressing Islamic Values in Medical Imaging Professionalism</a>	350.1 KiB	33
<a href="#">Allah's Wondrous Creatures, the Holy Qur'an and Technological Inventions: Ultrasound Imaging</a>	923.3 KiB	34
<a href="#">An Overview of the Application of Maqasid Al-Shariah into Cartilage Tissue Engineering</a>	365.6 KiB	34
<a href="#">Anatomy of Musculoskeletal System in the Light of the Qur'an and Hadith</a>	696.9 KiB	25
<a href="#">Assessment of Prophetic Foods Consumption among Lactating Mothers: Combining Quantitative &amp; Qualitative Approaches</a>	303.0 KiB	33
<a href="#">Breastfeeding Videos as a New Training Tool for Malaysian Breastfeeding Peer Counsellor Program in Empowering Breastfeeding Communities</a>	283.8 KiB	33
<a href="#">Challenges on Learning Islam among Deaf Muslim in an Indonesian Context: Deaf Teachers Perspective</a>	345.2 KiB	24
<a href="#">Engaging and Empowering Parents of Children with Down Syndrome-The Role of Parents Support Group</a>	216.7 KiB	43

Showing 1 to 10 of 32 entries

Previous

1

2

3

4

Next

# Identifying the Potential of *Qur'anic* Recitation on the Proliferation of Chondrocytes Derived from Rabbit Articular Cartilage: *Work in Progress*

Rosyafirah Hashim<sup>1</sup>, Munirah Sha'ban<sup>2</sup>, Sarah Rahmat<sup>3</sup>, Zainul Ibrahim Zainuddin<sup>4</sup>

<sup>1</sup>Department of Biomedical Science, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan Pahang, <sup>2</sup>Department of Physical Rehabilitation Sciences, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan Pahang, <sup>3</sup>Department of Audiology and Speech-Language Pathology, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan Pahang, <sup>4</sup>Department of Diagnostic Imaging and Radiotherapy, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan Pahang.

## ABSTRACT

**Introduction:** In Islamic practice, the use of *Qur'anic* recitation in treatment can be traced back to the times of Prophet Muhammad (PBUH). This preliminary study aims to identify the potential of *Qur'anic* recitation of *Surah Al-Fatihah* on the proliferation of chondrocytes derived from rabbit articular cartilage. Cartilage tissue engineering offers an alternative way to facilitate cartilage regeneration *in-vitro*. **Materials and Methods:** The cellular model was established using a serially cultured and expanded chondrocytes *in-vitro*. The model was assigned into three groups. The first group was exposed to the *Surah Al-Fatihah*, recited 17 times based on the five times daily prayer unit (*Raka'ah*) obligated upon Muslims. The second group was exposed to an Arabic poem recitation. The third group was not exposed to any sound and served as the control. All groups were subjected to the growth profile analysis. The analysis was conducted at different passages starting from passage 0 to passage 3. **Results:** The results showed that the cells proliferation based on the growth kinetic analysis is higher for the cells exposed with *Qur'anic* recitation as compared to the Arabic poem and control groups. **Conclusions:** The proliferation process of the rabbit articular cartilage might be influenced with the use of *Qur'anic* recitation and as well as Arabic poem recitation. Exposure to the Western poem recitation and mute sound will be added for future study. It is hoped that this study could shed some light on the potential use of the *Qur'anic* recitation to facilitate cartilage regeneration in tissue engineering studies.

**KEYWORDS:** Qur'anic Recitation; Al-Fatihah; Sound Healing; Chondrocytes; Tissue Engineering

## INTRODUCTION

Articular cartilage or also known as hyaline cartilage is the thin, smooth, glistening white tissue that covers the surface of all diarthrodial joints in the human body.<sup>1</sup> The only cell that is made up of articular cartilage is the chondrocytes which will produce and maintain the large amount of extracellular matrix around the joint.<sup>2</sup> Although the articular cartilage is a metabolically active tissue but the chondrocytes themselves are lack of blood supply.<sup>2</sup> Therefore it has a very limited ability to repair itself whereby a minor injury may lead to progressive damage and degeneration of the articular cartilage.<sup>1</sup>

When the chondrocytes grow in monolayer cultures, they will proliferate and have a tendency to dedifferentiate.<sup>3</sup> After the chondrocytes attach to

the tissue culture plate, they begin to spread out, dividing, changing the gene expression and rapidly lose the chondrogenic phenotype over several passages in the culture process.<sup>4</sup> The expression of the specialized proteins will be delayed and the chondrocytes will become fibroblastic in appearance.<sup>3</sup> The chondrocytes cells will lose their rounded shape, experience a reduction in extracellular matrix synthesis, loss of chondrocytes genes expression and increase in expression of fibroblastic genes.<sup>4</sup>

The recitation of the Holy *Qur'an* is a special practice that is valued by Muslim across the world. In Islam, *Qur'anic* recitation has already been used as the method for treatment since the times of Prophet Muhammad (PBUH). It was reported that listening to *Qur'anic* recitation can have therapeutic effects as the growth hormone crucial for the tissues repair processes can be triggered.<sup>5</sup> The *Qur'anic* recitation is considered as an effective healing power because it will affect those who listened to it.<sup>6</sup> The type and frequency of the sound plays a very important consideration towards the healing power.<sup>6</sup> The effectiveness of the sound healing might be different depending on the range

Zainul Ibrahim Zainuddin  
Department of Diagnostic Imaging and Radiotherapy,  
Kulliyah of Allied Health Sciences,  
International Islamic University Malaysia,  
Kuantan Campus.  
Email: zainul@iiu.edu.my

and the level of the sound used during the healing process.<sup>7</sup>

There is a specific way to recite the Holy *Qur'an* as the reciters must know the proper '*makhraj*' and '*tajwid*' in order to read it correctly.<sup>8</sup> The authors also indicated that '*makhraj*' is the correct position of the organs of speech in order to produce a letter so that it can be differentiated from others. Meanwhile, '*tajwid*' is the correctness of diction or proper pronunciation and technique during recitation. The same authors also state that it is important to recite the Holy *Qur'an* according to the law of '*makhraj*' and '*tajwid*' because even a small difference of the sound in the Arabic words may cause the meaning to be different. Although the *Qur'anic* recitation will probably differ from a person to another due to the difference in each person's voice, the meaning will still be the same as long as the laws of '*makhraj*' and '*tajwid*' are obeyed.

In Islam, a Muslim is obliged to pray at least five times a day. One *hadith* narrated by al-Bukhari, Muslim and at-Tarmizi mentions that the recitation of *Surah Al-Fatihah* is compulsory in daily prayers for a Muslim as without it the prayer becomes invalid. This means that a Muslim recites these verses at least seventeen times a day during the prayers. A recording of the *Surah Al-Fatihah* by a *Qur'anic* reciter known to all Muslim scholars around the world was used in this study as the '*makhraj*' and '*tajwid*' for the recitation has already been accepted.

This study was aimed to identify the potential of *Qur'anic* recitation, particularly *Surah Al-Fatihah* on the proliferation of chondrocytes derived from the rabbit articular cartilage. It is believed that *Surah Al-Fatihah* is able to act as a healing medium. This is attributed to one *hadith* of Abu Saeed al-Khudri. *He narrated that some of the Companions of the Prophet (PBUH) went on a journey till they reached some Arab tribes at night. They asked to be treated as guests but the tribe refused. The chief of that tribe was then bitten by a poisonous animal (snake or scorpion) and nothing had benefited him. Some members of the tribe went to the group of the Companions to seek help. One of the companions agreed to recite Ruqya in return for a flock of sheep. The companion recited Surah Al-Fatihah to treat the Chief. All the praises and thanks to Allah, as the chief who was sick became all right as if he was released from a chain, and started to walk and showed no signs of sickness. The chief paid the Companions what was agreed. Some of the Companions suggested that the earnings to be divided among themselves, but eventually they agreed to consult Prophet Muhammad (PBUH). Upon reaching Madinah, they narrated the story to the Prophet Muhammad (PBUH) whereupon he remarked, "How did he come to know that Surah Al-Fatihah can be used as a cure (Ruqya)? You have done the right thing. Distribute your reward amongst yourselves and assign a share for me as well".<sup>9</sup>*

The Holy *Qur'an* mentions the regulations and laws related to the human health in aspects involving the body and mind.<sup>10</sup> The Holy *Qur'an* mentions in *Surah Al-Hijr* (15) verse 82, "*And we sent down in the Qur'an such things that have healing and mercy for the believers*".<sup>11</sup> In *Surah Yunus* (10) verse 57, "*Mankind there has come to you a guidance from your Lord and a healing for (the diseases) in your hearts, and for those who believe a guidance and a mercy*" and in *Surah Fussilat* (41) verse 44, "*And declare (O Muhammad) that (the Qur'an) is guidance and healing for the believers*".<sup>11</sup> Thus, *Qur'anic* verses can be used as the guideline in order to improve health or treating diseases. A Muslim has faith and strong belief in the words of Allah<sup>12</sup> as evident in the Holy *Qur'an*. Allah tells mankind in the Holy *Qur'an* in *Surah Al-An'aam* (6) verse 17, "*And if Allah touches you with harm, none can remove it but He, and if He touches you with good, then He is Able to do all things*" and in *Surah Ash-Shu'ara* (26) verse 80, "*And when I am ill, it is (God) who cures me*".<sup>11</sup> Muslims believe that in every disease or illness, Allah has made the cure.<sup>13</sup> From these verses, it can be said that sound healing with *Qur'anic* verses can possibly be one of the media to treat human diseases.

## MATERIALS AND METHOD

### Sample Collection

With the ethical approval IUM /IACUCAPPROVAL /2015/ (5) (24) rabbit articular cartilage was obtained from a commercial source. This preliminary study used only two samples (n=2). The cartilage samples were collected in 50ml Falcon tube (Falcon, Franklin Lakes, NJ) filled with normal saline to avoid the sample from drying. In order to maintain the temperature between 4-8°C, a thermal container (Coleman, Sunbeam, Canada) was used to transport the cartilage back to the laboratory. The cartilages then were kept in refrigerator at 4°C before the start of the harvesting process. The samples were processed within the first 24 hours of sample collection to minimize the reduction of cell viability.

### Chondrocytes Isolation

A sterile scalpel was used to cut and minced the articular cartilage into smaller fragments averaging 1 mm<sup>3</sup> each. Then, the minced cartilage was washed twice with the phosphate buffered saline (PBS; pH 7.2) (Gibco, Grand Island, NY) containing 100 U/ml penicillin, 100 µg/ml streptomycin (Gibco) and 0.25 µg/ml amphotericin B (Gibco). The samples were then submitted to enzymatic digestion in order to isolate the cells. They were digested using 0.6% Collagenase A (Gibco, Grand Island, NY) in an orbital incubator (Stuart Scientific, Redhill, UK) at 37°C for 4 hours. After digestion, the chondrocytes suspension was centrifuged (Jouan, Duguay Trouin, SH) at 6000 rpm, at 37°C for 5 minutes. The digested tissues were then allowed to settle and the supernatant containing the cells was removed. The resultant cells pellet was washed again with PBS to

remove any remaining enzyme. After final centrifugation, the cells pellet was re-suspended in PBS for total cell count and viability test using haemocytometer (Weber Scientific International, Ltd., Middlx, England) and Trypan Blue Dye (Gibco, Invitrogen, USA) exclusion test.

### Chondrocytes Culture

The isolated chondrocytes cells were then cultured in 6 well plate (Thermo Scientific, Nunclon Delta Surface, Denmark) at a concentration of 5000 cells/cm<sup>2</sup> using Dulbecco's Modified Eagle Medium / Nutrient Mixture F-12 (Ham) (1:1) DMEM/F12 (Gibco, Life Technologies, USA) supplemented with 10% foetal bovine serum (FBS) (Gibco, Life Technologies, USA), 1% of antibiotics-antimycotic (Gibco, Life Technologies, USA), 4-(2-hydroxyethyl)-1-piperazineethanesulphonic acid (HEPES) buffer (Gibco, Life Technologies, USA), L-glutamine (Gibco, Life Technologies, USA) and L-ascorbic acid (Sigma®, Sigma-Aldrich Corporation, USA). Initial seeding per well was 50000 cells. The cells were maintained at 37°C in a humidified incubator (Thermo Scientific, BarnsteadLab Line, USA) at 5% CO<sub>2</sub> atmosphere. In order to grow the chondrocytes, culture media was replaced every 2 days to maintain the phenotypic stability to the cells.

### Chondrocytes Treatment

The chondrocytes cultures were divided into three groups; (1) not exposed to any sound and serves as control group, (2) exposed to recitation of *Surah Al-Fatihah* and (3) exposed to recitation of an Arabic poem. These groups were monitored at different passages (P0, P1, P2, and P3). The exposure duration of the recitations to the exposed groups, groups 2 and 3, were standardized to 14 minutes. To introduce the recitations to the cells, a speaker was used. The speaker was wrapped with a plastic wrapper and sprayed with 70% ethanol before use to maintain the sterility. The speaker was then placed at a fixed distance under the 6-well plate of the respective groups. The same speaker was used to introduce the different recitations to the cells.

The exposure of the sound was done inside an incubator. To control the sound level exposure, the intensity of the sound (for different volume settings) emitted from the speaker located at a fixed distance from the cell culture was measured using sound level meter (SLM). The volume that produced the intensity in the range of normal human conversation level (-60 dB SPL) was selected to be the constant volume setting throughout the study. This measurement was done for both *Qur'anic* and Arabic poem recitation.

Two incubators were used for this experiment. One incubator was used to keep all the cell cultures in a standard environment. The other incubator was used to expose groups 2 and 3 individually to the assigned recitation. The *Qur'anic* recitation featured a recitation of *Surah Al-Fatihah* by Sheikh

Abdul Basit Abdus Samad, a renowned *Qur'anic* reciter. A well-known Arabic poem recitation was downloaded from the internet.

### Cell Count and Viability

Trypan Blue Exclusion method is a rapid and simple measurement. The dead cells will appear blue as it retains the dyes inside the cells. Viable cells will not absorb the dyes and will have intact cell membranes. As the cells reached the 80% - 90% confluence, total cell count and viability for each passage were recorded. Growth kinetics analysis was evaluated using parameters that include (1) percentage of cell viability, (2) total cell count, (3) cell growth rate, (4) total number of cell doubling (TNCD) and (5) population doubling time (PDT).

Cell viability can be defined as the number of viable cell in the total cell which includes dead and viable cells. Meanwhile, total cell count is the cumulative cells yield at the end of this experiment. Growth rate was calculated based on the amount of cells/total area/days. Total number of cells doubling can be expressed by  $\log(\text{cell number} - \text{initial seeding}) / \log_2$ . Rates of cell division were calculated assuming that chondrocytes develop from single cell. The average population doubling times are given by days/TNCD. The values obtained can provide a minimum estimated for the maximum rate of cell division. All the calculations for the growth kinetic analysis are given in the Table 1 below. The normal trend for the cell growth of the chondrocytes sample would be an increase in cell growth with subsequent passages. The results were recorded and expressed as average as the results represent two samples (n=2).

Table 1. Formula for each parameter.

### RESULTS

Parameter	Formula
Cell viability	$= \frac{\text{Total viable cells}}{\text{Total cells (viable + dead)}} \times 100$
Total Cells Count	$= \frac{\text{Total viable cells}}{9} \times 10^4 \times \text{cell suspension}$
Growth Rate	$= \frac{(\text{Total cells count} - \text{Initial seeding})}{\text{Area (9.6cm}^2\text{)} \times \text{Days}}$
TNCD	$= \frac{\text{Log}(\text{Total cells count}) - \text{Log}(\text{Initial seeding})}{\text{Log } 2}$
PDT	$= \frac{\text{Days}}{\text{TNCD}}$

Five different aspects of cell reaction was observed and taken as the dependent variables of this study. They are the percentage of cell viability, total cell count, growth rate, total number of cell doubling and population doubling time.

### Cell Viability (%)

It was observed that the viability of chondrocytes varied with different groups. However, the viability of every group in different passages was always higher than 84%, which indicated that all cells in each group were healthy cells. The results for each group can be appreciated as follows: (1) *Qur'anic* group: P0= 94%, P1= 95%, P2= 94%, P3= 92%, (2) Arabic poem: P0= 89%, P1= 92%, P2= 93%, P3= 92%, and (3) Control group: P0= 88%, P1= 91%, P2=89%, P3=84%. The results can be summarized as in Figure 1.




Figure 1: Cell viability (%) of rabbit chondrocytes monolayer in every passage

### Total Cell Count

The chondrocytes cells exposed to the *Qur'anic* recitation show the highest value as compared to the Arabic poem recitation and control group. The result showed a similar trend between all groups. The chondrocytes cells proliferation showed a linear increment. The result for each total cell count in each group were given as the average number. It can be appreciated as follows: (1) *Qur'anic* group = 1075000; (2) Arabic poem = 972222; and (3) Control group = 850000. The results are summarized as in Figure 2.




Figure 2: Total cell count of rabbit chondrocytes monolayer in every group

### Growth Rate

The results for the growth rate of primary chondrocytes showed that the *Qur'anic* and Arabic poem recitation both are higher as compared to the control group. It was observed that the growth rate for the control and Arabic poem recitation groups were increased until P2 and started to decrease at P3. It showed the normal trend as the graph demonstrates a sigmoidal curve. The results for each group can be appreciated as follows: (1) *Qur'anic* group: P0= 3703, P1= 5425, P2= 8295, P3= 8681; (2) Arabic poem: P0= 3241, P1= 4991, P2= 7619, P3= 7137; and (3) Control group: P0= 2836, P1= 4557, P2= 6655, P3= 6076. The results are summarized as in Figure 3.




Figure 3: Growth rate of rabbit chondrocytes monolayer in every passage

### Total Number of Cell Doubling (TNCD)

The TNCD indicates the magnitude of chondrocytes expansion in culture. It was observed that the TNCD for all groups increased in different passages. The results revealed that the both *Qur'anic* and Arabic poem recitation groups showed similar pattern as compared to the control group. The cells exposed to the *Qur'anic* recitation showed the highest value as compared to the Arabic poem recitation and control group in every passage. The results for each group can be appreciated as follows: (1) *Qur'anic* group= 9.670643555; (2) Arabic poem= 9.093755026; and (3) Control group= 8.503862965. The results are summarized as in Figure 4.




Figure 4: Total number of cell doubling (TNCD) of chondrocytes in every passage

## Population doubling time (PDT)

The PDT value revealed that the both *Qur'anic* and Arabic poem recitation groups showed similar pattern as compared to the control group. Slight differences were observed between the groups when compared between the different passages. It was observed that the PDT for all groups decreased with passages. The *Qur'anic* recitation group required the shortest time to double the population as compared to Arabic poem and control groups within the same passage. The results for each group can be appreciated as follows: (1) *Qur'anic* group: P0= 2.29, P1= 1.69, P2= 1.19, P3= 1.16; (2) Arabic poem: P0= 2.45, P1= 1.76, P2= 1.24, P3= 1.28; and (3) Control group: P0= 2.64, P1= 1.85, P2= 1.32, P3= 1.38. The results are summarized as in Figure 5.




Figure 5: Population doubling time (PDT) of chondrocytes monolayer in every passage

## DISCUSSION

The use of sound has been utilized for treatment purposes even though the effect of the sound itself is still not fully understood. This experiment was conducted to analyse the growth kinetic profile on the healthy rabbit articular chondrocytes after being exposed to the *Qur'anic* and Arabic poem recitation. This experiment setting uses serially cultured chondrocytes in order to relate the ageing of the chondrocytes with the loss of cells phenotype. The chondrocytes cells were quantified using hemocytometer and trypan blue dye.

The study examined the effect of *Qur'anic* recitation on chondrocytes viability after several passages. The percentage of cell viability was noticeably higher than that seen in control and Arabic poem groups. Cell viability is taken as acceptable or satisfactory if it reached more than 80% for that indicates the cells were healthy. The findings for the cell viability in the control group show a normal trend as the viability is more than 84% in every passages. The trend can also be observed in the other 2 groups. However, the group exposed to the *Qur'anic* recitation shows a better result as compared to the group exposed to the

Arabic poem recitation and control group. This means that the amount of dead cells in the *Qur'anic* group is low. This increased the percentage of cell viability. The results highly suggest that *Qur'anic* recitation can have therapeutic effect as it was able to increase the amount of viable cells and reduce the number of dead cells.

Total cells count is the total number of viable and dead cells in a given area. It can also be described as the total amount of cell proliferation. The normal trend should indicate an increment along the passages. The findings suggest that the proliferation of rabbit articular chondrocytes for the control group experienced a reduction at P2, but an increase was noted at P3. The group exposed to the Arabic poem recitation also shows similar trend. The *Qur'anic* recitation group shows the normal trend as the total cell count increased along the passages. This might suggest that with the *Qur'anic* recitation facilitates cells to exhibit normal growth.

The growth rate measurement is the rate of change in the number of cells in a culture per unit time. An increasing trend in cell proliferation is observed within the groups. This can be explained with the increment of the total count of the chondrocytes cell in the group exposed to the *Qur'anic* recitation. This might suggest that the lifespan of the cell can be expanded further until it shows reduction in the value.

Total number of cell doubling is the amount of the cells that doubled in size or number. The control group shows a normal trend which demonstrates an increase in total number of cell doubling from P0 to P3. The cells exposed to *Qur'anic* recitation show a higher total number of cell doubling compared to the group exposed with the Arabic poem recitation and control group. This result supports the result obtained from the total cell count calculation. An increase in the total number of cell doubling between the passages indicates that the cells experienced an increase in proliferation. Thus it is suggestive that proliferation is better with *Qur'anic* recitation. A possible application of this result could be in the therapeutic effect on wound healing process. It is anticipated that the increased cell proliferation could speed up the recovery at the wounded site.

Population doubling time is the time required for cells to double in size or number. The control group demonstrates a normal trend of the population doubling time. A shorter population doubling time was recorded in the group exposed to the *Qur'anic* recitation as compared to other two groups. This result reveals the positive effect of the *Qur'anic* recitation where the cells require shorter time to proliferate after being exposed to the recitation. Shorter time in population doubling time could contribute positively in healing mechanism.

Based on the overall results, it can be inferred that sound, in this case *Qur'anic* and Arabic poem recitations, can indeed alter the proliferation of the

rabbit chondrocytes cells. The same results were showed by Lestard, et al. (2013) when the audible sound was able to alter the cell proliferation.<sup>14</sup> However, the cells exposed to the *Qur'anic* recitation shows a better result in the cell proliferation of rabbit chondrocytes as compared to the Arabic poem recitation and control group. It also shows that the *Qur'anic* recitation have more impact than the Arabic poem recitation. The results highlight the higher potential in *Qur'anic* recitation on healing effect as compared to the Arabic poem recitation. However, it could not be ruled out that the possibility of different effects observed between *Qur'anic* and Arabic poem exposed groups due to different reciters. Different voices contain different frequency and wavelength profiles.<sup>5</sup> Further work is indicated to verify these postulations.

## CONCLUSION

This study was carried out to identify the potential of *Qur'anic* recitation on the proliferation of chondrocytes derived from rabbit articular cartilage. This study demonstrates that the rabbit chondrocytes cells exposed to the *Qur'anic* recitation shows a positive effect as proliferation of the cells. The proliferation was found to be increased when compared to the Arabic poem recitation and control groups. The chondrocytes culture results optimized in this preliminary work could pave the way for possible applications in treating injuries to articular cartilage and also be a potential approach for cartilage regeneration in tissue engineering.

## Future Work

Based on the preliminary work above, it is felt that there exists justification to examine the effect of *Qur'anic* recitation as the medium for sound healing on cell cultures in a larger scale. At the moment due to limited number of samples, any significant relationship cannot be generated. Further experiments involving more samples are needed to be conducted in order to further validate these finding. Based on the findings, studies relating to wound healing assay can be conducted to observe the proliferation from photomicrograph aspect.

Under normal circumstances, the original chondrocytic characteristics of the chondrocytes will change to fibroblastic traits. These changes on the cells morphology can be clearly seen through the microscopic observation. Gene expression analysis using chondrogenic (cartilaginous) markers shall be proposed to see whether or not *Qur'anic* recitation has an influence on cells (original) phenotype in monolayer culture. Furthermore, this study used the chondrocytes obtained from healthy rabbits. Future work should also be directed to replicate this study on diseased cells. A new variable, a Western poem recitation shall be introduced. The results of this preliminary study is only based on two samples (n=2). With only two values, there is a lot of uncertainty. No statistical

analysis was conducted and no significant relationship can be observed at this juncture. A bigger sample size will be used for the future work to determine the statistical difference. The work is underway.

## ACKNOWLEDGEMENT

The authors thanked the Kulliyah of Allied Health Sciences (KAHS), International Islamic University Malaysia (IIUM), Kuantan Campus. The authors also thanked Ministry of Education, MyRA Incentive Research Grant Scheme (MIRGS13-01-002-0003) and the Ministry of Science, Technology and Innovation (MOSTI) Malaysia (06-01-08-SF0238/SF14-012-0062).

## REFERENCES

1. Ruiz EL, Peran M, Molinos JC, et al. Chondrocytes extract from patients with osteoarthritis induces chondrogenesis in infrapatellar fat pad-derived stem cells. *Osteoarthritis and Cartilage* 2013; 21: 246-258.
2. Lozito TP, Alexander PG, Lin H, et al. Three-dimensional osteochondral micro tissue to model pathogenesis of osteoarthritis. *Stem Cell Research & Therapy* 2013; 4: 1-6.
3. Heidari M, Tahmasebi M, Etemad S, et al. In-vitro human chondrocytes culture; a modified protocol. *Middle-East Journal of Scientific Research* 2011; 9: 102-109.
4. Tew SR, Murdoch AD, Rauchenberg RP, et al. Cellular methods in cartilage research: primary human chondrocytes in culture and chondrogenesis in human bone marrow stem cells. *Elsevier, Methods* 2008; 45: 2-9.
5. Fauzan N, Shahidan SN, Amran NH, et al. The therapeutic effects of listening to Quranic verse and rhythmic Zikr. *University of Malaysia Sarawak* 2014.
6. Kamal NF, Mahmood NH, Zakaria NA. Modeling brain activities during reading working memory task: Comparison between reciting Quran and reading book. *Procedia Social and Behavioral Sciences* 2013; 97: 83-89.
7. Erkkila J, Punkanen M, Fachner J, et al. Individual music therapy for depression: randomized controlled trial. *The British Journal of Psychiatry* 2011; 199: 132-139.
8. Arshad NW, Sukri SM, Muhammad LN, et al. Makhraj recognition for Al-Quran recitation using MFCC. *International Journal of Intelligent Processing* 2013; 4(2): 45-53.
9. Khan MM. The translation of the meaning of Sahih Al-Bukhari Arabic-English Volume 7. Riyadh, Saudi Arabia: Darussalam publishers and distributors 1997.
10. Akhlaqi E. Scientific remarks of the Quran to health and medical teachings. *Journal of Applied Science and Agriculture* 2014; 9: 960-964.
11. Ali AY. The Holy Qur'an original Arabic text with English translation and selected commentaries. Saba Islamic Media, Kuala Lumpur 2013.

12. Tumiran MA, Mohamad SP, Saat RM, et al. Addressing sleep disorder of autistic children with Qur'anic sound therapy. *Health* 2013; 5: 73-79.
13. Ateeq M, Jehan S, Mehmmod R. Faith healing; Modern health care. *Professional Medical Journal* 2014; 21: 295-301.
14. Lestard NDR, Valente RC, Lopes AG. Direct effects of music in non-auditory cells in culture. *Noise & Health* 2013; 15: 307-314.