

[< Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)[Full Text](#) [View at Publisher](#)Studia Universitatis Babes-Bolyai Chemia [Open Access](#)  
Volume 63, Issue 3, 2018, Pages 95-116

## Gum arabic: An optimization of ultrasonic-assisted extraction of antioxidant activity (Article) [\(Open Access\)](#)

Elmour, A.A.M.<sup>a,b</sup>, Mirghani, M.E.S.<sup>a,c</sup> [✉](#), Kabbashi, N.A.<sup>a</sup>, Alam, M.Z.<sup>a</sup>, Musa, K.H.<sup>d</sup> [👤](#)<sup>a</sup>Bioenvironmental Engineering Research Centre (BERC), Biotechnology Engineering, Faculty of Engineering, International Islamic University, Malaysia (IIUM), P. O. Box 10, Gombak, Kuala Lumpur, 50728, Malaysia<sup>b</sup>Department of Biochemistry & Gum Processing, Gum Arabic Research Centre, University of Kordofan, Box: 160, Elobied, Sudan<sup>c</sup>International Institute for Halal Research and Training (INHART), IIUM, P. O. Box 10, Gombak, Kuala Lumpur, 50728, Malaysia[View additional affiliations](#) [v](#)

### Abstract

[v](#) [View references \(49\)](#)

Gum Arabic (GA), also known as Acacia seyal gum (ASG), is a dried exudate from trees of Acacia senegal and Acacia seyal. It provides a rich source of non-viscous soluble fiber with significant health benefits and high antioxidant properties. Tonnes of raw GA are exported annually at a high cost with limited utilization in extraction form. Techniques for the extraction of the bioactive components of GA are available but the high extraction time and the capacity and quality of extraction hinders these procedures. Ultrasonic-assisted extraction is one of the most effective techniques for the recovery of antioxidant and phenolic compounds from ASG. A comparatively low extraction time has been reported for ultrasonication, but the influence of several extraction conditions such as temperature, time and ultrasonic power on the yield of extraction has not been thoroughly studied. This study investigates the optimal ultrasonic extraction conditions for maximum recovery of antioxidant and phenolic compounds from ASG using Response Surface Methodology (RSM) under the Central Composite Design (CCD). Three ultrasonic parameters, namely time in the range of (1-3 hours), power in the range of (1-3 level or 12 to 40 kHz) and temperature from (25-60 °C) were tested for their impact on antioxidant activity. The capacity of the extracts was determined by the scavenging activity of 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical, ferric reducing antioxidant power (FRAP) assay, and total phenolic compounds (TPC). The results indicated that ultrasonic time, power and temperature had a positive impact on antioxidant capacity and phenolic compounds. The optimum ultrasonic conditions were found to be a time of 3 hours, a power of 40 kHz, and a temperature of 42.50°C, under which, forty-eight bioactive compounds from the ASG extract were separated by Gas Chromatography coupled to Tandem Mass Spectrometry (GC-MS/MS). © 2018, Universitatea Babes-Bolyai, Catedra de Filosofie Sistemtica. All rights reserved.

### SciVal Topic Prominence [ⓘ](#)

Topic: [Spray drying](#) | [Drug Compounding](#) | [Wall materials](#)Prominence percentile: 99.358 [ⓘ](#)

### Author keywords

[Acacia seyal gum](#) [Bioactive compounds](#) [Gum Arabic](#) [Response Surface Methodology](#) [Ultrasonication](#)

### Funding details

Funding sponsor	Funding number	Acronym
-----------------	----------------	---------

### Metrics [ⓘ](#)

0 Citations in Scopus

0 Field-Weighted Citation Impact



### PlumX Metrics [v](#)

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

### Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

### Related documents

Optimum conditions for microwave assisted extraction for recovery of phenolic compounds and antioxidant capacity from macadamia (macadamia tetraphylla) skin waste using water

Dailey, A. , Vuong, Q.V. (2016) *Processes*

The effect of sodium bisulfite immersion to the potential of Candi banana peel ethanol extract as radical scavenger and UV protection

Laeliocattleya, R.A. , Wijaya, J. (2019) *IOP Conference Series: Earth and Environmental Science*

Biological activities of Umbilicaria crustulosa (Ach.) Frey acetone extract

Zlatanović, I. , Stanković, M. , Jovanović, V.S. (2017) *Journal of the Serbian Chemical Society*[View all related documents based on references](#)

---

International Islamic University Malaysia

Authors > Keywords >

---

Faculty of Engineering, Alexandria University

---

#### Funding text

The authors would like to acknowledge express their gratitude to the Department of Biotechnology Engineering (BTE), Faculty of Engineering at the International Islamic University Malaysia (IIUM) and the International Institute for Halal Research and Training (INHART) at IIUM. The 1st author would like to express his gratitude to Dr. Elbasheir Sallam for his continuous and unlimited financial support in conducting this research.

---

ISSN: 12247154

DOI: 10.24193/SUBBCHEM.2018.3.08

Source Type: Journal

Document Type: Article

Original language: English

Publisher: Universitatea Babes-Bolyai, Catedra de Filosofie Sistematica

---

#### References (49)

[View in search results format >](#)

All | [Export](#) [Print](#) [E-mail](#) [Save to PDF](#) [Create bibliography](#)

- 
- 1 Anderson, D.  
(1993) *Forest Ecology and Management*, 58.
- 
- 2 Elmqvist, B., Olsson, L., Elamin, E.M., Warren, A.  
(2005) *Agroforestry Systems*, p. 64.
- 
- 3 Islam, A., Phillips, G., Sljivo, A., Snowden, M., Williams, P.  
(1997) *Food Hydrocolloids*, 11.
- 
- 4 Yasseen, G.A., Salih, A.A., Ahmed, M.E.  
(2014) *Procedia-Social and Behavioral Sciences*, p. 120.
- 
- 5 Al Assaf, S., Phillips, G.O., Williams, P.A.  
(2005) *Food Hydrocolloids*, p. 19.
- 
- 6 Ali, B.H., Beegam, S., Al-Lawati, I., Waly, M., Al Za'abi, M., Nemmar, A.  
(2013) *Physiological Research*, p. 62.
-