

Free Full Text from Publisher

 Look Up Full Text

Full Text Options



Save to EndNote online

Add to Marked List

◀ 1 of 1 ▶

Antioxidant Activities of *Dialium indum* L. Fruit and Gas Chromatography-Mass Spectrometry (GC-MS) of the Active Fractions

By: **Osman, MF** (Osman, Muhamad Faris)^[1]; **Hassan, NM** (Hassan, Norazian Mohd)^[1]; **Khatib, A** (Khatib, Alfi)^[1]; **Tolos, SM** (Tolos, Siti Marponga)^[2]

[View ResearcherID and ORCID](#)

ANTIOXIDANTS

Volume: 7 Issue: 11

Article Number: 154

DOI: 10.3390/antiox7110154

Published: NOV 2018

Document Type: Article

Abstract

The fruit of *Dialium indum* L. (Fabaceae) is one of the edible wild fruits native to Southeast Asia. The mesocarp is consumed as sweets while the exocarp and seed are regarded as waste. This study aimed to evaluate the antioxidant activities of the fruit by using four assays, which measure its capabilities in reducing phosphomolybdic-phosphotungstic acid reagents, neocuproine, 2,2-diphenyl-picrylhydrazyl (DPPH), and inhibiting linoleic acid peroxidation. The active fractions were then analyzed by gas chromatography-mass spectrometry (GC-MS). The results showed that the seed methanol fraction (SMF) exhibited the strongest antioxidant activity with significantly higher ($p < 0.05$) gallic acid equivalence (GAE), total antioxidant capacity (TAC), and DPPH radical scavenging activity (IC₅₀ 31.71; 0.88 $\mu\text{g/mL}$) than the other fractions. The exocarp dichloromethane fraction (EDF) was the discriminating fraction by having remarkable linoleic acid peroxidation inhibition (IC₅₀ 121.43; 2.97 $\mu\text{g/mL}$). A total of thirty-eight metabolites were detected in derivatized EDF and SMF with distinctive classes of phenolics and amino acids, respectively. Bioautography-guided fractionation of EDF afforded five antioxidant-enriched subfractions with four other detected phenolics. The results revealed the antioxidant properties of *D. indum* fruit, which has potential benefits in pharmaceutical, nutraceutical, and cosmeceutical applications.

Keywords

Author Keywords: *Dialium indum*; exocarp; seed; antioxidant; phenolic acids; amino acids; GC-MS analysis

KeyWords Plus: PHENOLIC-COMPOUNDS; AMINO-ACIDS; CAPACITY; PLANTS; PARTS; QUANTIFICATION; IDENTIFICATION; EXTRACTION; FLAVONOIDS; ALCOHOL

Author Information

Reprint Address: Hassan, NM (reprint author)

+ Int Islamic Univ Malaysia, Kulliyah Pharm, Dept Pharmaceut Chem, Kuantan 25200, Pahang, Malaysia.

Addresses:

+ [1] Int Islamic Univ Malaysia, Kulliyah Pharm, Dept Pharmaceut Chem, Kuantan 25200, Pahang, Malaysia

+ [2] Int Islamic Univ Malaysia, Kulliyah Sci, Dept Computat & Theoret Sci, Kuantan 25200, Pahang, Malaysia

E-mail Addresses: farisosman@iium.edu.my; norazianmh@iium.edu.my; alfikhatib@iium.edu.my; smtolos@iium.edu.my

Funding

Funding Agency	Grant Number
International Islamic University Malaysia (IIUM)	RIGS16-123-0287

[View funding text](#)

Publisher

MDPI, ST ALBAN-ANLAGE 66, CH-4052 BASEL, SWITZERLAND

Categories / Classification

Research Areas: Biochemistry & Molecular Biology; Pharmacology & Pharmacy; Food Science & Technology

Web of Science Categories: Biochemistry & Molecular Biology; Chemistry, Medicinal; Food Science & Technology

Citation Network

In Web of Science Core Collection

0

Times Cited

 Create Citation Alert

50

Cited References

[View Related Records](#)

Use in Web of Science

Web of Science Usage Count

0

Last 180 Days

0

Since 2013

[Learn more](#)

This record is from:

Web of Science Core Collection
- Science Citation Index Expanded

Suggest a correction

If you would like to improve the quality of the data in this record, please suggest a correction.

[See more data fields](#)

◀ 1 of 1 ▶

Cited References: 50Showing 30 of 50 [View All in Cited References page](#)*(from Web of Science Core Collection)*

1. **Antioxidant activity and phenolic profile of various morphological parts of underutilised *Baccaurea angulata* fruit** Times Cited: 12
By: Ahmed, Idris Adewale; Mikail, Maryam Abimbola; Bin Ibrahim, Muhammad; et al.
FOOD CHEMISTRY Volume: 172 Pages: 778-787 Published: APR 1 2015
2. **Evaluation of antioxidant and antiulcerogenic activities of *Opuntia ficus indica* f. *inermis* flowers extract in rats** Times Cited: 38
By: Alimi, Hichem; Hfaiedh, Najla; Bouoni, Zouhour; et al.
ENVIRONMENTAL TOXICOLOGY AND PHARMACOLOGY Volume: 32 Issue: 3 Pages: 406-416 Published: NOV 2011
3. **Cupric Ion Reducing Antioxidant Capacity Assay for Food Antioxidants: Vitamins, Polyphenolics, and Flavonoids in Food Extracts** Times Cited: 23
By: Apak, Resat; Gueclue, Kubilay; Oezyuerek, Mustafa; et al.
Advanced Protocols in Oxidative Stress I Book Series: Methods in Molecular Biology Volume: 477 Pages: 163-193 Published: 2008
4. **The cupric ion reducing antioxidant capacity and polyphenolic content of some herbal teas** Times Cited: 175
By: Apak, Resat; Guclu, Kubilay; Ozyurek, Mustafa; et al.
INTERNATIONAL JOURNAL OF FOOD SCIENCES AND NUTRITION Volume: 57 Issue: 5-6 Pages: 292-304 Published: AUG-SEP 2006
5. **Evaluation of antioxidant potentials of different solvent-fractions of *Dialium indum* (African black velvet tamarind) fruit pulp *in vitro*** Times Cited: 1
By: Bamikole, A.O.; Ibidun, O.O.; Ibitayo, O.A.; et al.
Potravinarstvo Slovak J. Food Sci. Volume: 12 Pages: 70-78 Published: 2018
[\[Show additional data\]](#)
6. **Impact of Hydrophobicity on Antioxidant Efficacy in Low-Moisture Food** Times Cited: 4
By: Barden, Leann; Barouh, Nathalie; Villeneuve, Pierre; et al.
JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY Volume: 63 Issue: 24 Pages: 5821-5827 Published: JUN 24 2015
7. **High Correlation of 2,2-diphenyl-1-picrylhydrazyl (DPPH) Radical Scavenging, Ferric Reducing Activity Potential and Total Phenolics Content Indicates Redundancy in Use of All Three Assays to Screen for Antioxidant Activity of Extracts of Plants from the Malaysian Rainforest** Times Cited: 54
By: Clarke, Garry; Ting, Kang Nee; Wiart, Christophe; et al.
ANTIOXIDANTS Volume: 2 Issue: 1 Pages: 1-10 Published: MAR 2013
8. **Serviceberry, a berry fruit with growing interest of industry: Physicochemical and quali-quantitative health-related compound characterisation** Times Cited: 5
By: Donno, D.; Cerutti, A. K.; Mellano, M. G.; et al.
JOURNAL OF FUNCTIONAL FOODS Volume: 26 Pages: 157-166 Published: OCT 2016
9. **Determination of some phenolic compounds in *Crocus sativus* L. corms and its antioxidant activities study** Times Cited: 19
By: Esmaili, N.; Ebrahimzadeh, H.; Abdi, K.; et al.
PHARMACOGNOSY MAGAZINE Volume: 7 Issue: 25 Pages: 74-80 Published: JAN-MAR 2011
10. **Simultaneous identification and quantification of the sugar, sugar alcohol, and carboxylic acid contents of sour cherry, apple, and ber fruits, as their trimethylsilyl derivatives, by gas chromatography-mass spectrometry** Times Cited: 31
By: Fuzfai, Z.; Katona, ZF; Kovacs, E; et al.
JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY Volume: 52 Issue: 25 Pages: 7444-7452 Published: DEC 15 2004
11. **Non-Nutritive Compounds in Fabaceae Family Seeds and the Improvement of Their Nutritional Quality by Traditional Processing - a Review** Times Cited: 16
By: Gulewicz, Piotr; Martinez-Villaluenga, Cristina; Kasproicz-Potocka, Malgorzata; et al.
POLISH JOURNAL OF FOOD AND NUTRITION SCIENCES Volume: 64 Issue: 2 Pages: 75-89 Published: JUN 2014