

Document details

< Back to results | 1 of 1

Export Download Print E-mail Save to PDF Add to List More... >

[Full Text](#) View at Publisher

Journal of Family Violence
Volume 34, Issue 2, 15 February 2019, Pages 81-92

Prevalence and Factors Associated with Intimate Partner Violence among Postpartum Women Attending Government Primary Health Care Clinics in Malaysia (Article) [\(Open Access\)](#)

Chan, Y.Y.^a, Rosman, A.^b, Ahmad, N.A.^a, Mohamad Kasim, N.^a, Abd Razak, M.A.^a, Omar, M.^c, Abdul Aziz, F.A.^a, Awaluddin, S.M.^a, Yusof, M.^a, Jamaluddin, R.^a, Ab Wahab, N.A.^a, Rosli, N.H.^d

^aCentre for Family Health Research, Institute for Public Health, National Institutes of Health, Ministry of Health Malaysia, Kuala Lumpur, Malaysia

^bViolence and Injury Prevention (VIP) Unit, Disease Control Division, Ministry of Health Malaysia, Putrajaya, Malaysia

^cIjok Health Clinic, Kuala Selangor, Selangor, Malaysia

[View additional affiliations](#) v

Abstract

[v View references \(47\)](#)

Intimate partner violence (IPV) is an important gender-based, social, and public health problem that affects women worldwide, including women who are pregnant or have recently given birth. Studies have shown that violence against women often increases during pregnancy and the postpartum period. This study aims to examine lifetime and past-year prevalence of IPV among postpartum women in Malaysia, and to determine the socio-demographic as well as husband's/ partner's behavioral factors associated with IPV exposure. This is a nationwide, cross-sectional and clinic-based study involving a total of 5727 women at 6 to 16 weeks postpartum, who attended randomly selected government health clinics between July to November 2016. Face-to-face interviews were conducted by trained female enumerators based on a pre-validated structured questionnaire, using mobile devices as data collection tools. Chi-squared tests and multivariable logistic regressions were used to investigate selected factors associated with IPV exposure. The lifetime and past-year prevalence of any form of IPV among postpartum women were 4.94% (95% CI [3.81,6.39]) and 2.42% (95% CI [1.74,3.35]) respectively, with the highest prevalence being emotional violence, followed by physical and sexual violence. Multivariable analysis showed that husband's/ partner's behaviors, such as frequent alcohol use, drug use, fighting habits and controlling behaviour were significantly associated with both lifetime and past-year IPV (all $p < 0.001$ for past-year IPV). These findings suggest that prevention and intervention strategies for IPV should consider the prevention of substance use and reducing controlling behaviors by husband/ partner, as well as raising awareness to build healthy relationships through education. © 2018, The Author(s).

SciVal Topic Prominence [i](#)

Topic: Violence | Women | wife beating

Prominence percentile: 89.637 [i](#)

Author keywords

Factors Intimate partner violence Malaysia Postpartum Prevalence Women

Funding details

Funding sponsor	Funding number	Acronym
-----------------	----------------	---------

Metrics [?](#)

0 Citations in Scopus
0 Field-Weighted Citation Impact



PlumX Metrics [v](#)
Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

[Set citation feed >](#)

Related documents

Controlling Behavior and Lifetime Physical, Sexual, and Emotional Violence in sub-Saharan Africa

McClintock, H.F., Trego, M.L., Wang, E.M. (2019) *Journal of Interpersonal Violence*

Alcohol use among partner violent adults: Reviewing recent literature to inform intervention

Langenderfer, L. (2013) *Aggression and Violent Behavior*


Prevalence, risk factors, and complications of violence against pregnant women in a hospital in Peninsular Malaysia


Khaironisak, H., Zaridah, S., Hasanain, F.G. (2017) *Women and Health*

[View all related documents based on references](#)

[Find more related documents in Scopus based on:](#)

[Authors >](#) [Keywords >](#)

National Institutes of Health
See opportunities 

Institute of Population and Public Health
See opportunities 

Health Research

NMRR-15-2404-26677

Funding text #1

Centre for Family Health Research, Institute for Public Health, National Institutes of Health, Ministry of Health Malaysia, Kuala Lumpur, Malaysia

Funding text #2

We used the bilingual (Bahasa Malaysia and English) WHO Multi-country Study on Women's Health and Life Events Questionnaire (World Health Organization 2000) which has been locally validated for use in Malaysia (Saddki et al. 2013). The respondents were interviewed by trained female nurses (enumerators) face-to-face on a one-to-one basis in private without the presence of their husband/partner at the selected government health clinics. Prior to conducting the survey, the enumerators were trained to conduct interviews specifically related to physical and sexual abuse. The questionnaire was installed on mobile devices and all answers from the respondents were digitally recorded. The survey procedure was approved by the Medical Research and Ethics Committee (MREC), Ministry of Health Malaysia (NMRR-15-2404-26677). Informed consent was sought and obtained from all the study participants and confidentiality was assured.

ISSN: 08857482

Source Type: Journal

Original language: English




DOI: 10.1007/s10896-018-0014-0

Document Type: Article

Publisher: Springer New York LLC

References (47)

[View in search results format >](#)

All [Export](#)  Print  E-mail  Save to PDF [Create bibliography](#)

-
- 1 Abdullah, R., Raj-Hashim, R., Schmitt, G. (1995) *Battered women in Malaysia: Prevalence, problems and public attitudes*. Cited 8 times. A summary report of Women's Aid Organization Malaysia's National Research on Domestic Violence. Petaling Jaya: Women's Aid Organization (WAO)
-
- 2 Abeya, S.G., Afework, M.F., Yalew, A.W. **Intimate partner violence against women in western Ethiopia: Prevalence, patterns, and associated factors** ([Open Access](#)) (2011) *BMC Public Health*, 11, art. no. 913. Cited 34 times. doi: 10.1186/1471-2458-11-913 [View at Publisher](#)
-
- 3 Al Serkal, F., Hussein, H., El Sawaf, E., Al Faisal, W., Hasan Mahdy, N., Wasfy, A. **Intimte partner violence against women in Dubai: Prevalence, associated factors and health consequences, 2012-2013** (2014) *Middle East Journal of Psychiatry and Alzheimers*, 5 (3), pp. 19-27. Cited 2 times.
-