

## Document details

[Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More...](#)[View at Publisher](#)Industrial and Engineering Chemistry Research  
Volume 58, Issue 2, 16 January 2019, Pages 762-770

## Prediction of Mefenamic Acid Solubility and Molecular Interaction Energies in Different Classes of Organic Solvents and Water (Article)

Abdul Mudalip, S.K.<sup>a</sup>  Abu Bakar, M.R.<sup>b</sup>  Jamal, P.<sup>c</sup> Adam, F.<sup>a</sup> <sup>a</sup>Centre of Excellence for Advanced Research in Fluid Flow, Faculty of Chemical and Natural Resources Engineering, University Malaysia Pahang, Kuantan, Pahang, Gambang, 26300, Malaysia<sup>b</sup>Department of Pharmaceutical Technology, Kulliyah of Pharmacy, International Islamic University Malaysia, Kuantan, Pahang, Bandar Indera Mahkota, 25200, Malaysia<sup>c</sup>Department of Biotechnology Engineering, Kulliyah of Engineering, International Islamic University Malaysia, Kuala Lumpur, 50728, Malaysia

## Abstract

[View references \(46\)](#)

Determination of solubility data either through experimental- or model-based approaches becomes a necessity in the crystallization of pharmaceutical compounds. The current work predicts the mefenamic acid solubility and molecular interaction energy, namely electrostatic (H-MF), hydrogen bonding (H-HB), and van der Waals (H-vdW), in different solvents at temperatures from 298 to 323 K using the conductor-like screening model for real solvents (COSMO-RS). The solvents used were N,N-dimethylacetamide, N,N-dimethylformamide, acetone, ethyl acetate, ethanol, iso-propyl alcohol, n-hexane, n-heptane, cyclohexane, and water. The Gibbs free energy of fusion required in COSMO-RS computation was determined using differential scanning calorimetry and reference solubility method. The accuracy of methods employed in prediction of solubility were evaluated using mean squared quadratic error (MSE). The mefenamic acid solubility predicted using COSMO-RS with reference solubility method showed a small MSE value of less than 2%. The predicted solubility also follows the same trend as the experimental values and increases with temperature. The predicted H-HB energy and Gibbs free energy changes of mefenamic acid dissolution in the solvents studied highly influence the solubility data. Therefore, COSMO-RS with reference solubility method is a promising approach to predict the solubility and intermolecular interaction energy of mefenamic acid in different solvents. © Copyright 2018 American Chemical Society.

SciVal Topic Prominence 

Topic: Solubility | Ethanol | fraction solubility

Prominence percentile: 99.006



## Reaxys Database Information

[View Compounds](#)

## Indexed keywords

Engineering controlled terms:

[Acetone](#) [Differential scanning calorimetry](#) [Dimethylformamide](#) [Forecasting](#) [Free energy](#)  
[Gibbs free energy](#) [Heptane](#) [Hexane](#) [Hydrogen bonds](#) [Molecular interactions](#)  
[Molecular structure](#) [Organic solvents](#) [Van der Waals forces](#)

Metrics  [View all metrics](#)

1 Citation in Scopus

6.19 Field-Weighted Citation Impact



## PlumX Metrics

Usage, Captures, Mentions,  
Social Media and Citations  
beyond Scopus.

## Cited by 1 document

2018 International Conference of Chemical Engineering and Industrial Biotechnology (ICCEIB) Preface

Cheng, C.K. , Rahman Khan, M.M. , Rasid, R.A.  
*(2019) Industrial and Engineering Chemistry Research*[View details of this citation](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

## Related documents

Optimal solvent screening for the crystallization of pharmaceutical compounds from multisolvent systems

Sheikholeslamzadeh, E. , Chen, C.-C. , Rohani, S.  
*(2012) Industrial and Engineering Chemistry Research*

Vapour-liquid and vapour-liquid-liquid equilibrium modeling for binary, ternary, and quaternary systems of solvents

Sheikholeslamzadeh, E. , Rohani,

Reaxys PhD Prize 2019  
The global award for ambitious young chemists is now open!

[Apply now](#)

Phase Equilibria

Prediction of pharmaceutical and chemical



Engineering  
uncontrolled terms

Conductor-like screening model for real solvents | Gibbs free energy changes  
Intermolecular interaction energies | Model based approach | Molecular interaction energy  
N,N-dimethylacetamide | N,N-Dimethylformamide | Pharmaceutical compounds

Engineering main  
heading:

Solubility

compounds in pure and mixed  
solvents using predictive models  
Sheikholeslamzadeh, E., Rohani,  
S.  
(2012) *Industrial and Engineering  
Chemistry Research*

View all related documents based  
on references

Find more related documents in  
Scopus based on:

Authors > Keywords >

## Funding details

Funding sponsor	Funding number	Acronym
Ministry of Higher Education, Malaysia	RDU1200607	MOHE
International Islamic University Malaysia		IIUM
Universiti Malaysia Pahang	RDU150359,RDU1803107	UMP
Universiti Malaysia Pahang		UMP

### Funding text #1

\*E-mail: kholijah@ump.edu.my. Tel.: (+609)-549 2356. Fax: (+609)-549 2889. ORCID Siti K. Abdul Mudalip: 0000-0002-9759-2756 Author Contributions The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript. Funding The work was financially supported by the Ministry of Education Malaysia (Exploratory Research Grant Scheme RDU1200607), International Islamic University Malaysia (Endowment Fund B12-363-0841) and the Universiti Malaysia Pahang (RDU150359 and RDU1803107). Notes The authors declare no competing financial interest.

### Funding text #2

S.K.A.M. is grateful to the Ministry of Education Malaysia and the Universiti Malaysia Pahang. The authors are grateful to F. Eckert and A. Klamt, COSMOtherm, version C3.0, release 13.01; COSMOlogic GmbH & Co. KG, Leverkusen, Germany, 2013, for their technical support.

ISSN: 08885885

DOI: 10.1021/acs.iecr.8b02722

CODEN: IECRE

Document Type: Article

Source Type: Journal

Publisher: American Chemical Society

Original language: English

## References (46)

[View in search results format >](#)

All    [Export](#)    [Print](#)    [E-mail](#)    [Save to PDF](#)    [Create bibliography](#)


- 1 Tang, W., Xie, C., Wang, Z., Wu, S., Feng, Y., Wang, X., Wang, J., (...), Gong, J.  
**Solubility of androstenedione in lower alcohols**

(2014) *Fluid Phase Equilibria*, 363, pp. 86-96. Cited 24 times.  
doi: 10.1016/j.fluid.2013.11.008

[View at Publisher](#)

- 2 Bennet, R.C., Myerson, A.S.

Crystallizer Selection  
(2007) *Handbook*  
Butterworth-Heinemann



Reaxys PhD Prize 2019  
The global award for ambitious  
young chemists is now open!

[Apply now](#)

X