

Document details

< Back to results | 1 of 1

Export Download Print E-mail Save to PDF Add to List More... >

[Full Text](#) View at Publisher

Plant physiology
Volume 179, Issue 2, 1 February 2019, Pages 544-557

Characterization of CRISPR Mutants Targeting Genes Modulating Pectin Degradation in Ripening Tomato (Article)

Wang, D.^a, Samsulrizal, N.H.^b, Yan, C.^c, Allcock, N.S.^d, Craigon, J.^a, Blanco-Ulate, B.^e, Ortega-Salazar, I.^e, Marcus, S.E.^f, Bagheri, H.M.^a, Perez Fons, L.^g, Fraser, P.D.^g, Foster, T.^a, Fray, R.^a, Knox, J.P.^f, Seymour, G.B.^a ✉

^aSchool of Biosciences, University of Nottingham, Sutton Bonington, Loughborough, LE12 5RD, United Kingdom
^bDepartment of Plant Science, Kulliyah of Science, International Islamic University Malaysia, Kuantan, Pahang 25200, Malaysia
^cInstitution of Vegetable Research, Shanxi Academy of Agricultural Sciences, 030031, China

[View additional affiliations](#) ▾

Abstract

Tomato (*Solanum lycopersicum*) is a globally important crop with an economic value in the tens of billions of dollars, and a significant supplier of essential vitamins, minerals, and phytochemicals in the human diet. Shelf life is a key quality trait related to alterations in cuticle properties and remodeling of the fruit cell walls. Studies with transgenic tomato plants undertaken over the last 20 years have indicated that a range of pectin-degrading enzymes are involved in cell wall remodeling. These studies usually involved silencing of only a single gene and it has proved difficult to compare the effects of silencing these genes across the different experimental systems. Here we report the generation of CRISPR-based mutants in the ripening-related genes encoding the pectin-degrading enzymes pectate lyase (PL), polygalacturonase 2a (PG2a), and β-galactanase (TBG4). Comparison of the physiochemical properties of the fruits from a range of PL, PG2a, and TBG4 CRISPR lines demonstrated that only mutations in PL resulted in firmer fruits, although mutations in PG2a and TBG4 influenced fruit color and weight. Pectin localization, distribution, and solubility in the pericarp cells of the CRISPR mutant fruits were investigated using the monoclonal antibody probes LM19 to deesterified homogalacturonan, INRA-RU1 to rhamnogalacturonan I, LM5 to β-1,4-galactan, and LM6 to arabinan epitopes, respectively. The data indicate that PL, PG2a, and TBG4 act on separate cell wall domains and the importance of cellulose microfibril-associated pectin is reflected in its increased occurrence in the different mutant lines. © 2019 American Society of Plant Biologists. All Rights Reserved.

ISSN: 15322548
Source Type: Journal
Original language: English

DOI: 10.1104/pp.18.01187
PubMed ID: 30459263
Document Type: Article
Publisher: NLM (Medline)

© This record is sourced from MEDLINE®/PubMed®, a database of the U.S. National Library of Medicine

< Back to results | 1 of 1

^ Top of page

Metrics

0 Citations in Scopus
0 Field-Weighted Citation Impact



PlumX Metrics ▾
Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

[Set citation feed >](#)

Related documents

Find more related documents in Scopus based on:

[Authors >](#)

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API

Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

Customer Service

- Help
- Contact us

