

Document details

[< Back to results](#) | 1 of 1[↗ Export](#) [↓ Download](#) [🖨 Print](#) [✉ E-mail](#) [📄 Save to PDF](#) [☆ Add to List](#) [More... >](#)[View at Publisher](#)Marine Pollution Bulletin
Volume 141, April 2019, Pages 472-481

Comparison of prediction model using spatial discriminant analysis for marine water quality index in mangrove estuarine zones (Article)

Samsudin, M.S.^{a,c}, Azid, A.^a [✉](#), Khalit, S.I.^a, Sani, M.S.A.^b, Lananan, F.^a [👤](#)^aFaculty Bioresources and Food Industry, Universiti Sultan Zainal Abidin (UniSZA), Besut Campus, Besut, Terengganu 22200, Malaysia^bInternational Institute for Halal Research and Training, International Islamic University Malaysia, Selangor, Malaysia^cDr. F.A.S. Technologies, Block D1, 2nd Floor UniSZA Digital Hub, UniSZA Besut Campus, Besut, Terengganu 222000, Malaysia

Abstract

[View references \(64\)](#)

The prediction models of MWQI in mangrove and estuarine zones were constructed. The 2011–2015 data employed in this study entailed 13 parameters from six monitoring stations in West Malaysia. Spatial discriminant analysis (SDA) had recommended seven significant parameters to develop the MWQI which were DO, TSS, O&G, PO₄, Cd, Cr and Zn. These selected parameters were then used to develop prediction models for the MWQI using artificial neural network (ANN) and multiple linear regressions (MLR). The SDA-ANN model had higher R² value for training (0.9044) and validation (0.7113) results than SDA-MLR model and was chosen as the best model in mangrove estuarine zone. The SDA-ANN model had also demonstrated lower RMSE (5.224) than the SDA-MLR (12.7755). In summary, this work suggested that ANN was an effective tool to compute the MWQI in mangrove estuarine zone and a powerful alternative prediction model as compared to the other modelling methods. © 2019 Elsevier Ltd

Author keywords

Artificial neural networks Discriminant analysis Mangrove estuarine zone Marine water quality

Multiple linear regression

ISSN: 0025326X
 CODEN: MPNBA
 Source Type: Journal
 Original language: English

DOI: 10.1016/j.marpolbul.2019.02.045
 Document Type: Article
 Publisher: Elsevier Ltd

References (64)

[View in search results format >](#)

All [Export](#) [🖨 Print](#) [✉ E-mail](#) [Save to PDF](#) [Create bibliography](#)

- 1 Abdullah, A.R., Tahir, N.M., Loong, T.S., Hoque, T.M., Sulaiman, A.H.
 The GEF/UNDP/IMO Malacca Straits demonstration project: Sources of pollution
 (1999) *Marine Pollution Bulletin*, 39 (1-12), pp. 229-233. Cited 46 times.
 doi: 10.1016/S0025-326X(98)00206-9
[View at Publisher](#)

Metrics [?](#)

0 Citations in Scopus
 0 Field-Weighted
 Citation Impact

PlumX Metrics [▼](#)Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 0 documents

Inform me when this document
is cited in Scopus:[Set citation alert >](#)[Set citation feed >](#)

Related documents

Spatial air quality modelling using chemometrics techniques: A case study in Peninsular Malaysia | Pemodelan ruang kualiti udara menggunakan teknik-teknik kemometrik: Satu kajian kes di semenanjung Malaysia

Azid, A. , Juahir, H. , Amran, M.A.
 (2015) *Malaysian Journal of Analytical Sciences*

Applied chemometric approach in identification sources of air quality pattern in Selangor, Malaysia

Hua, A.K.
 (2018) *Sains Malaysiana*

Identification source of variation on regional impact of air quality pattern using chemometric

Azid, A. , Juahir, H. , Ezani, E.
 (2015) *Aerosol and Air Quality Research*

View all related documents based on references

- 2 Aertsen, W., Kint, V., van Orshoven, J., Özkan, K., Muys, B.
Comparison and ranking of different modelling techniques for prediction of site index in Mediterranean mountain forests

(2010) *Ecological Modelling*, 221 (8), pp. 1119-1130. Cited 144 times.
doi: 10.1016/j.ecolmodel.2010.01.007

[View at Publisher](#)

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

- 3 Ali, M.M., Ali, M.L., Islam, M.S., Rahman, M.Z.
Preliminary assessment of heavy metals in water and sediment of Karnaphuli River, Bangladesh ([Open Access](#))

(2016) *Environmental Nanotechnology, Monitoring and Management*, 5, pp. 27-35. Cited 81 times.
<http://www.journals.elsevier.com/environmental-nanotechnology-monitoring-and-management/>
doi: 10.1016/j.enmm.2016.01.002

[View at Publisher](#)

- 4 Alizadeh, M.J., Kavianpour, M.R.
Development of wavelet-ANN models to predict water quality parameters in Hilo Bay, Pacific Ocean

(2015) *Marine Pollution Bulletin*, 98 (1-2), pp. 171-178. Cited 33 times.
www.elsevier.com/locate/marpolbul
doi: 10.1016/j.marpolbul.2015.06.052

[View at Publisher](#)

- 5 Amiri, B.J., Nakane, K.
Comparative prediction of stream water total nitrogen from land cover using artificial neural network and multiple linear regression approaches

(2009) *Polish Journal of Environmental Studies*, 18 (2), pp. 151-160. Cited 15 times.

- 6 An, Q., Wu, Y., Wang, J., Li, Z.
Assessment of dissolved heavy metal in the Yangtze River estuary and its adjacent sea, China

(2010) *Environmental Monitoring and Assessment*, 164 (1-4), pp. 173-187. Cited 36 times.
doi: 10.1007/s10661-009-0883-z

[View at Publisher](#)

- 7 Azid, A., Juahir, H., Toriman, M.E., Kamarudin, M.K.A., Saudi, A.S.M., Hasnam, C.N.C., Aziz, N.A.A., (...), Yamin, M.
Prediction of the level of air pollution using principal component analysis and artificial neural network techniques: A case study in Malaysia

(2014) *Water, Air, and Soil Pollution*, 225 (8), art. no. 2063. Cited 40 times.
<http://www.kluweronline.com/issn/0049-6979/>
doi: 10.1007/s11270-014-2063-1

[View at Publisher](#)