

Document details

[< Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)Journal of Pharmaceutical Sciences and Research
Volume 10, Issue 1, January 2018, Pages 167-174Potential oral wound healing of topical application of dental gel prepared from *Baccaurea angulata* fruit in diabetic rats (Article)Rahman, M.N.A.^a, Sukmasari, S.^b, Doolaanea, A.A.^{a,c}, Qader, O.A.J.A.^d ^aAdvanced Drug Delivery Lab, Department of Pharmaceutical Technology, Kulliyah of Pharmacy International Islamic University Malaysia, Bandar Indera Mahkota, Kuantan, Pahang 25200, Malaysia^bDepartment of Paediatric Dentistry, Kulliyah of Dentistry, International Islamic University Malaysia, Bandar Indera Mahkota, Kuantan, Pahang 25200, Malaysia^cIKOP Sdn Bhd, Kulliyah of Pharmacy, International Islamic University, Malaysia[View additional affiliations](#) 

Abstract

[View references \(52\)](#)

Objective: This study investigates the effect of *Baccaurea angulata* fruit extract towards wound healing activity at tooth extractions sites. Significance: High antioxidant content in underutilised *Baccaurea angulata* were expected to open up new possibilities for tooth extraction procedure among diabetic patients. Methods: Sprague dawley rats were divided into 6 groups, and their left mandibular first molar tooth were extracted. The alveolar socket were treated accordingly where group 1, 2, 3, 4, and 5 were diabetic rats (induced using Streptozotocin) applied with plain gel, alveolex, 3%, 5%, and 10% *Baccaurea angulata* extract respectively. Group 6 were non diabetic rats applied with plain gel. On day 4, 7 and 18, rats were euthanized, and the tissue samples were processed for histomorphological analysis. Results: It was found that *Baccaurea angulata* does significantly promotes tooth extraction wound healing in diabetic rats comparable to diabetic group treated with alveolex and non-diabetic rats at early phase of the wound. However the positive effect were not obvious at day 7, while at day 18, groups treated with *Baccaurea angulata* treated group had higher mean scores in most categories although may not statistically significant different. BADG concentration of 4% was found to give the best effects on wound healing. Conclusion: To a certain extent, *Baccaurea angulata* extract does promotes tooth extraction wound healing in diabetic rats. © 2018, Pharmainfo Publications. All rights reserved.

SciVal Topic Prominence 

Topic: Propolis | Bees | green propolis

Prominence percentile: 98.177 

Reaxys Database Information

[View Compounds](#)

Author keywords

[Baccaurea angulata](#) [Dental gel](#) [Diabetes](#) [Extracts](#) [Tooth extraction](#) [Wound healing](#)

Indexed keywords

EMTREE drug terms: [acetylcysteine](#) [dental gel](#) [herbaceous agent](#) [unclassified drug](#)
[wound healing promoting agent](#)Metrics 

0 Citations in Scopus

0 Field-Weighted
Citation ImpactPlumX Metrics Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 0 documents

Inform me when this document
is cited in Scopus:[Set citation alert >](#)[Set citation feed >](#)

Related documents

Headspace analysis and
characterisation of South African
propolis volatile compounds
using GCxGC–ToF–MSKamatou, G. , Sandasi, M. ,
Tankeu, S.
(2019) *Brazilian Journal of
Pharmacognosy*Assessment of bone width for
implants in the posterior
mandibleBlock, M.S. , Scoggin, Z.D. , Yu,
Q.
(2015) *Journal of Oral and
Maxillofacial Surgery*Effects of coenzyme Q10
encapsulated in nanoliposomes
on wound healing processes after
tooth extractionTodorovic, K. , Jovanovic, G. ,
Todorovic, A.
(2018) *Journal of Dental Sciences*[View all related documents based
on references](#)[Find more related documents in
Scopus based on:](#)

EMTREE medical terms:

- animal experiment
- animal model
- animal tissue
- Article
- Baccaurea angulata
- controlled study
- Euphorbiaceae
- fibroblast
- fruit
- histopathology
- neovascularization (pathology)
- nonhuman
- rat
- tooth extraction
- wound healing

Chemicals and CAS Registry Numbers:

acetylcysteine, 616-91-1

Funding details

| Funding sponsor | Funding number | Acronym |
|---|------------------|---------|
| International Islamic University Malaysia | | IIUM |
| | RIGS 15-047-0047 | |

Funding text

This research was funded by the Research and Innovation Grant Scheme: RIGS 15-047-0047 of International Islamic University Malaysia.

ISSN: 09751459

Source Type: Journal

Original language: English

Document Type: Article

Publisher: Pharmainfo Publications

References (52)

[View in search results format >](#)

- All
 Export
 Print
 E-mail
 Save to PDF
 Create bibliography

- 1 Dunnill, C., Patton, T., Brennan, J., Barrett, J., Dryden, M., Cooke, J., Leaper, D., (...), Georgopoulos, N.T.
Reactive oxygen species (ROS) and wound healing: the functional role of ROS and emerging ROS-modulating technologies for augmentation of the healing process

(2017) *International Wound Journal*, 14 (1), pp. 89-96. Cited 64 times.
[http://onlinelibrary.wiley.com.ezproxy.um.edu.my/journal/10.1111/\(ISSN\)1742-481X](http://onlinelibrary.wiley.com.ezproxy.um.edu.my/journal/10.1111/(ISSN)1742-481X)
 doi: 10.1111/iwj.12557

[View at Publisher](#)

- 2 Baltzis, D., Eleftheriadou, I., Veves, A.
Pathogenesis and Treatment of Impaired Wound Healing in Diabetes Mellitus: New Insights

(2014) *Advances in Therapy*, 31 (8), pp. 817-836. Cited 88 times.
<http://www.springer.com.ezproxy.um.edu.my/springer+healthcare/journal/12325>
 doi: 10.1007/s12325-014-0140-x

[View at Publisher](#)

- 3 Boulton, A.J.M.
The pathway to foot ulceration in diabetes

(2013) *Medical Clinics of North America*, 97 (5), pp. 775-790. Cited 46 times.
 doi: 10.1016/j.mcna.2013.03.007

[View at Publisher](#)