

## Document details

[< Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)[View at Publisher](#)Malaysian Orthopaedic Journal [Open Access](#)

Volume 13, Issue 1, March 2019, Pages 25-29

Outcome of islanded gastrocnemius musculocutaneous flap in orthopaedic practice (Article) [\(Open Access\)](#)Yusof, N.M.<sup>a</sup> [✉](#), Ahmad-Alwi, A.A.<sup>b</sup> [👤](#)<sup>a</sup>Department of Orthopaedics, Traumatology and Rehabilitation, International Islamic University of Malaysia, Kuantan, Malaysia<sup>b</sup>Department of Surgery, International Islamic University of Malaysia, Kuantan, Malaysia

## Abstract

[View references \(16\)](#)

Introduction: Large wounds in the leg require combination of local flaps or free flap for wound coverage. Gastrocnemius musculocutaneous flap (GMCF) allows a large wound to be covered by a single local flap. However, the conventional GMCF is often associated with donor site morbidity where the exposed soleus raphe causes poor uptake of the skin graft. Islanding the skin on the muscles allows the donor site to be closed primarily, thus avoiding the donor site morbidity. Materials and Methods: Medical records of twelve patients who underwent islanded GMCF surgery from 2004 till 2018 were reviewed retrospectively. Results: The mean age was 31 years old. Eight cases were with open fracture of the tibia, two degloving injury exposing the patella, one open fracture of patella and necrotising soft tissue infection. The wound size ranged from 12cm<sup>2</sup> to 120cm<sup>2</sup>. All flaps survived. Three patients required skin grafting at the donor site while in the rest the donor sites were able to be closed primarily. Four patients developed deep infection, one healed after vacuum dressing, one after bone transport and one after split thickness skin graft. One patient ended up with below knee amputation after developing chronic osteomyelitis of the tibia. Conclusion: Islanded gastrocnemius musculocutaneous flap is an effective simple alternative for coverage of large soft tissue defects from the knee to half of the leg distally with minimal donor site morbidity. Aggressive debridement of unhealthy tissue is necessary to prevent infection following wound coverage with this flap. © 2019, Malaysian Orthopaedic Association. All rights reserved.

## Author keywords

[Leg injuries](#) [Surgical flaps](#) [Wounds](#)

## Funding details

Funding sponsor	Funding number	Acronym
International Islamic University Malaysia	RIGS 16-110-0274	

## Funding text

This study was funded by International Islamic University of Malaysia Research Initiative Grant (RIGS 16-110-0274).

Metrics [?](#)

0 Citations in Scopus

0 Field-Weighted Citation Impact

PlumX Metrics [v](#)

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

## Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

## Related documents

Baropodometric analysis of the functional donor-site morbidity after gastrocnemius or soleus muscle-flap procedure | Analyse baropodométrie des séquelles fonctionnelles liées au prélèvement des lambeaux gastrocnémiens et soléaire

Lasserre, G. , Cornu, J.-Y. , Vidal, C.

*(2011) Annales de Chirurgie Plastique et Esthétique*

Soft-tissue coverage: Gastrocnemius and soleus rotational muscle flaps

Sherman, R. , Li, W.-Y. *(2012) Master Techniques in Orthopaedic Surgery: Fractures: Third Edition*

## Gastrocnemius flap

Masquelet, A.-C. , Sassu, P. *(2009) Flaps and Reconstructive Surgery*[View all related documents based on references](#)

ISSN: 19852533

Source Type: Journal

Original language: English

DOI: 10.5704/MOJ.1903.004

Document Type: Article

Publisher: Malaysian Orthopaedic Association

All  Export  Print  E-mail  Save to PDF  Create bibliography

Authors > Keywords >

- 
- 1 Pu, L.L.Q.  
Soft-tissue coverage of an extensive mid-tibial wound with the combined medial gastrocnemius and medial hemisoleus muscle flaps: The role of local muscle flaps revisited  
  
(2010) *Journal of Plastic, Reconstructive and Aesthetic Surgery*, 63 (8), pp. e605-e610. Cited 10 times.  
doi: 10.1016/j.bjps.2010.05.003  
  
[View at Publisher](#)
- 
- 2 Ger, R.  
Muscle transposition for treatment and prevention of chronic post-traumatic osteomyelitis of the tibia  
  
(1977) *Journal of Bone and Joint Surgery - Series A*, 59 (6), pp. 784-791. Cited 113 times.  
doi: 10.2106/00004623-197759060-00012  
  
[View at Publisher](#)
- 
- 3 McCraw, J.B., Fishman, J.H., Sharzer, L.A.  
The versatile gastrocnemius myocutaneous flap  
  
(1978) *Plastic and Reconstructive Surgery*, 62 (1), pp. 15-23. Cited 116 times.  
doi: 10.1097/00006534-197807000-00002  
  
[View at Publisher](#)
- 
- 4 Mathes, S.J., Nahai, F.  
Classification of the vascular anatomy of muscles: Experimental and clinical correlation  
  
(1981) *Plastic and Reconstructive Surgery*, 67 (2), pp. 177-187. Cited 433 times.  
doi: 10.1097/00006534-198167020-00007  
  
[View at Publisher](#)
- 
- 5 Kim, H.H., Jeong, J.H., Seul, J.H., Cho, B.C.  
New design and identification of the medial sural perforator flap: An anatomical study and its clinical applications  
  
(2006) *Plastic and Reconstructive Surgery*, 117 (5), pp. 1609-1618. Cited 50 times.  
doi: 10.1097/01.prs.0000207077.19601.86  
  
[View at Publisher](#)
- 
- 6 Hallock, G.G.  
Anatomic basis of the gastrocnemius perforator-based flap  
  
(2001) *Annals of Plastic Surgery*, 47 (5), pp. 517-522. Cited 82 times.  
<http://journals.lww.com/annalsplasticsurgery/pages/default.aspx>  
doi: 10.1097/00000637-200111000-00008  
  
[View at Publisher](#)
-