

[< Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)[Full Text](#) [View at Publisher](#)Pharmaceutical biology [Open Access](#)
Volume 57, Issue 1, 1 December 2019, Pages 335-344

Methanol extract of *Muntingia calabura* leaves attenuates CCl₄-induced liver injury: possible synergistic action of flavonoids and volatile bioactive compounds on endogenous defence system (Article)

[\(Open Access\)](#)

Zakaria, Z.A.^{a,b,c}, Mahmood, N.D.^c, Omar, M.H.^d, Taher, M.^e, Basir, R.^f^aLaboratory of Halal Science Research, Halal Products Research Institute, Universiti Putra Malaysia, Serdang, Malaysia^bIntegrative Pharmacogenomics Institute (iPROMISE), Universiti Teknologi MARA, Malaysia^cDepartment of Biomedical Science, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, Serdang, Malaysia[View additional affiliations](#)

Abstract

CONTEXT: *Muntingia calabura* L. (Muntingiaceae) exerts antioxidant and anti-inflammatory activities, thus, it might be a good hepatoprotective agent. OBJECTIVE: This study investigates the effect of methanol extract of *M. calabura* leaves (MMCL) on hepatic antioxidant and anti-inflammatory activities in CCl₄-induced hepatotoxic rat. MATERIALS AND METHODS: Sprague Dawley rats (n = 6) were treated (p.o.) with 10% DMSO (Groups 1 and 2), 50 mg/kg N-acetylcysteine (Group 3) or, 50, 250, or 500 mg/kg MMCL (Groups 4-6) for 7 consecutive days followed by pretreatment (i.p.) with vehicle (Group 1) or 50% CCl₄ in olive oil (v/v) (Groups 2-6) on day 7th. Plasma liver enzymes and hepatic antioxidant enzymes and pro-inflammatory cytokines concentrations were measured while liver histopathology was examined. RESULTS: MMCL, at 500 mg/kg, significantly (p < 0.05) ameliorated CCl₄-induced hepatotoxicity by decreasing the plasma level of alanine transaminase (429.1 versus 168.7 U/L) and aspartate transaminase (513.8 versus 438.1 U/L) as well as the tissue level of nitric oxide (62.7 versus 24.1 nmol/g tissue). At 50, 250, or 500 mg/kg, MMCL significantly (p < 0.05) reduced the tumour necrosis factor α (87.8 versus 32.7 pg/mg tissue), interleukin-1 β (1474.4 versus 618.3 pg/mg tissue), and interleukin-6 (136.7 versus 30.8 pg/mg tissue) while increased the liver catalase (92.1 versus 114.4 U/g tissue) and superoxide dismutase (3.4 versus 5.5 U/g tissue). Additionally, qualitative phytochemicals analysis showed that MMCL contained gallic acid, ferulic acid, quercetin, and genistein. DISCUSSION AND CONCLUSIONS: MMCL ability to attenuate CCl₄-induced hepatotoxicity could be helpful in the development of hepatoprotective agents with fewer side effects.

SciVal Topic Prominence

Topic: Antioxidants | Methanol | Malabathricum leaves

Prominence percentile: 58.289

Author keywords

[antioxidant activity](#) [GCMS](#) [hepatoprotective activity](#) [Muntingiaceae](#) [oxidative stress markers](#) [phytoconstituents](#)
[pro-inflammatory mediators](#) [UHPLC-ESI](#)

Metrics

0 Citations in Scopus
0 Field-Weighted Citation Impact

PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

Related documents

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

© This record is sourced from MEDLINE/PubMed, a database of the U.S. National Library of Medicine

< Back to results | 1 of 1

^ Top of page

About Scopus

What is Scopus
Content coverage
Scopus blog
Scopus API
Privacy matters

Language

日本語に切り替える
切换到简体中文
切换到繁體中文
Русский язык

Customer Service

Help
Contact us

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

 RELX