

Document details

[< Back to results](#) | 1 of 1[↗ Export](#) [↓ Download](#) [🖨 Print](#) [✉ E-mail](#) [📄 Save to PDF](#) [★ Add to List](#) [More... >](#)[View at Publisher](#)Journal of Environmental Management
Volume 232, 15 February 2019, Pages 1012-1020

Citizen science frontiers horseshoe crab population regain at their spawning beach in East Peninsular Malaysia (Article)

Mat Zauki, N.A.^a, Satyanarayana, B.^a ✉, Fairuz-Fozi, N.^a, Nelson, B.R.^b ✉, Martin, M.B.^c, Akbar-John, B.^d, Chowdhury, A.J.K.^e ^aMangrove Research Unit, Institute of Oceanography and Environment (INOS), Universiti Malaysia Terengganu, Kuala Nerus, Terengganu 21030, Malaysia^bInstitute of Tropical Biodiversity and Sustainable Development (ITBSD), Universiti Malaysia Terengganu, Kuala Nerus, Terengganu 21030, Malaysia^cSchool of Marine and Environmental Sciences (PPSMS), Universiti Malaysia Terengganu, Kuala Nerus, Terengganu 21030, Malaysia[View additional affiliations](#) 

Abstract

[View references \(93\)](#)

Carcinoscorpius rotundicauda and *Tachypleus gigas* may co-exist and share common spawning grounds elsewhere but at Balok (East Coast of Peninsular Malaysia), *C. rotundicauda* is an understudied species. Neglected as research candidate because of inaccessible spawning grounds, smaller size and less commercial value than *T. gigas* and also, difficulty to attain from the wild has made *C. rotundicauda* population status remaining unidentified at Balok. This standpoint drove the present attempt because anthropic activities like structure placement and mining are point-source for runoffs that load sediments into Balok River. While erosion-accretion events have altered Balok River width, the shore sediments in Balok Beach were transitioned between medium-fine and fine sand between years 2012 and 2016. Eventually by year 2016, the *C. rotundicauda* were depositing 5117 eggs in 91 nests from 200 to 1000 m range along this corridor facing South China Sea. From this yield, *C. rotundicauda* released 2880 eggs in 56 nests during the Southwest monsoon, 1254 eggs in 19 nests during the Northeast monsoon and 983 eggs in 16 nests during the Inter-monsoon seasons. Though female *C. rotundicauda* opted to lay their eggs in shallow burrows at lower shorelines, the absence of erosion and substantial silt and clay (>20%) deposition facilitates *C. rotundicauda* embryogenesis with brief periods of temperature and salinity shocks during day-time falling tides. This encourages *C. rotundicauda* to emerge with increasing abundance and carry out bi-monthly spawning at Balok Beach. In short, shore restoration initiatives like systematic boat docking, proper disposal of nets and waste and, periodic fish-catching operations were effectively led by the Balok fisher citizen scientist. This successful community joint-cooperation proves that citizen-led caretaking of degraded beaches offers marine life protection and are practical for coastal area management especially at areas where other oviparous animals such as turtles and crocodiles are harboured. © 2018 Elsevier Ltd

SciVal Topic Prominence Topic: [Limulus polyphemus](#) | [crab](#) | [Limulus](#)Prominence percentile: 65.908 

Author keywords

[Citizen science](#) [Fisheries management](#) [Horseshoe crab](#) [Seasonal ecology](#) [Sedimentation](#) [Shore recovery](#)

Indexed keywords

Metrics  [View all metrics >](#)

2 Citations in Scopus

5.11 Field-Weighted
Citation ImpactPlumX Metrics Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 2 documents

Compendium of hand, foot and mouth disease data in Malaysia from years 2010–2017

Nelson, B.R., Edinur, H.A., Abdullah, M.T. (2019) *Data in Brief*Effects of shore sedimentation to *Tachypleus gigas* (Müller, 1785) spawning activity from Malaysian watersNelson, B.R., Zhong, J.M.H., Zauki, N.A.M. (2019) *Journal of Sustainability Science and Management*[View all 2 citing documents](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

Related documents

Effects of shore sedimentation to *Tachypleus gigas* (Müller, 1785) spawning activity from Malaysian watersNelson, B.R., Zhong, J.M.H., Zauki, N.A.M. (2019) *Journal of Sustainability Science and Management**Carcinoscorpius rotundicauda* (Latreille, 1802) population status and spawning behaviour at

GEOBASE Subject Index:

- accretion
- catch statistics
- coastal erosion
- coastal sediment
- coastal zone management
- coexistence
- commercial species
- cooperative sector
- crab
- crocodilian
- fishing community
- human activity
- salinity
- spawning ground
- temperature effect
- turtle
- wild population

EMTREE medical terms:

- adult
- article
- Carcinoscorpius
- citizen scientist
- crocodile
- ecology
- embryo development
- female
- fishery management
- human
- joint
- Malaysia
- mining
- nonhuman
- oviparity
- river
- salinity
- seashore
- season
- sediment
- sedimentation
- South China Sea
- spawning
- Tachypleus
- turtle

Regional Index:

- Malaysia
- Pacific Ocean
- South China Sea
- West Malaysia

Species Index:

- Animalia
- Carcinoscorpius rotundicauda
- Crocodylidae (all crocodiles)
- Merostomata
- Tachypleus gigas
- Testudines

Pendas coast, Peninsular Malaysia

Fairuz-Fozi, N. , Satyanarayana, B. , Mat Zauki, N.A. (2018) *Global Ecology and Conservation*

Compendium of hand, foot and mouth disease data in Malaysia from years 2010–2017

Nelson, B.R. , Edinur, H.A. , Abdullah, M.T. (2019) *Data in Brief*

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

Funding details

Funding sponsor	Funding number	Acronym
Ministry of Higher Education, Malaysia	FRGS-59380	

Funding text

Authors would like to thank the Malaysian Ministry of Higher Education for funding this project under the FRGS grant (FRGS-59380, 2015). Courtesies extended by the authorities at Institute of Oceanography and Environment (INOS), School of Marine and Environmental Sciences (PPSMS) and Universiti Malaysia Terengganu (UMT) Central Lab are appreciable. Special thanks goes to Mr Raja Haris Raja Salim, the fisher head of Balok, in Pahang. Also, cooperation and assistance provided by the local fishermen communities is always remembered.

ISSN: 03014797
CODEN: JEVMA
Source Type: Journal
Original language: English

DOI: 10.1016/j.jenvman.2018.12.002
Document Type: Article
Publisher: Academic Press

References (93)

View in search results format >

- All
- Export
- Print
- E-mail
- Save to PDF
- Create bibliography

View all 93 references

1 Adibah, A.B., Ng, W.L., Tan, S.G.
The Malay Peninsula as a barrier to gene flow in an Asian horseshoe crab species, *Carcinoscorpius rotundicauda* Latreille

(2015) *Biochemical Systematics and Ecology*, 60, pp. 204-210. Cited 6 times.
www.elsevier.com/inca/publications/store/3/6/4
doi: 10.1016/j.bse.2015.04.026

View at Publisher