Document details

International Medical Journal Malaysia Open Access
Volume 18, Issue 2, 2019, Pages 101-106

Periodic assessment of antenatal and post natal serum Endothelin-1 (ET-1) and nitric oxide (NO) levels in Hypertensive Disorders of Pregnancy (HDP) (Article)

Hidayatul, R.I.[a], Tariq, A.R.[a], Jamani, N.A.[b], Maizura, M.Z.[a]

[a]Department of Basic Medical Sciences, Kulliyyah of Medicine, International Islamic University Malaysia, Malaysia
[b]Department of Family Medicine, Kulliyyah of Medicine, International Islamic University Malaysia, Malaysia

Abstract

View references (21)

Introduction: Hypertensive Disorders of Pregnancy (HDP) is an independent risk factor of cardiovascular (CVS) disease with endothelial dysfunction postulated to be the pathophysiology. Endothelin-1 (ET-1), a potent vasoconstrictor, has been identified as a pivotal mediator in HDP. Disturbances in nitric oxide (NO) bioavailability found in endothelial dysfunction may increase susceptibility to cardiovascular diseases such as hypertension. The study aims to determine serial ET-1 and NO levels in patients with HDP and its role in persistent endothelial dysfunction. Materials and Methods: Thirty-six pregnant women from the following categories (i) normal pregnant women (Control) (ii) chronic hypertension during pregnancy (CH) and (iii) pregnancy induced hypertension (PIH) participated in this study. Blood pressure indices measurements and sample collection were done at antepartum (32 weeks) and postpartum (8 weeks and 12 weeks). ET-1 and serum NO were measured using the Human ET-1 (Endothelin-1) ELISA Kit and Nitric Oxide (total) detection kit respectively. Results: Serum ET-1 was significantly higher in patients with CH (55.3 pg/ml) and PIH (35.6 pg/ml) compared to Control (11.8 pg/ml) during antenatal until 3 months postpartum (CH 38.3 pg/ml, PIH 29.5 pg/ml, Control 1.9 pg/ml). This was accompanied by significantly lower levels of serum NO in HDP patients. Conclusion: Persistently higher than normal levels of ET-1 and lower than normal levels of NO up to 3 months postpartum in patients with history of HDP indicate presence of persistent endothelial dysfunction despite BP normalisation in PIH patients. Long term NO/ET-1 imbalance may account for the increased CVS disease risk. © 2019 Default.

SciVal Topic Prominence

Topic: Pre-Eclampsia | Pregnancy | Chronic hypertension

Prominence percentile: 96.220

Chemistry database information

Substances

ET-1

Related documents

Protective effect of Theobroma cacao on nitric oxide and endothelin-1 level in endothelial cells induced by plasma from preeclamptic patients: In silico and in vitro studies
Barokah, L. , Baktiyani, S.C.W. , Kalsum, U.
(2016) European Journal of Integrative Medicine

Emerging drugs for preeclampsia - The endothelium as a target
Sasser, J.M. , Murphy, S.R. , Granger, J.P.
(2015) Expert Opinion on Emerging Drugs

Endothelin type B (ETB) receptors: Friend or foe in the pathogenesis of pre-eclampsia and future cardiovascular disease (CVD) risk?
Colafella, K.M.M.
(2018) Clinical Science

Author keywords

Endothelin-1 (ET-1) | Hypertensive Disorders of Pregnancy (HDP) | Nitric Oxide (NO) | Persistent Endothelial Dysfunction

ISSN: 18234631
Source Type: Journal
Original language: English

Document Type: Article
Publisher: International Islamic University Malaysia

References (21)

View in search results format >

1 Romundstad, P.R., Magnussen, E.B., Smith, G.D., Vatten, L.J.

Hypertension in pregnancy and later cardiovascular risk: Common antecedents? (Open Access)

(2010) *Circulation*, 122 (6), pp. 579-584. Cited 193 times.
doi: 10.1161/CIRCULATIONAHA.110.943407

View at Publisher

2 (2014) *CLINICAL PRACTICE GUIDELINES Management of Hypertension*, 13, pp. 36-42.

3 Lin, Y.-S., Tang, C.-H., Yang, C.-Y.C., Wu, L.-S., Hung, S.-T., Hwa, H.-L., Chu, P.-H.

Effect of pre-eclampsiaeclampsia on major cardiovascular events among peripartum women in taiwan

(2011) *American Journal of Cardiology*, 107 (2), pp. 325-330. Cited 48 times.
doi: 10.1016/j.amjcard.2010.08.073

View at Publisher

4 Heida, K.Y., Franx, A., Van Rijn, B.B., Eijkemans, M.J.C., Boer, J.M.A., Verschuren, M.W.M., Oudijk, M.A., (...), Van Der Schouw, Y.T.

Earlier Age of Onset of Chronic Hypertension and Type 2 Diabetes Mellitus After a Hypertensive Disorder of Pregnancy or Gestational Diabetes Mellitus (Open Access)

(2015) *Hypertension*, 66 (6), pp. 1116-1122. Cited 56 times.
http://hyper.ahajournals.org/
doi: 10.1161/HYPERTENSIONAHA.115.06005

View at Publisher

5 Veerbeek, J.H.W., Hermes, W., Breimer, A.Y., Van Rijn, B.B., Koenen, S.V., Mol, B.W., Franx, A., (...), Koster, M.P.H.

Cardiovascular disease risk factors after early-onset preeclampsia, late-onset preeclampsia, and pregnancy-induced hypertension (Open Access)

(2015) *Hypertension*, 65 (3), pp. 600-606. Cited 95 times.
http://hyper.ahajournals.org/
doi: 10.1161/HYPERTENSIONAHA.114.04850

View at Publisher

6 Männistö, T., Mendola, P., Vääräsmäki, M., Järvelin, M.-R., Hartikainen, A.-L., Pouta, A., Suvanto, E.
Elevated blood pressure in pregnancy and subsequent chronic disease risk (Open Access)

(2013) *Circulation*, 127 (6), pp. 681-690. Cited 143 times.
doi: 10.1161/CIRCULATIONAHA.112.128751

View at Publisher

7 Valdiviezo, C., Garovic, V.D., Ouyang, P.
Preeclampsia and hypertensive disease in pregnancy: Their contributions to cardiovascular risk

(2012) *Clinical Cardiology*, 35 (3), pp. 160-165. Cited 39 times.
doi: 10.1002/clc.21965

View at Publisher

8 George, E.M., Granger, J.P.
Endothelin: Key mediator of hypertension in preeclampsia (Open Access)

(2011) *American Journal of Hypertension*, 24 (9), pp. 964-969. Cited 104 times.
doi: 10.1038/ajh.2011.99

View at Publisher

9 Kohan, D.E., Rossi, N.F., Inscho, E.W., Pollock, D.M.
Regulation of blood pressure and salt homeostasis by endothelin

(2011) *Physiological Reviews*, 91 (1), pp. 1-77. Cited 233 times.
http://physrev.physiology.org/content/91/1/1.full.pdf+html
doi: 10.1152/physrev.00060.2009

View at Publisher

10 Stow, L.R., Jacobs, M.E., Wingo, C.S., Cain, B.D.
Endothelin-1 gene regulation

(2011) *FASEB Journal*, 25 (1), pp. 16-28. Cited 105 times.
http://www.fasebj.org/content/25/1/16.full.pdf+html
doi: 10.1096/fj.10-161612

View at Publisher

11 Bernardi, F., Constantino, L., MacHado, R., Petronilho, F., Dal-Pizzol, F.
Plasma nitric oxide, endothelin-1, arginase and superoxide dismutase in pre-eclamptic women

(2008) *Journal of Obstetrics and Gynaecology Research*, 34 (6), pp. 957-963. Cited 43 times.
doi: 10.1111/j.1447-0756.2008.00860.x

View at Publisher

12 Silva, B.R., Pernomian, L., Bendhack, L.M.
Contribution of oxidative stress to endothelial dysfunction in hypertension
(Open Access)

(2012) *Frontiers in Physiology*, 3 DEC, art. no. Article 441. Cited 50 times.
http://www.frontiersin.org/Journal/FullText.aspx?ART_DOI=10.3389/fphys.2012.00441&x=y#h3
doi: 10.3389/fphys.2012.00441

View at Publisher

13 Lankhorst, S., Kappers, M.H.W., Van Esch, J.H.M., Danser, A.H.J., Van Den Meiracker, A.H.
Hypertension during vascular endothelial growth factor inhibition: Focus on nitric oxide, endothelin-1, and oxidative stress

(2014) *Antioxidants and Redox Signaling*, 20 (1), pp. 135-145. Cited 45 times.
doi: 10.1089/ars.2013.5244

View at Publisher

14 Verdonk, K., Saleh, L., Lankhorst, S., Smilde, J.E.I., Van Ingen, M.M., Garrelds, I.M., Friesema, E.C.H., (...), Danser, A.H.J.
Association Studies Suggest a Key Role for Endothelin-1 in the Pathogenesis of Preeclampsia and the Accompanying Renin-Angiotensin-Aldosterone System Suppression (Open Access)

(2015) *Hypertension*, 65 (6), pp. 1316-1323. Cited 67 times.
http://hyper.ahajournals.org/
doi: 10.1161/HYPERTENSIONAHA.115.05267

View at Publisher

15 George, E.M., Granger, J.P.
Linking Placental Ischemia and Hypertension in Preeclampsia: Role of Endothelin-1
(2012) *NIH Public Access*, 29, pp. 997-1003.

16 George, E.M., Granger, J.P.
Endothelin: Key mediator of hypertension in preeclampsia (Open Access)

(2011) *American Journal of Hypertension*, 24 (9), pp. 964-969. Cited 104 times.
doi: 10.1038/ajh.2011.99

View at Publisher

17 Kohan, D.E.
Endothelin, hypertension and chronic kidney disease: New insights

(2010) *Current Opinion in Nephrology and Hypertension*, 19 (2), pp. 134-139. Cited 75 times.
doi: 10.1097/MNH.0b013e328335f91f

View at Publisher

18 López Jaramillo, P., Arenas, W.D., García, R.G., Rincon, M.Y., López, M.
The role of the L-arginine-nitric oxide pathway in preeclampsia (Open Access)

(2008) *Therapeutic Advances in Cardiovascular Disease*, 2 (4), pp. 261-275. Cited 55 times.
doi: 10.1177/1753944708092277

View at Publisher

19 Baksu, B., Davas, I., Baksu, A., Akyol, A., Gulbaba, G.
Plasma nitric oxide, endothelin-1 and urinary nitric oxide and cyclic guanosine monophosphate levels in hypertensive pregnant women

(2005) *International Journal of Gynecology and Obstetrics*, 90 (2), pp. 112-117. Cited 40 times.
www.elsevier.com/locate/ijgo
doi: 10.1016/j.ijgo.2005.04.018

View at Publisher

20 Rapoport, R.M.

Nitric oxide inhibition of endothelin-1 release in the vasculature: In vivo relevance of in vitro findings (Open Access)

(2014) *Hypertension*, 64 (5), pp. 908-914. Cited 12 times.
http://hyper.ahajournals.org/
doi: 10.1161/HYPERTENSIONAHA.114.03837

View at Publisher

21 Bourque, S.L., Davidge, S.T., Adams, M.A.

The interaction between endothelin-1 and nitric oxide in the vasculature: New perspectives

(2011) *American Journal of Physiology - Regulatory Integrative and Comparative Physiology*, 300 (6), pp. 1288-1295. Cited 110 times.
http://ajpregu.physiology.org/content/300/6/R1288.full.pdf+html
doi: 10.1152/ajpregu.00397.2010

View at Publisher

Hidayatul, R.I.; Ismawi Department of Basic Medical Sciences, Kulliyyah of Medicine, International Islamic University Malaysia, Kuantan, Pahang, Malaysia; email:hidayatulradziah@iium.edu.my