



# Document details

< Back to results | 1 of 1

↗ Export ↓ Download 🖨 Print ✉ E-mail 📄 Save to PDF ☆ Add to List More... >

[Full Text](#) View at Publisher

Optics and Laser Technology  
Volume 115, July 2019, Pages 97-103

## Transition from saturable absorption to reverse saturable absorption of carmoisine dye under low-powered continuous wave laser excitation

(Article)

Abdullah, M.<sup>a,b</sup>, Bakhtiar, H.<sup>b,c</sup> ✉, Krishnan, G.<sup>b,c</sup>, Aziz, M.S.A.<sup>b,c</sup>, Danial, W.H.<sup>d</sup>, Islam, S.<sup>b,c</sup> 🔍

<sup>a</sup>Institute of Nano Optoelectronics Research and Technology (INOR), Universiti Sains Malaysia, USM, Penang 11800, Malaysia

<sup>b</sup>Laser Center, Ibnu Sina for Scientific and Industrial Research, Universiti Teknologi Malaysia, Johor Bahru, Johor 81300, Malaysia

<sup>c</sup>Department of Physics, Faculty of Sciences, Universiti Teknologi Malaysia, Johor Bahru, Johor, Malaysia

View additional affiliations ▾

### Abstract

▾ View references (31)

Unique nonlinear optics (NLO) properties i.e. intensity-dependent nonlinear absorption and refraction of carmoisine (food dye) is studied using a single beam z-scan technique. A switchover behavior from a saturable absorption (SA) to reverse saturable absorption (RSA) is observed by increasing concentration of carmoisine dye. The flip in the absorption response is attributed to the aggregated dye molecules under intense laser beam, which induces the formation of carmoisine dimers. In the UV-Vis absorption analysis, the appearance of two bands at higher concentration confirms the formation of carmoisine dimers. Fourier transform infrared spectroscopy (FTIR) suggests the intermolecular charge transfer (ICT) within the naphthyl-azo bonds. Huge magnitude of  $\chi^{(3)}$  is calculated in the order of  $10^{-5}$  esu due to ICT process within the dye molecules. Optical limiting (OL) behavior is observed with low OL action threshold  $\sim 0.01$  kW/cm<sup>2</sup> under continuous wave laser beam. The experimental findings shows that carmoisine dye has potential as an optical material for photonics applications such as an optical limiter under low-powered continuous wave laser. © 2019


### SciVal Topic Prominence ⓘ

Topic: Nonlinear optics | Refractive index | Coefficient  $\beta$

Prominence percentile: 95.505 ⓘ

### Chemistry database information ⓘ

#### Substances



#### Author keywords

Metrics ⓘ View all metrics >



PlumX Metrics ▾

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

[Set citation feed >](#)

### Related documents

Nonlinear properties of Acid Blue 29 Dye in CTAB/1-butanol/water microemulsion

Shavakandi, S.M., Sharifi, S. (2017) *Journal of Optoelectronics and Advanced Materials*

Nonlinear optical response of hydroxy substituted anthraquinone/PMMA thin films using z-scan technique

Zafar, S., Khan, H.Z., Khan, M.S. (2015) *Advanced Science Letters*

Third-order nonlinear optical properties of acid green 25 dye by Z-scan method

Jeyaram, S., Geethakrishnan, T. (2017) *Optics and Laser Technology*

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

## Indexed keywords

Engineering controlled terms:

Charge transfer Continuous wave lasers Dimers  
Fourier transform infrared spectroscopy Laser beams Molecules Nonlinear optics

Engineering uncontrolled terms

Carmoisine FTIR Optical limiters Optical switching Z-scan technique

Engineering main heading:

Laser excitation

## Funding details

Funding sponsor	Funding number	Acronym
-----------------	----------------	---------

Ministry of Higher Education, Malaysia

Universiti Teknologi Malaysia

### Funding text

The authors would like to thank the Malaysia Ministry of Education and Universiti Teknologi Malaysia for their financial support through Tier 1 with vote 18H67. Special thanks also to Universiti Teknologi Malaysia for the postdoctoral scheme under vote number 04E08 for Mundzir Abdullah. Corresponding author is grateful to the Malaysia Ministry of Education through the FRGS fund with vote 03EE89. Appendix A

ISSN: 00303992

CODEN: OLTC A

Source Type: Journal

Original language: English

DOI: 10.1016/j.optlastec.2019.01.032

Document Type: Article

Publisher: Elsevier Ltd

## References (31)

[View in search results format >](#)

- 1 Jeyaram, S., Geethakrishnan, T.  
Third-order nonlinear optical properties of acid green 25 dye by Z-scan method

(2017) *Optics and Laser Technology*, 89, pp. 179-185. Cited 30 times.  
doi: 10.1016/j.optlastec.2016.10.006

[View at Publisher](#)

- 2 Marder, S.R.  
Organic nonlinear optical materials: Where we have been and where we are going

(2006) *Chemical Communications*, (2), pp. 131-134. Cited 371 times.  
doi: 10.1039/b512646k

[View at Publisher](#)

- 3 Fernando, M.G.P.S., Wijewardena Gamalath, K.A.I.L.  
Modelling all-optical switching and limiting properties of AIs photonic crystals  
(2018) *Int. Lett. Chem. Phys. Astron.*, 77, pp. 1-14.