

IJOHS

International Journal of Orofacial and Health Sciences

A scientific journal published by Kulliyyah of Dentistry,IIUM,Malaysia







VOL. 1 ISSUE 1

2020

International Journal of Orofacial and Health Sciences

Editorial Board

Chief Editor	: Professor Dr. Zainul Ahmad Rajion
Editor	: Dr. Salwana Supa'at
	: Dr. Basma Ezzat Mustafa Alahmad
	: Assoc. Prof. Dr. Khairani Idah Mokhtar@Makhtar
	: Assoc. Prof. Dr. Solachuddin J.A. Ichwan
	: Assoc. Prof. Dr. Muhannad Ali Kashmoola
	: Dr. Widya Lestari
	: Dr. Azlini Ismail
	: Dr. Mohd Hafiz Arzmi
	: Dr. Mohamad Shafiq Mohd Ibrahim
	·

Aims and Scope:

International Journal of Orofacial and Health Sciences (IJOHS) is a peer reviewed biannual international journal dedicated to publish high quality of scientific research in the field of orofacial sciences, health sciences and interdisciplinary fields, including basic, applied and clinical research. The journal welcomes review articles, original research, case reports and letters to the editor. Areas that are covered include but are not limited to dental sciences, oral microbiology and immunology, oral maxillofacial and craniofacial surgery and imaging, dental stem cells and regenerative medicine, dental biomaterial, oral maxillofacial genetic and craniofacial deformities.

Published by:

Kulliyyah of Dentistry, International Islamic University Malaysia (IIUM), Indera Mahkota, 25200 Kuantan, Pahang, Malaysia.

Printed and distributed by:

Kulliyyah of Dentistry, International Islamic University Malaysia (IIUM), Indera Mahkota, 25200 Kuantan, Pahang, Malaysia.

All rights reserved; No part of this publication maybe reproduced, stored in a retrieval system, or transmitted in any form, or by any means, electronic or otherwise, without prior permission of the publisher.

CONTENTS

EDITORIAL	
Introduction to IJOHS	3
REVIEW ARTICLES	
Genetics of malocclusion: A review	4
ORIGINAL ARTICLES	
Potential antibacterial effects of flaxseed and Nigella sativa extracts on	11
Streptococcus pyogenes	
Dental treatment needs among patients undergoing screening at a	18
university-based dental institution in Kuantan, Pahang, Malaysia	
Analysis of the anti-cancer effect of ethyl-p-methoxycinnamate extracted	28
cekur (<i>Kaempferia galanga</i>) on cancer cell lines with wild-type and null p53	
Radiographic findings in panoramic radiographs of patients attending	34
Kulliyyah of Dentistry, IIUM	40
Isolation of Candida species in children and their biofilm-forming ability on	40
nano-composite surfaces	

Dental treatment needs among patients undergoing screening at a university-based dental institution in Kuantan, Pahang, Malaysia

Azlini Ismail^{1*}, Zurainie Abllah^{2,} Nur Aishah Muhammad Radhi³, Syazalina Musa³, Mohd Firdaus Akbar Abdul Halim⁴

¹Department of Fundamental Dental and Medical Sciences, Kulliyyah of Dentistry, International Islamic University Malaysia, Indera Mahkota, 25200 Kuantan, Pahang, Malaysia. ²Department of Paediatric Dentistry and Dental Public Health, Kulliyyah of Dentistry, International Islamic University Malaysia, Indera Mahkota, 25200 Kuantan, Pahang, Malaysia. ³Kulliyyah of Dentistry, International Islamic University Malaysia, Indera Mahkota, 25200 Kuantan, Pahang, Malaysia.

⁴Faculty Computer and Mathematical Sciences, Universiti Teknologi MARA Kelantan Branch, Kota Bharu Campus, Lembah Sireh, 15050 Kota Bharu, Kelantan, Malaysia.

Abstract

University-based dental institution in Malaysia receives large number of dental visits, however, dental treatment needs among patients attending this kind of institution is not usually reported. This study aimed to identify the trend of dental treatment needs in a university-based dental institution in Kuantan, Pahang situated in the East Coast region of Peninsular Malaysia. This cross-sectional study utilized secondary data, obtained from list of patients who underwent for screening at Outpatient Clinic, Kulliyyah of Dentistry, International Islamic University Malaysia from 1st January to 31st December 2016. Patient's age, gender, residential area and dental treatment needs were retrieved from the list. All patients were included except those with incomplete data, with old Malaysian identification card or foreign passport or those assigned to receive Orthodontic treatment. Descriptive statistics and Pearson's Chi Square test was run using SPSS® Version 20 software. Conservative care (22.7 %) was the major treatment need among 2,627 patients included in this study. Teenage and adult patients mostly required conservative care while the elderaged patients have major need for prosthodontics. Females outnumbered males in requiring all types of dental treatment, except for conservative care which was pre-dominantly required by males. Patients residing non-urban area majorly required all types of dental treatment except for endodontics and conservative care which were more frequently necessitated by patients from urban area. Conservative care was the major dental treatment need. The type of dental treatment needs has significant association with patient's age, gender and residential area.

Keywords: treatment need, screening, dental, age, gender

*Corresponding Author Email address: <u>dr_azlini@iium.edu.my</u> Tel: +6014-501 0081

Introduction

Need is classified into normative, felt, expressed or comparative needs according to Bradshaw (1972) taxonomy. Normative need for dental treatment is a category of need which is usually defined by the experts or professional. However, the process of dental screening usually takes into consideration the felt or

perceived need according to patients' complaints on their oral health conditions. Dental professional or a dental officer then assesses the suitability of dental treatment for the patients.

Dental treatment needs and the oral health status of Malaysians were periodically assessesed every ten years from 1990 to 2010 using National Oral Health Survey of Adults (NOHSA). Since 2010, the oral health status and dental treatment needs in Malaysia have not been assessed at any national scale study. In the meantime, there are local studies that studied on specific dental treatment need. for instance the orthodontic treatment need among various study populations including of school children aged 12 and 16 years old (Zreaqat et al., 2013), adolescents aged 13 to 14 years old (Zamzuri et al., 2014), adults aged 20 to 70 years old (Ravindranath et al., 2017) as well as among the special need people with Down Syndrome (Abdul Rahim et al., 2014). Nevertheless, there are also researches that analysed on general dental treatment needs in certain populations in Malaysia, for instance, among the special need children in Negeri Sembilan, a state which is situated in the centre of Peninsular Malaysia (Mokhtar et al., 2016); among normal (Oo et al., 2011) and hearingimpaired school children (Rahman et al., 2015) as well as among the elderly (Sinor, 2013; Sinor et al., 2018) in Kelantan, a state located in the North East Region of Peninsular Malaysia.

However, there is no specific study that has contextualizes the dental treatment needs based on any universitybased Malaysian dental institution. Dental Kulliyyah Clinic at of Dentistry. International Islamic University Malaysia which is established in 2006 is the only university-based dental service provider in the city of Kuantan, Pahang, Malaysia and it is one of the two university-based dental centres that cater for the East Coast

region in Malaysia. The number of patients may reach up to 10,000 visits per year indicative for a high demand and need for dental treatment among the surrounding community. In view of this, this study aims to analyse the trend of major dental treatment need in this population.

Materials and Methods

Study Location and Data Collection

Ethical approval (IREC 762) was obtained from IIUM Research Ethics Committee. This study was conducted at IIUM Dental Outpatient Clinic in Kuantan, Pahang. Kuantan is the capital city of Pahang, situated at the East Coast region of Peninsular Malaysia. Our study population was all first-visit patients undergoing screening at IIUM Dental Outpatient Clinic from 1st January 2016 until 31st December 2016. Information on patient's age, gender, residential area and the type of dental treatment needs was retrieved from Outpatient Clinic database. Type of treatment needs were decided by dental officer in-charge at this clinic after carrying out the usual procedure of screening and dental charting for patient's oral health condition. Treatment need was then notified in the patient's folder, and was remarked into the Patient Waiting List of any departments according to their treatment needs. Patients with incomplete data or with old Malaysian identification card or foreign passport and those attended Orthodontic department are excluded.

For demographic profile analysis, patients were classified based on their age group, gender and residential area in Kuantan. Age were classified into six groups of 14 years old and below, 15 to 19 years old, 20 to 34 years old, 35 to 44 years old, 45 to 64 years old, and 65 years old and above. Residential areas were classified into urban and non-urban areas according to Kuantan Municipal Council. Dental treatment needs was classified as stated in the clinic database; endodontics (anterior), endodontics (posterior), fixed prosthetic (bridge), fixed prosthetic (crown), partial denture, full denture, paediatric dentistry, oral surgery, general dental practice, oral maxillofacial, conservative care and periodontal need.

Statistical Analysis

Descriptive statistics was used to analyze the demographic profiles of this population. Association of demographic background (age, gender and residential areas) with the type of dental treatment needs was analysed using Chi-Square test in SPSS[®] software Version 20.

Results

Demographic Profiles

Demographic profiles of first-visit patients at IIUM Dental Outpatient Clinic for the year 2016 are tabulated in Table 1. The major age group of patients was from the age of 20 to 34 years old (39.5 %) while the least number of patients was from the age group of 65 and above (4.2 %). The mean age of the patients was 32 ± 17.61 years old. Female patients (55.5 %) outnumbered male patients (44.5 %) with majority of patients residing non-urban areas (37.6 %).

Dental Treatment Needs

Figure 1 shows distribution of dental treatment needs in which the highest treatment need was conservative care (22.7 %) while the lowest treatment need was for oral maxillofacial care (0.6 %).

Association of demographic profiles with types of dental treatment needs

Table 2 shows cross tabulation of age with the type of dental treatment needs. There is significant association between age, X^2 (55, N = 2,627) =

3,466.9, p < .0001 with the type of dental treatment needs. Paediatric patients (age group of \leq 14) were almost exclusively assigned to receive paediatric dentistry care (97.5 %). Younger-age patients (age group of 15-19, 20-34 and 35-44) mostly required conservative care (38.9 %, 30.9 % and 30.3 %, respectively), followed by periodontal need (17.2 %, 24.5 % and 17.0 %, respectively). Elder-age patients (age group of 45-64 and \geq 65) have major need for prosthodontics with 30.5 % and 40.5 %, respectively for partial dentures and 9.4 % and 31.5 %, respectively for full dentures. The need for partial dentures increases with increasing age, this is noticeable from the age group 35-44 (8.9 %), 45-64 (30.5 %) and ≥65 (40.5 %). This is also similar to the need for full dentures that begins from the age group of 45-64 (9.4 %) and drastically increased in the age group of ≥65 (31.5 %).

In contrast, the need for oral surgery decreases with increasing age; the need was highest among patients from the age groups of 20-34 (17.9 %), 35-44 $(16.7 \%), 45-64 (9.6 \%) \text{ and } \ge 65 (8.1 \%).$ The need for oral maxillofacial care is consistently low compared to other treatment needs across all age groups. This study also found that types of dental treatment needs also significantly differed by gender, X^2 (11, N = 2,627) = 49.8, p < .0001. As cross tabulated in Table 3, females outnumbered males in all types of dental treatment need. except for periodontal which need was predominated by males. The relation between residential areas and the type of dental treatment need was also significant, *X*² (11, N = 2,627) = 112.6, p <.0001.

As shown in Table 4, people residing non-urban areas outnumbered those residing urban areas in getting fixedprosthetic (bridge) (50.6 %), fixedprosthetic (crown) (58.5 %), partial denture (60.0 %), full denture (79.1 %), paediatric dentistry care (62.3 %), oral surgery (57.6 %), general dental practice (52.6 %) and periodontal care (51.7 %).

Demographic Profiles	n (% of total)
Age	
≤14	434(16.5 %)
15 to 19	157(6.0 %)
20 to 34	1,037 (39.5 %)
35 to 44	347 (13.2 %)
45 to 64	541 (20.6 %)
≥65	111 (4.2 %)
	Total = 2,627 (100.0 %)
Gender	
Male	1,168 (44.5 %)
Female	1,459 (55.5 %)
	Total = 2,627(100.0 %)
Residential Areas	
Urban	1,251 (47.6%)
Non-urban	1,376 (52.4%)
	Total=2,627 (100.0%)

Table 1. Demographic profiles of patients underwent for screening at IIUM Dental Outpatient Clinic for the year 2016.



Dental Treatment Need

Figure 1. Distribution of dental treatment needs among patients underwent for screening at IIUM Dental Outpatient Clinic for the year 2016.

Age	Dental Treatment Needs											Statistical		
group	а	b	С	d	е	f	g	h	i	j	k	I	Total	Analysis
≤14														
Count	0	0	0	1	0	0	423	0	0	3	2	5	434	
% within														
age	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	97.5%	0.0%	0.0%	0.7%	0.5%	1.2%	100.0%	
% within														
treatment	0.0%	0.0%	0.0%	1.1%	0.0%	0.0%	96.6%	0.0%	0.0%	1.3%	0.3%	1.2%	16.5%	
15 to 19														
Count	13	4	1	4	1	0	15	4	4	23	61	27	157	X² (55,
% within														N =
age	8.3%	2.5%	0.6%	2.5%	0.6%	0.0%	9.6%	2.5%	2.5%	14.6%	38.9%	17.2%	100.0%	2,627) =
% within														3,466.9,
treatment	15.3%	6.2%	2.9%	4.3%	0.4%	0.0%	3.4%	1.3%	23.5%	10.1%	10.2%	6.6%	6.0%	p <.0001
20 to 34														
Count	49	32	10	37	23	0	0	186	11	115	320	254	1,037	
% within														
age	4.7%	3.1%	1.0%	3.6%	2.2%	0.0%	0.00%	17.9%	1.1%	11.1%	30.9%	24.5%	100.0%	
% within														
treatment	57.6%	49.2%	28.6%	39.4%	8.7%	0.0%	0.00%	60.2%	64.7%	50.4%	53.6%	62.3%	13.2%	

Table 2. Cross tabulation of dental treatment needs versus age of patients underwent for screening at IIUM Dental Outpatient Clinic for the year 2016

Note: Treatment a-Endodontics (Anterior), b-Endodontics (Posterior) c-Fixed-prosthetics (Bridge), d-Fixed-Prosthetics (Crown), e- Partial Denture, f-Full Denture, g-Paedodontics, h-Oral Surgery, i-Oral Maxillofacial, j-General Dental Practice, k-Conservative Care and I-Periodontal Need. Chi-square test, X^2 (55, N = 2,627) = 3,466.9, p < .0001

Age	Dental Treatment Needs													Statistical
group	а	b	С	d	е	f	g	h	i	j	k	I	Total	Analysis
35 to 44														
Count	13	15	8	21	31	0	0	58	1	36	105	59	347	
% within														
age	3.7%	4.3%	2.3%	6.1%	8.9%	0.0%	0.0%	16.7%	0.3%	10.4%	30.3%	17.0%	100.0%	
% within														
treatment	15.3%	23.1%	22.9%	22.3%	11.17%	0.0%	0.0%	18.8%	5.9%	15.8%	17.6%	14.5%	39.5%	
45 to 64														
Count	9	14	15	30	165	51	0	52	0	44	105	56	541	X^{2} (55
% within														N =
age	1.7%	2.6%	2.8%	5.5%	30.5%	9.4%	0.0%	9.6%	0.0%	8.1%	19.4%	10.4%	100.0%	2.627)
% within														=
treatment	10.6%	21.5%	42.9%	31.9%	62.3%	59.3%	0.0%	16.8%	0.0%	19.3%	17.6%	13.7%	20.6%	3,466.9,
≥65														p <.0001
Count	1	0	1	1	45	35	0	9	1	7	4	7	111	
% within														
age	0.9%	0.0%	0.9%	0.9%	40.5%	31.5%	0.0%	8.1%	0.9%	6.3%	3.6%	6.3%	100.0%	
% within														
treatment	1.2%	0.0%	2.9%	1.1%	17.0%	40.7%	0.0%	2.9%	5.9%	3.1%	0.7%	1.7%	4.2%	
Total	85	65	35	94	265	86	438	309	17	228	597	408	2,627	

Note: Treatment a-Endodontics (Anterior), b-Endodontics (Posterior) c-Fixed-prosthetics (Bridge), d-Fixed-Prosthetics (Crown), e- Partial Denture, f-Full Denture, g-Paedodontics, h-Oral Surgery, i-Oral Maxillofacial, j-General Dental Practice, k-Conservative Care and I-Periodontal Need. Chi-square test, X^2 (55, N = 2,627) = 3,466.9, p < .0001

	Dental Treatment Needs											Statistical		
Gender	а	b	С	d	е	f	g	h	i	j	k	I	Total	Analysis
Male														
Count	32	26	15	32	93	37	178	152	2	93	319	189	1,168	
% within														
gender	2.7%	2.2%	1.3%	2.7%	8.0%	3.2%	15.2%	13.0%	0.2%	8.0%	27.3%	16.2%	100.0%	<i>X</i> ² (11,
% within														N =
treatment	37.6%	40.0%	42.9%	34.0%	35.1%	43.0%	40.6%	49.2%	11.8%	40.8%	53.4%	46.3%	44.5%	2,627)
Female														= 49.8,
Count	53	39	20	62	172	49	260	157	15	135	278	219	1,459	p<.0001
% within														
gender	3.6%	2.7%	1.4%	4.2%	11.8%	3.4%	17.6%	10.8%	1.0%	9.3%	19.1%	15.0%	100.0%	
% within														
treatment	62.4%	60.0%	57.1%	66.0%	64.9%	57.0%	59.4%	50.8%	88.2%	59.2%	46.6%	53.7%	55.5%	
Total	85	65	35	94	265	86	438	309	17	228	597	408	2,627	l

Table 3. Cross tabulation of dental treatment needs versus gender of patients underwent for screening at IIUM Dental Outpatient Clinic for the year 2016

Note: Treatment a-Endodontics (Anterior), b-Endodontics (Posterior) c-Fixed-prosthetics (Bridge), d-Fixed-Prosthetics (Crown), e- Partial Denture, f-Full Denture, g-Paedodontics, h-Oral Surgery, i-Oral Maxillofacial, j-General Dental Practice, k-Conservative Care and I-Periodontal Need. Chi-square test, X^2 (11, N = 2,627) = 49.8, p < .0001

Residen-	Dental Treatment Needs												Statistical	
tial Areas	а	b	С	d	е	f	g	h	i	j	k	I	а	Analysis
Urban														
Count	43	38	17	42	106	18	165	131	10	108	376	197	1,251	
% within														
gender	3.4%	3.0%	1.4%	3.4%	8.5%	1.4%	13.2%	10.5%	0.8%	8.6%	30.1%	15.7%	100.0%	
% within														<i>X</i> ² (11,
treatment	50.6%	58.5%	48.6%	44.7%	40.0%	20.9%	37.7%	42.4%	58.8%	47.4%	63.0%	48.3%	47.6%	N =
Non-urban														2,627)
Count	42	27	18	52	159	68	273	178	7	120	221	211	1,376	_ 112.6.
% within														p<.0001
gender	3.1%	2.0%	1.3%	3.8%	11.6%	4.9%	19.8%	12.9%	0.5%	8.7%	16.1%	15.3%	100.0%	-
% within														
treatment	49.4%	41.5%	51.4%	55.3%	60.0%	79.1%	62.3%	57.6%	41.2%	52.6%	37.0%	51.7%	52.4%	
Total	85	65	35	94	265	86	438	309	17	228	597	408	2,627	

Table 4. Cross tabulation of dental treatment needs versus residential area of patients underwent for screening at IIUM Dental Outpatient Clinic for the year 2016

Note: Treatment a-Endodontics (Anterior), b-Endodontics (Posterior) c-Fixed-prosthetics (Bridge), d-Fixed-Prosthetics (Crown), e- Partial Denture, f-Full Denture, g-Paedodontics, h-Oral Surgery, i-Oral Maxillofacial, j-General Dental Practice, k-Conservative Care and I-Periodontal Need. Chi-square test, *X*² (11, N = 2,627) = 112.6, p < .0001.

Discussion

The predominant need among teenage to adult patients was conservative care, followed closely by periodontal need. The main reason for the high demand for conservative and periodontal care was perhaps due to the high prevalence of caries and gingivitis among Malaysian adults. According to National Oral Health Survey of Adult (NOHSA) in 2010, about 94.0 % of Malaysian adult population had periodontal disease while 88.9 % of Malaysian population had dental caries experience.

This study also shows that majority of the first-visit patients at this clinic were females. This scenario was also observed in Turkish population (Pekiner et al., 2010), and in fact, a previous study in Southern China has also shown that females were more likely to visit dental service provider (Lo et al., 2001). Other than that, NOHSA 2010 also reported a higher proportion of females (53.1 %) that sought after dental treatment at public dental provider compared to males (49.3 %). This female predilection might be due to the difference of oral health between men and women, especially at certain age. The disparity in oral health increases between the genders as a population ages, which may relate to the combination of reproductive hormones influences, pregnancy, diet, as well as morning sickness during pregnancy in women (Lukacs, 2011). Other than pregnancy, menopause is also associated with increased risk of oral health complication, especially in women who developed osteoporosis (Branch-Elliman, 2012). Another plausible reason is that women visit more dental and oral health service providers because they usually are more attentive towards the aesthetics inclusive of the teeth (Akbar et al., 2019).

In addition to gender, age also have significant association with the type of dental treatment need. This study shows that younger-age patients mostly required conservative care while the elderage patients have major need for prosthodontics, either partial or full agreement dentures. In with this. prosthodontics was the most required treatment need among the elderly patients in Turkish (Pekiner et al., 2010) as well as in Northeast China (Liu et al., 2015) population. This study also shows that the need for partial and full dentures increases with increasing age. This is actually consistent with findings in NOHSA 2010 in which the study has shown that the number of edentulous significantly increases from the age of 35 (1.1 %) to the age 65 (35.6 %) and 75 (53.3 %). When the patient becomes edentulous, then the need for tooth extraction is nullified. The need for oral maxillofacial care which includes treatment for temporomandibular joint symptoms, operculectomy, pericoronitis and bruxism is consistently low compared to other treatment need across all age groups.

Residential area also has significant association with the type of dental treatment need. This study shows that people residing non-urban areas outnumbered those residing urban areas to receive fixed prosthetics of either bridge or crown; partial and full dentures; paedodontic care; periodontal treatment; and being referred to oral surgery and general dental practice departments. In general, people living in rural areas tend to have lower oral health problem, has more caries and fewer teeth compared to the urban residents (Akbar et al., 2019). However, the geographical location of this dental institution and the convenient of transportation made this institution still accessible to both the urban and nonurban residents.

Conclusion

Conservative care was the major dental treatment need identified in this study population. The type of dental treatment

need has significant association with patient's age, gender and residential area. substantially This finding helps in understanding of dental treatment need, within especially the context of а university-based dental care provider in the East Coast region of Peninsular Malaysia.

Acknowledgement

We are also thankful to Mrs. Sa'diah, the Head Officer of Kuantan Municipal Council, as well as to our dental surgical assistant for her support during data retrieving at the clinic.

References

- Abdul Rahim, F. S., Mohamed, A. M., Nor, M. M., & Saub, R. (2014). Malocclusion and orthodontic treatment need evaluated among subjects with Down syndrome using the Dental Aesthetic Index (DAI). *The Angle Orthodontist*, 84(4), 600-606.
- Akbar, F. H., Pasinringi, S., & Awang, A. H. (2019). Relationship between health service access to dental conditions in urban and rural areas in Indonesia. *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*, 19, 1-7.
- Bradshaw, J. (1972). Taxonomy of social need. In G. McLahlan (Ed.), *Problems and progress in medical care*. London: Oxford University Press. 69-82.
- Branch-Elliman, D. (2012). A gender-based approach to oral health changes across the lifespan. University of Pennsylvannia, USA.
- Liu, L., Zhang, Y., Wu, W., & Cheng, R. (2015). Characteristics of dental care-seeking behavior and related sociodemographic factors in a middle-aged and elderly population in Northeast China. *BMC Oral Health*, 15(1), 66.
- Lo, E., Lin, H., Wang, Z., Wong, M., & Schwarz, E. (2001). Utilization of dental services in Southern China. *Journal of Dental Research*, 80(5), 1471-1474.
 Lukacs, J. R. (2011). Sex differences in dental
- Lukacs, J. R. (2011). Sex differences in dental caries experience: Clinical evidence, complex etiology. *Clinical Oral Investigations*, 15(5), 649-656. Mokhtar, S. M., Jalil, L. A., Noor, N. M., Tan, B.,
- Mokhtar, S. M., Jalil, L. A., Noor, N. M., Tan, B., Shamdol, Z., & Hanafiah, H. A. (2016). Dental status and treatment needs of special needs children in Negeri Sembilan, Malaysia. *World Journal of Research and Review*, 2(6), 64-70.

- National Oral Health Survery of Adults (NOHSA). Oral Health Division, Ministry of Health, Malaysia.
- Oo, M. M. T., Naing, L., Mani, S. A., & Ismail, A. R. (2011). Dental caries experience and treatment needs in the mixed dentition in North East Malaysia. *Archives of Orofacial Sciences*, 6(2), 41-48.
- Pekiner, F., Gumru, B., Borahan, M. O., & Aytugar, E. (2010). Evaluation of demands and needs for dental care in a sample of the Turkish population. *European Journal of Dentistry*, 4(2), 143-149.
- Rahman, N. A., Yusoff, A., Daud, M. K. M., & Kamaruzaman, F. N. (2015). Salivary parameters, dental caries experience and treatment needs of hearing-impaired children in a special school for deaf in Kelantan, Malaysia. *Archives of Orofacial Sciences*, 10(1), 17-23.
- Ravindranath, S., En, J. T. S., & Heng, A. P. K. (2017). Orthodontic treatment need and self-perceived psychosocial impact of dental esthetics in a university adult population in Malaysia. *Journal of Indian Orthodontic Society*, 51(2), 69-74.
- Sinor, M. Z. (2013). Oral health assessment among elderly staying in shelter (Rumah Seri Kenangan), Kelantan, Malaysia. *Age*, 71, 8-38.
- Sinor, M. Z., Ahmad, B., Ariffin, A., & Hassan, A. (2018). Main medical illness, oral health complaints and treatment in elderly patients attending Hospital Universiti Sains Malaysia (HUSM) Dental Clinic. International Medical Journal, 25(4), 230-232.
- Zamzuri, S. Z. M., Razak, I. A., & Esa, R. (2014). Normative and perceived need for treatment of malocclusion among Malaysian adolescents. *Sains Malaysiana*, 43(7), 1037-1043.
- Zreaqat, M., Hassan, R., Ismail, A. R., Ismail, N. M., & Aziz, F. A. (2013). Orthodontic treatment need and demand among 12-and 16 yearold school children in Malaysia. Oral Health and Dental Management, 12(4), 217-221.

Instruction to Authors

Submission

Complete manuscripts, written in English along with tables and illustrations should be submitted as softcopy to <u>zainulrajion@iium.edu.my</u>

Conditions

All manuscripts submitted for publication must be accompanied by a cover letter declaring originality of the work and certifies that the manuscript has not been published or submitted elsewhere, free from conflict of interest and conduct of research in accordance to ethical established quidelines for human subjects and animal welfare. In addition, a statement of agreement of all authors to transfer all the copyrights to the journal should also accompany the manuscript.

A copy of the certification form, which certifies that the manuscript has not been published or submitted before and all authors have contributed and are in agreement with the content of the manuscript, must be submitted to the journal's editorial office by uploading it as a PDF file.

Manuscripts submitted for publication are understood to be offered only to **IJOHS** and which have not been sent to other journals for consideration. There is no publication fee to submit or publish content in **IJOHS**.

Manuscript should be typed in **Arial, font** size 11 with 1.5 spacing.

The manuscript should be divided into the following sections with the sequence; Title Page, Abstract, Introduction, Materials and Methods, Results, Discussion, Acknowledgements, References, Tables, Illustrations and Figure Legends must be incorporated near to the text they are cited

Title Page:

Title page should be arranged in the following order: (1) a concise and informative title not exceeding 80 characters; (2) authors' full names (without degrees and titles) and affiliation including city and country. Superscript numbers may be used to affiliate authors to different departments or institutions; and (3) a complete mailing address, telephone, fax and e-mail address of the corresponding author.

Abstract:

Abstract should be unstructured and should not exceed 250 words. It should be a self-standing summary of the work that may be republished by information retrieval services. A maximum of 5 keywords should be listed at the end of the abstract.

Introduction:

Introduction should briefly and clearly describe the background and objective(s) of the study. Avoid exhaustive review of the literature and include only relevant recent studies.

Materials and Methods:

This section should be described in sufficient details so that it is reproducible. It should include technical information such as the study design, and specific procedures. Sub-headings may be used to enhance the clarity or to categorize the Established methods. methods or procedures should be named and cited. New methods or modifications of old methods should be described with complete details. Authors are advised to use generic names and terms rather than commercial names. Statistical methods applied, software used must also be stated clearly and concisely.

Results:

This section should be presented in a logical sequence in text, tables and illustrations. Subheadings may be used

to enhance clarity or to call attention to the most significant findings. Data appearing in tables or figures may be summarized but not duplicated in the text. Tables and figures should be numbered in the order in which they are described and cited in the text. Example, For tables, examples are Table 1, Table 2 and for figures, examples are Figure 1, Figure 2.

Discussion:

This section should summarize, explain and interpret the results with a scientifically critical view of previously published works in the field. Avoid repetition of the data already presented in the result section.

Acknowledgements:

This section is to be given to those who have provided assistance to the project and sponsors who have given financial support and funds.

References:

This section should list all sources cited in the paper. The citations should be arranged in alphabetical order by the surname of the first author without numbering in the text. Cite article as e.g. (Ahmad, 2016) for a single author, (Chu & Kim, 2019) for two authors and (Karim *et al.*, 2013) for more than three authors.

Standard journal article

References to journals should provide the name(s) of the author(s), year, title of the paper and journal, volume and issue number, page numbers. Please provide full name of journal. If there are more than six authors, the first six authors should be listed and followed by *et al.* in the list of References.

Sudbery, P. E. (2011). Growth of *Candida albicans* Hyphae. *Nature Reviews Microbiology*, 9(10), 737. <u>Book</u>

References to books/monographs should give the name(s) of the author(s), year, title of book, edition number, place of publication, publisher and page numbers. Conn, E.E., Stumpf, P.K., Brueing, G., Doi, R.H. (1987). *Outlines of Biochemistry*, 3rd edn. New York: John Wiley & Sons, pp. 45–52.

Chapter in book

References to chapter in books/monographs should give the name(s) of the author(s), year, chapter title, editors, title of book, edition number, place of publication, publisher and page numbers.

Fejerskov O, Nyvad B, Kidd EAM (2003). Clinical and histological manifestations of dental caries. In: Fejerskov O, Kidd EAM (eds.), *Dental Caries – The Disease and Its Clinical Management*. London: Blackwell Munksgaard, pp. 71–97.

Internet resources

Authors are required to include as much electronic retrieval information as needed for others to locate the sources they have cited.

López-Jornet P (2006). Labial mucocele: a study of eighteen cases. The Internet Journal of Dental Science, 3(2). Retrieved 7 February 2007, from http://www.ispub.com/ostia/index.ph p?xmlFilePath=journals/ijds/vol3n2/labial. xml

Tables

Each table should have a suitable title with footnotes wherever necessary. Do not use vertical lines in the table.

Illustrations

All figures and illustrations could be original photographs, artwork or highquality digital images (submitted as CMYK - 8 bits per channel in TIFF format). Images must be at least 600 by 450 pixels (proportional height) in size when in landscape orientation with a resolution of at least 300 pixels per inch. Graphs should be approximately 500 pixels wide so that all labeling can be read with data points clearly visible. Figures should be numbered consecutively in the order in which they appear in the manuscript, using Arabic numerals. A list of figure legends must be included on a separate page following the illustrations. The legend should explain each figure in detail. All figures will be printed as black and white. Colour figures will only appear in the PDF file.

Page proofs

On acceptance of the manuscripts for publication, page proofs should be reviewed meticulously by the contributors. Changes made in proof should be limited to the correction of typographical errors. Proofs must be returned for publication with corrections and responses to queries on the date specified by the Editor.

