



# Document details

[Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)[View at Publisher](#)International Journal of Preventive Medicine [Open Access](#)

Volume 11, Issue 1, 1 January 2020, Article number 58

## A snapshot of pre - And post-intervention changes among cardiovascular disease patients participating in the new smoking cessation program (Article) [\(Open Access\)](#)

Azmi, J.<sup>a</sup>, Nurumal, M.<sup>b</sup> , Nik Mohamed, M.<sup>c</sup>, Ab Rahman, N.<sup>c</sup> <sup>a</sup>Department of Medical Surgical Nursing, Kulliyah of Nursing, International Islamic University Malaysia (IIUM), Selangor, Malaysia<sup>b</sup>Department of Critical Care Nursing, Kulliyah of Nursing, IIUM, Kuantan, Pahang, 25200, Malaysia<sup>c</sup>Department of Pharmacy Practice, Kulliyah of Pharmacy, IIUM, Kuantan, Pahang, 25200, Malaysia

### Abstract

[View references \(16\)](#)

**Background:** The study evaluates the changes of pre and post stages of behavioral change, motivation level, and smoking status among cardiovascular disease patients, participating in the new smoking cessation program in Hospital Tengku Ampuan Afzan, Kuantan. **Methods:** A total of 65 adult cardiac patients were randomly distributed into two groups, i.e., intervention and control group, who were baseline smokers and assessed in three phases. Initially, the first, second, and third phase collected the information about their demographic details, their smoking status, and smoking status using cotinine amylase strip, respectively. **Results:** It showed that behavioral change was significant for the control group ( $P$  value = 0.031), while motivation level improved for both groups (i.e., control,  $P$  value = 0.000 and intervention group,  $P$  value = 0.001). The smoke quitting status percentage was higher for intervention group 41.9% and lower for control group 20.6%; however, the  $P$  value was insignificant for both control group (1.000) and intervention group (0.250). **Conclusions:** This study suggests a need for more personal testimonial videos to focus on other smoking-related diseases. © 2020 BMJ Publishing Group. All rights reserved.

### Author keywords

[Cardiovascular disease](#) [intervention](#) [tobacco cessation program](#)

### Indexed keywords

EMTREE drug terms: [amylase](#) [carbon monoxide](#) [cotinine](#)

EMTREE medical terms:

[adult](#) [Article](#) [behavior change](#) [cardiovascular disease](#) [controlled study](#)  
[coronary artery disease](#) [current smoker](#) [educational status](#) [employment status](#) [ex-smoker](#)  
[human](#) [intervention study](#) [major clinical study](#) [Malaysia](#) [marriage](#) [motivation](#)  
[smoking](#) [smoking cessation](#) [smoking cessation program](#) [videorecording](#)

### Chemicals and CAS Registry Numbers:

amylase, 9000-90-2, 9000-92-4, 9001-19-8; carbon monoxide, 630-08-0; cotinine, 486-56-6

[Metrics](#) [View all metrics >](#)

### PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

### Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

### Related documents

Animation production and design based on metro television advertising LED

Li, X. (2018) *Journal of Advanced Oxidation Technologies*

Increasing the dose of television advertising in a national antismoking media campaign: Results from a randomised field trial

McAfee, T. , Davis, K.C. , Shafer, P. (2017) *Tobacco Control*

Does digital video advertising increase population-level reach of multimedia campaigns? Evidence from the 2013 tips from former smokers campaign

Davis, K.C. , Shafer, P.R. , Rodes, R. (2016) *Journal of Medical Internet Research*

View all related documents based on references

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

## References (16)

[View in search results format >](#) All     Export     Print     E-mail     Save to PDF     Create bibliography

- 1 Samaan, Z., Nowacki, B., Schulze, K., Magloire, P., Anand, S.S.  
Smoking cessation intervention in a cardiovascular hospital based clinical setting  
([Open Access](#))  
(2012) *Cardiovascular Psychiatry and Neurology*, art. no. 970108. Cited 6 times.  
doi: 10.1155/2012/970108  
[View at Publisher](#)
- 
- 2 Farrelly, M.C., Duke, J.C., Davis, K.C., Nonnemaker, J.M., Kamyab, K., Willett, J.G., Juster, H.R.  
Promotion of smoking cessation with emotional and/or graphic antismoking advertising  
(2012) *American Journal of Preventive Medicine*, 43 (5), pp. 475-482. Cited 68 times.  
doi: 10.1016/j.amepre.2012.07.023  
[View at Publisher](#)
- 
- 3 McAfee, T., Davis, K.C., Alexander Jr., R.L., Pechacek, T.F., Bunnell, R.  
Effect of the first federally funded US antismoking national media campaign  
(2013) *The Lancet*, 382 (9909), pp. 2003-2011. Cited 167 times.  
<http://www.journals.elsevier.com.ezproxy.um.edu.my/the-lancet/>  
doi: 10.1016/S0140-6736(13)61686-4  
[View at Publisher](#)
- 
- 4 Wakefield, M.A., Loken, B., Hornik, R.C.  
Use of mass media campaigns to change health behaviour  
(2010) *The Lancet*, 376 (9748), pp. 1261-1271. Cited 868 times.  
doi: 10.1016/S0140-6736(10)60809-4  
[View at Publisher](#)
- 
- 5 (2017) *Noncommunicable Diseases Country Profiles 2017*. Cited 3 times.  
WHO
- 
- 6 Lancaster, T., Stead, L.F.  
Individual behavioural counselling for smoking cessation  
(2017) *Cochrane Database of Systematic Reviews*, 2017 (3), art. no. CD001292. Cited 137 times.  
<http://as.wiley.com.ezproxy.um.edu.my/WileyCDA/Brand/id-6.html>  
doi: 10.1002/14651858.CD001292.pub3  
[View at Publisher](#)