



# Document details

< Back to results | 1 of 1

↗ Export ⬇ Download 🖨 Print ✉ E-mail 📄 Save to PDF ☆ Add to List More... >

View at Publisher

Egyptian Journal of Forensic Sciences **Open Access**  
Volume 10, Issue 1, 1 December 2020, Article number 15

## Chemical profiling of volatile organic compounds from shoe odour for personal identification (Article) **(Open Access)**

Mohd Kamal, M.H.<sup>a</sup> ✉, Zakaria, Y.<sup>b</sup> ✉, Md Isa, M.L.<sup>c</sup> ✉, Nik Hassan, N.F.<sup>a</sup> ✉ 🔍

<sup>a</sup>Forensic Science Programme, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia

<sup>b</sup>Biomedicine Programme, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia

<sup>c</sup>Department of Basic Medical Sciences, Kulliyah of Nursing, International Islamic University of Malaysia, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, Kuantan, Pahang, Malaysia

### Abstract

∨ View references (46)

**Background:** Body odour of an individual has a unique composition due to the combined influences of genetic, dietary, and environmental factors. This exploratory study was conducted to investigate the feasibility of obtaining unique chemical signatures of individuals from the shoe odour as a human chemical fingerprint for forensic identification. **Methods:** Five adult males and five adult females participated in this study and provided with two pairs of new canvas sports shoes. The participants were asked to do strenuous activities while wearing the shoes for 2 weeks. Two different extraction methods: swabbing technique, followed with direct extraction using different solvents, and air passive sampling technique, were compared for the effectiveness to extract the volatile organic compounds (VOCs) from the shoes. The variations of VOCs were investigated using gas chromatography coupled with flame ionisation detector (GC-FID), and the resultant chromatogram profiles collected from the shoe odour were further studied for individual identification purposes. In addition, principal component-discriminant analyses (PCA-DA) were then carried out on the GC dataset. **Results:** This study demonstrated that air passive sampling technique using methanol as the extraction solvent was effective for the detection of shoe odour. PCA-DA had successfully distinguished GC profile patterns and discriminated the shoe odour samples accordingly to their owners. The accuracy of DA for classification was 100%, with all samples were classified correctly to their groups. **Conclusion:** The chemical compounds of shoe odour have the potential to be utilised to aid the forensic investigation by identifying and narrowing down the possible suspect if shoes were recovered as evidence at the crime scene. © 2020, The Author(s).

### SciVal Topic Prominence ⓘ







Topic: Odors | Comprehensive Two-Dimensional Gas Chromatography | Deodorant Agent

Prominence percentile: 82.374 ⓘ

### Chemistry database information ⓘ

#### Substances

View all substances (9)



Metrics ⓘ View all metrics >



PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

### Related documents

Human scent samples for chemical analysis

Pojmanová, P., Ladislavová, N., Škeříková, V. (2020) *Chemical Papers*

New methods for field collection of human skin volatiles and perspectives for their application in the chemical ecology of human-pathogen-vector interactions

Dormont, L., Bessiere, J.-M., McKey, D. (2013) *Journal of Experimental Biology*

Determination and profiling of human skin odors using hair samples

Tavares, D.S., Mesquita, P.R.R., Salgado, V.R. (2019) *Molecules*

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

## Author keywords

Body odour

Discriminant analysis

Gas chromatography

Principal component analysis

Shoe odour

VOCs

## Funding details

| Funding sponsor                        | Funding number             | Acronym |
|--|----------------------------|---------|
| Ministry of Higher Education, Malaysia | FRGS/1/2019/STG04/USM/02/5 | MOHE    |

### Funding text

Financial support for this research by MOHE FRGS grant (FRGS/1/2019/STG04/USM/02/5)

ISSN: 2090536X

Source Type: Journal

Original language: English

DOI: 10.1186/s41935-020-00189-0

Document Type: Article

Publisher: Springer

## References (46)

[View in search results format >](#)

All  Export  Print  E-mail  Save to PDF  Create bibliography

- 1 Ara, K., Hama, M., Akiba, S., Koike, K., Okisaka, K., Hagura, T., Kamiya, T., (...), Tomita, F.

Foot odor due to microbial metabolism and its control

(2006) *Canadian Journal of Microbiology*, 52 (4), pp. 357-364. Cited 100 times.

doi: 10.1139/W05-130

[View at Publisher](#)

- 2 Brown, J.S., Prada, P.A., Curran, A.M., Furton, K.G.

Applicability of emanating volatile organic compounds from various forensic specimens for individual differentiation

(2013) *Forensic Science International*, 226 (1-3), pp. 173-182. Cited 24 times.

doi: 10.1016/j.forsciint.2013.01.008

[View at Publisher](#)

- 3 Buttery, R.G., Guadagni, D.G., Ling, L.C., Seifert, R.M., Lipton, W.

Additional Volatile Components of Cabbage, Broccoli, and Cauliflower

(1976) *Journal of Agricultural and Food Chemistry*, 24 (4), pp. 829-832. Cited 175 times.

doi: 10.1021/jf60206a037

[View at Publisher](#)

- 4 Caroprese, A., Gabbanini, S., Beltramini, C., Lucchi, E., Valgimigli, L.

HS-SPME-GC-MS analysis of body odor to test the efficacy of foot deodorant formulations

(2009) *Skin Research and Technology*, 15 (4), pp. 503-510. Cited 33 times.

doi: 10.1111/j.1600-0846.2009.00399.x

[View at Publisher](#)