



Document details

< Back to results | 1 of 1

↗ Export ⬇ Download 🖨 Print ✉ E-mail 📄 Save to PDF ☆ Add to List ⋮ More... >

International journal of pharmaceutical compounding
Volume 24, Issue 2, 1 March 2020, Pages 148-155

Comparison of Disintegrant -addition Methods on the Compounding of Orodispersible Tablets (Article)

Mahesparan, V.A.^a, Bin Abd Razak, F.S.^a, Ming, L.C.^b, Uddin, A.H.^c, Sarker, M.Z.I.^c, Bin, L.K.^d

^aDepartment of Pharmaceutical Technology & Industry, Faculty of Pharmacy, University of Cyberjaya, Cyberjaya, Selangor, Malaysia

^bPAPRSB Institute of Health Sciences, Universiti Brunei Darussalam, Brunei Darussalam

^cFaculty of Pharmacy, International Islamic University Malaysia, Bandar Indera Mahkota, Malaysia

View additional affiliations ▾

Abstract

Orodispersible tablets disintegrate rapidly (within 3 minutes) in the oral cavity and release the medicament before swallowing. The mode of disintegrant addition might affect the properties of orodispersible tablets. The objective of this study was to formulate and evaluate orodispersible tablets by studying different modes of disintegration addition with varying concentrations of disintegrants. The wet granulation method was used to produce the orodispersible tablets. Two methods of disintegration addition were compared (i.e., intragranular, extragranular). Three disintegrants (i.e., cornstarch, sodium starch glycolate, crospovidone) were used at three levels (5%, 10%, and 15%) in the study. The formulations were tested for the powder flowability (angle of repose) and characterized physically (hardness, weight, thickness, friability, disintegration time). The mangosteen pericarp extract was used as a model active pharmaceutical ingredient to be incorporated into the optimum formulation. It was observed that the extragranular method produced granules with better flowability compared to that of the intragranular method. Crospovidone was found as the most efficient disintegrant among the three. The optimum formulation selected was one with the highest concentration of crospovidone (15%), which showed the fastest disintegration time. The mode of disintegrant addition into the orodispersible tablets formulation was found to show a marked difference in the disintegration, as well as other physical characteristics of the orodispersible tablets where the extragranular mode of addition showed better property, which caused the orodispersible tablets to disintegrate the fastest. Copyright© by International Journal of Pharmaceutical Compounding, Inc.

SciVal Topic Prominence ⓘ

Topic: Tablets | Carboxymethylcellulose | Angle of Repose

Prominence percentile: 92.085 ⓘ

Indexed keywords

EMTREE drug terms: excipient

EMTREE medical terms: chemistry drug formulation hardness medicinal chemistry oral drug administration powder procedures solubility tablet

MeSH: Administration, Oral Chemistry, Pharmaceutical Drug Compounding Excipients Hardness Powders Solubility Tablets

Chemicals and CAS Registry Numbers:

Metrics ⓘ View all metrics >

PlumX Metrics ▾
Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

Related documents

Find more related documents in Scopus based on:

Authors > Keywords >

ISSN: 10924221
Source Type: Journal
Original language: English

PubMed ID: 32196477
Document Type: Article
Publisher: NLM (Medline)

© This record is sourced from MEDLINE/PubMed, a database of the U.S. National Library of Medicine

< Back to results | 1 of 1

^ Top of page

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

Customer Service

- Help
- Contact us

ELSEVIER

[Terms and conditions](#) [Privacy policy](#)

Copyright © Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.
We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

