



Document details

[Back to results](#) | 1 of 1

[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)

[Full Text](#) | [View at Publisher](#)

BMC Complementary Medicine and Therapies [Open Access](#)
Volume 20, Issue 1, 18 September 2020, Article number 285

Use of complementary and alternative therapies for the treatment of dysmenorrhea among undergraduate pharmacy students in Malaysia : A cross sectional study (Article) [\(Open Access\)](#)

Abubakar, U., Zulkarnain, A.I., Samri, F., Hisham, S.R., Alias, A., Ishak, M., Sugiman, H., Ghozali, T.

Department of Pharmacy Practice, Kulliyah of Pharmacy, International Islamic University Malaysia, Kuantan, Pahang, 25200, Malaysia

Abstract

[View references \(36\)](#)

Background: Dysmenorrhea is a common problem that affects female students' quality of life and academic activities. Complementary and alternative therapies (CATs) are used for the treatment of dysmenorrhea. This study investigated the practices and perceptions of female undergraduate students with dysmenorrhea towards CATs. **Methods:** This was a cross-sectional study conducted among undergraduate pharmacy students in a public university in Malaysia using a validated and pre-tested self-administered questionnaire. The study was conducted in November and December 2019. The data was analysed using descriptive and inferential statistical tests. **Results:** Of the 318 female undergraduate students invited, 219 completed the questionnaire (response rate: 68.9%) with 52% aged between 21 and 23 years. The prevalence of dysmenorrhea was 72.1%, and the prevalence of ever-use and current use of CATs was 70.3 and 54.4%, respectively. Bed rest (71.5%), hot compress/heating pad (47.5%) and massage (43.0%) were the most common CATs used by the respondents. The most common reasons for using CAT were to reduce the need for analgesics (61.4%), efficacy (37.3%) and recommendation by others (32.9%). About 23 and 9% of the respondents believed that CATs were equally "effective" and "more effective" than analgesics, respectively. Reducing the need for analgesics (AOR: 4.066, 95% CI: 2.136-7.739) and those who agreed that CATs are effective (AOR: 2.701, 95% CI: 1.337-5.457) were independently associated with the current use CATs for the treatment of menstrual pain. **Conclusion:** The prevalence of ever-use and current use of CATs is high among female undergraduate pharmacy students. Bed rest and heat applications are the most common CATs used. Reducing the need for analgesics and efficacy are the factors associated with the current use of CATs. Students should be educated about the safe and effective use of CATs to reduce adverse effects and improve their quality of life. © 2020 The Author(s).

SciVal Topic Prominence [①](#)

Topic: Dysmenorrhea | Acupuncture | Mefenamic Acid

Prominence percentile: 91.720



Author keywords

[Complementary and alternative therapy](#) [Dysmenorrhea](#) [Malaysia](#) [Perception](#) [Pharmacy students](#) [Predictors](#)

Indexed keywords

EMTREE medical terms:

[adult](#) [alternative medicine](#) [Article](#) [bed rest](#) [cross-sectional study](#) [dysmenorrhea](#)
[fatigue](#) [female](#) [headache](#) [human](#) [major clinical study](#) [Malaysia](#) [menstrual cycle](#)
[nausea and vomiting](#) [pain intensity](#) [pharmacy \(shop\)](#) [prevalence](#) [quality of life](#)
[questionnaire](#) [risk factor](#) [undergraduate student](#)

Metrics [②](#) [View all metrics >](#)



PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

Related research data [②](#)

Use of complementary and alternative therapies for the treatment of dysmenorrhea among undergraduate pharmacy students in Malaysia: a cross sectional study

Abubakar, Usman , et al
[figshare](#)

Use of complementary and alternative therapies for the treatment of dysmenorrhea among undergraduate pharmacy students in Malaysia: a cross sectional study

Abubakar, Usman , et al
[figshare](#)

Data linking provided by OpenAIRE's Scholarexplorer

Related documents

Management of primary dysmenorrhea among university students in the south of spain and family influence

Parra-Fernández, M.L. , Onieva-Zafra, M.D. , Abreu-Sánchez, A. (2020) *International Journal of Environmental Research and Public Health*

References (36)

[View in search results format >](#)

All [Export](#)  Print  E-mail  Save to PDF [Create bibliography](#)

- 1 De Sanctis, V., Soliman, A.T., Elsedfy, H., Soliman, N.A., Elalaily, R., El Kholy, M. **Dysmenorrhea in adolescents and young adults: A review in different countries**

(2016) *Acta Biomedica*, 87 (3), pp. 233-246. Cited 40 times.

<http://www.mattioli1885.com/onlinejournals/index.php/actabiomedica/issue/archive>

[View at Publisher](#)

- 2 Iacovides, S., Avidon, I., Baker, F.C. **What we know about primary dysmenorrhea today: A critical review** ([Open Access](#))

(2015) *Human Reproduction Update*, 21 (6), art. no. dmv039, pp. 762-778. Cited 176 times.

<http://humupd.oxfordjournals.org/>

doi: 10.1093/humupd/dmv039

[View at Publisher](#)

[View all related documents based on references](#)

[Find more related documents in Scopus based on:](#)

[Authors >](#) [Keywords >](#)

- 3 Abu Helwa, H.A., Mitaeb, A.A., Al-Hamshri, S., Sweileh, W.M. **Prevalence of dysmenorrhea and predictors of its pain intensity among Palestinian female university students** ([Open Access](#))

(2018) *BMC Women's Health*, 18 (1), art. no. 18. Cited 24 times.

<http://www.biomedcentral.com/bmcwomenshealth/>

doi: 10.1186/s12905-018-0516-1

[View at Publisher](#)

- 4 Ameade, E.P.K., Amalba, A., Mohammed, B.S. **Prevalence of dysmenorrhea among University students in Northern Ghana; its impact and management strategies** ([Open Access](#))

(2018) *BMC Women's Health*, 18 (1), art. no. 39. Cited 11 times.

<http://www.biomedcentral.com/bmcwomenshealth/>

doi: 10.1186/s12905-018-0532-1

[View at Publisher](#)

- 5 Najafi, N., Khalkhali, H., Moghaddam Tabrizi, F., Zarrin, R. **Major dietary patterns in relation to menstrual pain: A nested case control study** ([Open Access](#))

(2018) *BMC Women's Health*, 18 (1), art. no. 69. Cited 7 times.

<http://www.biomedcentral.com/bmcwomenshealth/>

doi: 10.1186/s12905-018-0558-4

[View at Publisher](#)

- 6 Alsaleem, M.A. **Dysmenorrhea, associated symptoms, and management among students at King Khalid University, Saudi Arabia: An exploratory study**
(2018) *J Family Med Prim Care*, 7 (4), p. 769. Cited 5 times.