



# Document details

< Back to results | 1 of 1

↗ Export Download Print E-mail PDF Save to PDF ☆ Add to List More... >

International Medical Journal Malaysia [Open Access](#)  
Volume 19, Issue 2, 1 July 2020, Pages 3-12

## High prevalence of thyroid antibodies in urban population of peninsular Malaysia (Article)

Shahar, M.A.<sup>a</sup> ✉, Omar, A.M.<sup>b</sup>, Wahab, A.B.N.<sup>c</sup>, Sukor, N.<sup>c</sup>, Kamaruddin, N.A.<sup>c</sup> 👤

<sup>a</sup>AVISENA Specialist Hospital, 4, Jalan Ikhtisas, Seksyen 14, Shah Alam, Selangor, 40000, Malaysia

<sup>b</sup>Department of Internal Medicine, Kulliyah of Medicine, International Islamic University Malaysia (IIUM), Kuantan, Pahang, Malaysia

<sup>c</sup>Endocrine Unit, Universiti Kebangsaan Malaysia Medical Centre (UKMMC), Kuala Lumpur, Malaysia

### Abstract

View references (14)

Thyroid antibodies are closely related to autoimmune thyroid disorders. To date, there are no data on the prevalence of these antibodies among the Malaysian population. This study aimed to determine the prevalence of thyroid antibodies; and the factors associated with thyroid antibodies in the Malaysian adult population. MATERIALS AND METHODS: A cross-sectional study was performed in 5 pre-assigned regions in Peninsular Malaysia. Participants' sociodemographic profile and medical history were recorded. Physical examinations were done looking for abnormalities of the thyroid gland and signs of thyroid dysfunctions. Fifteen mls of blood were withdrawn and analysed for thyroid function, anti-thyroperoxidase (anti-TPO) and anti-thyroglobulin (anti-TG) antibodies at a central laboratory. RESULTS: Among the total of 2190 respondents, the overall prevalence of positive anti-TPO and anti-TG antibodies were 12.2% and 12.1%, respectively; mainly found in urban and coastal areas. Only 7% to 9% of those with positive anti-TPO or anti-TG antibodies had either hypo- or hyperthyroidism. The predictors for positive anti-TPO antibody were female [adjusted OR 1.7 (95%CI: 1.2-2.4); p=0.001], Indian [adjusted OR 1.9 (95%CI: 1.1-3.1); p=0.020], and having a goitre [adjusted OR 1.8 (95%CI: 1.2-2.8), p=0.004]. The predictors of positive anti-TG antibody was female [adjusted OR 2.3 (95%CI: 1.6-3.3); p<0.001], and having a goitre [adjusted OR 2.0 (95%CI: 1.3-3.4), p=0.001]. Those living in rural areas had 38% lower risk of having positive anti-TG. CONCLUSIONS: The prevalence of positive anti-TPO and anti-TG antibodies were 12.2% and 12.1%, respectively; were higher among the urban and coastal populations. Those living in rural areas had lower risks of anti-TG antibody. (258 words). © 2020 International Islamic University Malaysia.


### SciVal Topic Prominence ⓘ

Topic: Hypothyroidism | Thyrotropin | Thyroid Function

Prominence percentile: 96.930 ⓘ

### Chemistry database information ⓘ

#### Substances



Metrics ⓘ View all metrics >

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

### Related documents

The association between thyroid autoantibodies in serum and abnormal function and structure of the thyroid

Yan, Y.-R. , Gao, X.-L. , Zeng, J. (2015) *Journal of International Medical Research*

Incidence of thyroid dysfunction in an Iranian adult population: The predictor role of thyroid autoantibodies: Results from a prospective population-based cohort study

Aminorroaya, A. , Meamar, R. , Amini, M. (2017) *European Journal of Medical Research*

The incidence of subclinical thyroid dysfunction and its natural course in the tehranian adults

Assadi, M. , Delshad, H. , Tohidi, M. (2010) *Iranian Journal of Endocrinology and Metabolism*

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

## Author keywords

Anti-TG

Anti-TPO

## Funding details

### Funding text

The study was funded by the Malaysian Endocrine and Metabolic Society (MEMS).

ISSN: 18234631

Source Type: Journal

Original language: English

Document Type: Article

Publisher: International Islamic University Malaysia

## References (14)

[View in search results format >](#)

All  Export  Print  E-mail  Save to PDF  Create bibliography

- 1 Vanderpump, M.P.J., Tunbridge, W.M.G., French, J.M., Appleton, D., Bates, D., Clark, F., Grimley Evans, J., (...), Young, E.T.  
The incidence of thyroid disorders in the community: A twenty-year follow-up of the Wickham Survey  
(1995) *Clinical Endocrinology*, 43 (1), pp. 55-68. Cited 1698 times.  
[http://onlinelibrary.wiley.com.ezlib.iiu.edu.my/journal/10.1111/\(ISSN\)1365-2265](http://onlinelibrary.wiley.com.ezlib.iiu.edu.my/journal/10.1111/(ISSN)1365-2265)  
doi: 10.1111/j.1365-2265.1995.tb01894.x  
[View at Publisher](#)
- 2 Li, Y., Teng, D., Shan, Z., Teng, X., Guan, H., Yu, X., Fan, C., (...), Teng, W.  
Antithyroperoxidase and antithyroglobulin antibodies in a five-year follow-up survey of populations with different iodine intakes ([Open Access](#))  
(2008) *Journal of Clinical Endocrinology and Metabolism*, 93 (5), pp. 1751-1757. Cited 105 times.  
<http://jcem.endojournals.org/cgi/reprint/93/5/1751>  
doi: 10.1210/jc.2007-2368  
[View at Publisher](#)
- 3 Population and Housing Census of Malaysia 2010  
(2011) *Dep Stat Malaysia*, p. 77.
- 4 Mafauzy, M., Wan Mohamad, W.B., Yasmin Anum, M.Y., Musalmah, M., Mustafa, B.E.  
The prevalence of endemic goitre in Kelantan, Malaysia.  
(1993) *The Medical journal of Malaysia*, 48 (1), pp. 64-70. Cited 4 times.
- 5 (2002) *WORLD HEALTH SURVEY C-Appendices (KISH tables*  
Vignettes, Cards
- 6 (2014) *Goitre as a determinant of the prevalence and severity of iodine deficiency disorders in populations*, pp. 14-19. Cited 12 times.

- 7 Kocak, M., Erem, C., Deger, O., Topbas, M., Ersoz, H.O., Can, E.  
Current prevalence of goiter determined by ultrasonography and associated risk factors in a formerly iodine-deficient area of Turkey  
(2014) *Endocrine*, 47 (1), pp. 290-298. Cited 7 times.  
<http://www.springer.com.ezlib.iium.edu.my/humana+press/journal/12020>  
doi: 10.1007/s12020-013-0153-2  
View at Publisher
- 
- 8 Delitala, A.P., Pilia, M.G., Ferrelli, L., Loi, F., Curreli, N., Balaci, L., Schlessinger, D., (...), Cucca, F.  
Prevalence of unknown thyroid disorders in a Sardinian cohort (Open Access)  
(2014) *European Journal of Endocrinology*, 171 (1), pp. 143-149. Cited 25 times.  
<http://www.eje-online.org/content/171/1/143.full.pdf+html>  
doi: 10.1530/EJE-14-0182  
View at Publisher
- 
- 9 Liu, Y., Huang, H., Zeng, J., Sun, C.  
Thyroid volume, goiter prevalence, and selenium levels in an iodine-sufficient area: A cross-sectional study (Open Access)  
(2013) *BMC Public Health*, 13 (1), art. no. 1153. Cited 26 times.  
doi: 10.1186/1471-2458-13-1153  
View at Publisher
- 
- 10 Kasagi, K., Takahashi, N., Inoue, G., Honda, T., Kawachi, Y., Izumi, Y.  
Thyroid function in Japanese adults as assessed by a general health checkup system in relation with thyroid-related antibodies and other clinical parameters  
(2009) *Thyroid*, 19 (9), pp. 937-944. Cited 56 times.  
doi: 10.1089/thy.2009.0205  
View at Publisher
- 
- 11 Bjoro, T., Holmen, J., Kruger, O., Midthjell, K., Hunstad, K., Schreiner, T., Sandnes, L., (...), Brochmann, H.  
Prevalence of thyroid disease, thyroid dysfunction and thyroid peroxidase antibodies in a large, unselected population. The health study of Nord-Trøndelag (HUNT) (Open Access)  
(2000) *European Journal of Endocrinology*, 143 (5), pp. 639-647. Cited 333 times.  
<http://www.eje-online.org/>  
doi: 10.1530/eje.0.1430639  
View at Publisher
- 
- 12 Hollowell, J.G., Staehling, N.W., Dana Flanders, W., Harry Hannon, W., Gunter, E.W., Spencer, C.A., Braverman, L.E.  
Serum TSH, T<sub>4</sub>, and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III) (Open Access)  
(2002) *Journal of Clinical Endocrinology and Metabolism*, 87 (2), pp. 489-499. Cited 2538 times.  
<http://jcem.endojournals.org>  
doi: 10.1210/jcem.87.2.8182  
View at Publisher
-

□ 13 Yan, Y.-R., Gao, X.-L., Zeng, J., Liu, Y., Lv, Q.-G., Jiang, J., Huang, H., (...), Tong, N.-W.

The association between thyroid autoantibodies in serum and abnormal function and structure of the thyroid ([Open Access](#))

(2015) *Journal of International Medical Research*, 43 (3), pp. 412-423. Cited 6 times.

<http://imr.sagepub.com/content/by/year>

doi: 10.1177/0300060514562487

[View at Publisher](#)

□ 14 Selamat, R., Wan Mohamud, W.N., Zainuddin, A.A., Rahim, N.S.C.A., Ghaffar, S.A., Aris, T.

Iodine deficiency status and iodised salt consumption in Malaysia: Findings from a national iodine deficiency disorders survey

(2010) *Asia Pacific Journal of Clinical Nutrition*, 19 (4), pp. 578-585. Cited 13 times.

[http://apjcn.nhri.org.tw/server/APJCN/Volume19/vol19.4/Finished/18\\_1782\\_578-585.pdf](http://apjcn.nhri.org.tw/server/APJCN/Volume19/vol19.4/Finished/18_1782_578-585.pdf)

📍 Shahr, M.A.; AVISENA Specialist Hospital, 4, Jalan Ikhtisas, Seksyen 14, Shah Alam, Selangor, Malaysia;  
email:arifshahr2013@gmail.com

© Copyright 2020 Elsevier B.V., All rights reserved.

< Back to results | 1 of 1

^ Top of page

## About Scopus

[What is Scopus](#)  
[Content coverage](#)  
[Scopus blog](#)  
[Scopus API](#)  
[Privacy matters](#)

## Language

[日本語に切り替える](#)  
[切换到简体中文](#)  
[切换到繁體中文](#)  
[Русский язык](#)

## Customer Service

[Help](#)  
[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

 RELX