

Brought to you by [INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA](#)



Scopus



[Back](#)

The Feasibility of Using Human Primary Chondrocytes Derived from Osteoarthritic Patients Overexpressed with SOX9 Seeded on PLGA-Fibrin Hybrid Scaffolds for Cartilage Engineering

[Malaysian Journal of Fundamental and Applied Sciences](#) • Article • Open Access • 2022 •

DOI: 10.11113/mjfas.v18n3.2344

[Sukri, Norhamiza Mohamad](#)^a; [Shaban, Munirah](#)^b ; [Radzi, Muhammad Aazamuddin Ahmad](#)^a; [Rahman, Rozlin Abdul](#)^b; [Zulkifly, Ahmad Hafiz](#)^c

^a Department of Biomedical Science, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, Jalan Sultan Haji Ahmad Shah, Bandar Indera Mahkota, Pahang, Kuantan, 25200, Malaysia

[Show all information](#)

2 33th percentile

Citations

0.21

FWCI

[View PDF](#)

[Full text](#)

[Export](#)

[Save to list](#)

[Document](#)

[Impact](#)

[Cited by \(2\)](#)

[References \(83\)](#)

[Similar documents](#)

Abstract

This study aimed to find an optimal formulation to form 3D hyaline-like cartilage substitutes using the tissue engineering triads. The primary cells taken from osteoarthritic patients were overexpressed with transcriptional factor SRY (Sex Determining Region Y)-Box 9 (SOX9) using Lipofectamine 2000™ through a non-viral transfection method. The transfected and non-transfected cells were seeded on poly(lactic-co-glycolic acid) (PLGA) based scaffolds with and without fibrin. The arrangement resulted in four experimental groups. The 3D 'cells-scaffolds' tissue constructs were cultured for three weeks and implanted ectopically in nude mice for four weeks. The evaluations include macroscopic and microscopic study, gene expression analyses, and sulfated glycosaminoglycan (sGAG) assay, focusing on the cartilage properties. A biomechanical evaluation was performed only on post-implanted constructs. All in vitro, two- and four-week post-implanted constructs exhibited firm and smooth hyaline-like cartilage appearance. In vitro constructs showed sparse cells distribution with minimal cartilaginous tissue formation. However, a high density, lacunae-encapsulated chondrocytes embedded within the basophilic ground substances was observed in all post-implanted constructs. It is supported by positive-brownish precipitation immunolocalisation against collagen type II. Besides, molecular analysis showed that COL2A1 and other cartilaginous markers were also expressed. Increased sGAG content and compressive strain could be observed in vitro and in vivo. Although quantitatively, no significant statistical differences were found between the four groups, the qualitative results indicated that SOX9-overexpressed cells, PLGA, and fibrin combination guides hyaline-like cartilage formation better than other groups. Hence, the combination may be studied in a big animal model to develop its potential for future clinical application. ©Copyright Sukri et al.

Author keywords

Articular cartilage; chondrocytes; ectopic implantation model; fibrin; gene transfer; osteoarthritis; Overexpression; PLGA; SOX9

Funding details

Details about financial support for research, including funding sources and grant numbers as provided in academic publications.

Funding sponsor	Funding number	Acronym
IIUM Tissue Engineering and Regenerative Medicine Research Team		
University of Chicago Medical Center		

Funding sponsor	Funding number	Acronym
Ministry of Higher Education, Malaysia See opportunities by MOHE ↗		MOHE
International Islamic University Malaysia See opportunities by IIUM ↗		IIUM

Funding text

The authors thank Malaysia Research Assessment Instrument (MyRA) Research Grant Scheme (MIRGS13-01-002-0003), Ministry of Higher Education (MOHE) Malaysia, and International Islamic University Malaysia (IIUM) for research funding and facilities, IIUM Tissue Engineering and Regenerative Medicine Research Team, and Assoc. Professor Dr Tong-Chuan He from the University of Chicago Medical Center, USA, for the pcDNA3-SOX9 plasmid gift.

Corresponding authors

Corresponding author	M. Shaban
Affiliation	Physical Rehabilitation Sciences, Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, Jalan Sultan Haji Ahmad Shah, Bandar Indera Mahkota, Pahang, Kuantan, 25200, Malaysia
Email address	munirahshaban@iium.edu.my

© Copyright 2022 Elsevier B.V., All rights reserved.

Abstract

Author keywords

Funding details

Corresponding authors