English Products

Web of Science™ Smart Search Advanced Search Research Assistant Noor Afiqah Musa

MENU

Smart Search › Results for Automated fract… › … › Refine results for Lifetime tr… ›

Lifetime trauma, symptoms of psychological disturbance, and suicidal idea…

Full text at publisher View Full Text on ProQuest

Export Add To Marked List

1 of 101

Lifetime trauma, symptoms of psychological disturbance, and suicidal ideation among university students in Malaysia

By Ghazali, SR (Ghazali, Siti Raudzah) [1] ; Chen, YY (Chen, Yoke Yong) [1] ; Mohamad, M (Mohamad, Mardiana) [2] ; Yein, LP (Yein, Lee Ping) [3]

Are you this author?

Citation Network

In Web of Science Core Collection

6

Citations

Create citation alert

8

; Razali, ZA (Razali, Zul Azlin) [4] ; Pauzi, N (Pauzi, Noraskin) [1] ; Majani, AF (Majani, Alia Fadaliana) [1]

View Web of Science ResearcherID and ORCID (provided by Clarivate)

Source CURRENT PSYCHOLOGY

← View Journal Impact

Volume: 42 Issue: 25 Page: 22127-22136

DOI: 10.1007/s12144-022-03297-3

Published SEP 2023

Early Access JUN 2022

Indexed 2022-06-24

Document Type Article

Jump to ↓ Enriched Cited References

Abstract In many countries, mental disorders are high among university students. We conducted a cross-sectional research design among 1767 university students (71.2% were females and the remaining were males) who attended public universities in Sarawak. Questionnaires related to a history of lifetime trauma, posttraumatic stress disorder (PTSD), and depressive symptoms together with suicide behavior were surveyed. Results showed that at least 87% of students had lifetime trauma once, 9.6% exhibited PTSD symptoms, 15.5% exhibited depressive symptoms, and 12.9% of participants reported having suicidal behavior. There were significantly more females who reported having suicidal behaviors than

Times Cited in All Databases

+ See more times cited

64

Cited References

→ View Related Records

How does this document's citation performance compare to peers?

← Open comparison metrics panel

Data is from InCites Benchmarking & Analytics

Citing items by classification

males. Ethnicity was significantly associated with PTSD symptoms. PTSD and depressive symptoms were significant predictors of suicidal behavior. Other significant predictors for suicide behavior among those who were traumatized were near drowning, sexual assault, losing a family member, neglect, and bullying. This study provides a significant understanding of the link between PTSD, depressive symptoms, and suicidal behavior among Malaysian university students.

Keywords

Author Keywords: PTSD; Depression; University students; Trauma

Keywords Plus: POSTTRAUMATIC-STRESS-DISORDER; MEDIATING ROLE; DEPRESSIVE SYMPTOMS; INTERNET ADDICTION; TREATMENT-SEEKING; CHILDHOOD TRAUMA; MENTAL-DISORDERS; SEX-DIFFERENCES; PREVALENCE; RISK

Author Information

Corresponding Address: Ghazali, Siti Raudzah (corresponding author)

Univ Malaysia Sarawak, Fac Med & Hlth Sci, Dept Psychol Med, Sarawak 94300, Malaysia

E-mail Addresses :

gsraudzah@unimas.my

Addresses :


[1] Univ Malaysia Sarawak, Fac Med & Hlth Sci, Dept Psychol Med, Sarawak 94300, Malaysia

[2] Int Islamic Univ Malaysia, Fac Islamic Revealed Knowledge & Human Sci, Dept Psychol, Gombak, Selangor, Malaysia

[3] Univ Malaya, Fac Med, UMeHlth Unit, Kuala Lumpur, Malaysia

[4] Univ Sains Islam Malaysia, Fac Med & Hlth Sci, Nilai, Negeri Sembilan, Malaysia

Breakdown of how this article has been mentioned, based on available citation context data and snippets from 3 citing item(s).



You may also like...

Eagle, DE; Hybels, CF; Proeschold-Bell, RJ; Perceived social support, received social support, and depression among clergy JOURNAL OF SOCIAL AND PERSONAL RELATIONSHIPS

Dong, X; Su, F; Wang, ZH; et al.